

Correction to: “Prediction of the Clinical Risk of Drug-Induced Cholestatic Liver Injury Using an In Vitro Sandwich Cultured Hepatocyte Assay”

In the above article [Susukida T, Sekine S, Nozaki M, Tokizono M, and Ito K (2015) *Drug Metab Dispos* **43**: 1760-1768], the authors would like to make the below corrections.

In Table 2, composition of the 1×BA mixture standard was incorrectly calculated as 1.5×BA mixture standard. A corrected Table 2 is provided below.

The Full Text is corrected with the issue of this erratum.

The authors apologize for any inconvenience this may have caused.

TABLE II
Composition of the 1 × BA mixture standard.

Concentrations of each BA are set based on the standard BA constituents of human serum (Scherer et al., 2009).

Bile acid	1 × Standard concentration (μM)
Cholic acid	0.20
Chenodeoxy cholic acid	0.34
Glycochenodeoxycholic acid	1.71
Deoxycholic acid	0.73
Lithocholic acid	0.030
Ursodeoxycholic acid	0.11
Glycocholic acid	0.41
Glycodeoxycholic acid	0.38
Taurocholic acid	0.048
Taurochenodeoxycholic acid	0.21
Taurolithocholic acid	0.087
Tauroursodeoxycholic acid	0.29