

## AMINO ACID CONJUGATES: METABOLITES OF 2-PROPYLPENTANOIC ACID (VALPROIC ACID) IN EPILEPTIC PATIENTS

V. S. GOPAUL,<sup>1</sup> W. TANG,<sup>2</sup> K. FARRELL, AND F. S. ABBOTT

*Faculties of Pharmaceutical Sciences (V.S.G., W.T., F.S.A.) and Medicine (K.F.), University of British Columbia, Vancouver, Canada*

(Received September 3, 2002; accepted October 7, 2002)

This article is available online at <http://dmd.aspetjournals.org>

### ABSTRACT:

In this study, spectroscopic and chromatographic evidence is presented for the identification and characterization of the metabolites, valproyl glutamate (2-propylpentanoyl glutamate, VPA-GLU) and valproyl glutamine (2-propylpentanoyl glutamine, VPA-GLN) in the urine, serum, and cerebrospinal fluid (CSF) of patients on valproic acid (VPA) therapy. Moreover, the identification of valproyl glycine (2-propylpentanoyl glycine, VPA-GLY) in the serum and urine of patients on VPA, albeit in trace concentrations, is also reported here. The three amino acid conjugates excreted in urine accounted for about 1% of the VPA dose in four patients who were on VPA therapy chronically and had reached steady state. VPA-

GLU was quantitatively the most prominent metabolite (0.66–13.1  $\mu\text{g}/\text{mg}$  creatinine) compared with VPA-GLN (0.78–9.93  $\mu\text{g}/\text{mg}$  creatinine) and VPA-GLY (trace–1.0  $\mu\text{g}/\text{mg}$  creatinine) in overnight urine samples of all patients studied ( $n = 29$ ). The relatively low serum concentrations of the three amino acid conjugates of VPA in six patients suggest that the metabolites are readily excreted once formed. In contrast, whereas VPA GLY was absent in the CSF of one patient on VPA, the concentrations of VPA-GLU and VPA-GLN in this CSF sample were 9 and 5 times, respectively, their corresponding serum concentrations.

The antiepileptic drug, valproic acid (VPA<sup>3</sup>), is primarily used for the management of generalized and absence seizures in children. Concerns over its teratogenicity and hepatotoxicity side effect have encouraged the need to develop analogs of VPA devoid of the adverse effects. However, this task is complicated by the fact that the mechanism of action of the drug itself remains to be understood.

A large body of evidence has emerged to link the hepatotoxicity of VPA to its metabolism and has been reviewed over the years (Eadie

et al., 1988; Abbott and Anari, 1999; Radatz and Nau, 1999). Moreover, a review of mass-balance studies indicated that the total recovery of a VPA dose cannot be completely accounted for in humans (Baillie and Sheffels, 1995), suggesting the existence of yet unknown metabolites. For all these reasons, despite the extensive work that has been conducted on the metabolism of VPA, the subject remains an area that warrants further investigation.

To date, the role of the amino acid conjugation pathway in the biotransformation of VPA, particularly in humans, has remained relatively unexplored. 2-Propylpentanoyl glycine (VPA-GLY) is the only amino acid conjugate of the drug that has been detected and studied in rats (Granneman et al., 1984a,b). The role of amino acid conjugation of VPA in human is significant owing to the potential for CoA consumption. The depletion of mitochondrial CoA with VPA therapy is one of several hypotheses put forth to explain VPA hepatotoxicity (Turnbull et al., 1983; Harris et al., 1991). Furthermore, the conjugation of VPA with amino acids known to be neurotransmitters involved in epilepsy also need to be explored as a potential to further understand the mechanism of action of VPA.

Although in general, bile acids form conjugates with glycine and taurine by the action of microsomal enzymes, amino acid conjugation of xenobiotics is believed to be a mitochondrial reaction occurring primarily in the liver or the kidney (Williams, 1989). Conjugation of organic acids is initiated by the obligatory activation of the acid to a CoA ester followed by the acyl transfer to an amino acid residue with concurrent release of CoA. The last step is catalyzed by *N*-acyl transferase specific for each amino acid and defines the structural requirement for amino acid conjugation of xenobiotics (Hutt and Caldwell, 1990). Glycine-*N*-acyltransferase (Tishler and Goldman, 1970; Mawal et al., 1997) and glutamine-*N*-phenylacetyltransferase

This work was supported by a program grant from the Medical Research Council of Canada and a program grant from the Medical Research Council of Canada, and was part of the doctoral dissertation of Gopaul (1998). A preliminary account of these studies was presented at the 6th European ISSX meeting, 1997 June 30–July 3, Gothenburg, Sweden.

<sup>1</sup> Current address: Department of Drug Metabolism, Johnson & Johnson Pharmaceutical Research and Development, Spring House, PA, 19477

<sup>2</sup> Current address: Department of Drug Metabolism, Merck & Co., Rahway, NJ 07065-0900.

<sup>3</sup> Abbreviations used are: VPA, valproic acid; VPA-GLY, valproyl glycine (2-propylpentanoyl glycine); LC-MS/MS, liquid-chromatography tandem mass-spectrometry; GC-MS, gas-chromatography mass-spectrometry; CSF, cerebrospinal fluid; PFBBr, 1-bromo-2,3,4,5,6-pentafluorotoluene; DIPEA, diisopropylethylamine; TFA, trifluoroacetic acid; IS, internal standard; GC-MS NICI, gas chromatography in negative ion chemical ionization; MS, mass spectrometer; HPLC, high performance liquid chromatography; MRM, multiple reaction monitoring; CID, collision-induced dissociation; FVPA-GLN, 2-fluoro valproyl glutamine; VPA-GLN, valproyl glutamine (2-propylpentanoyl glutamine); VPA-GLU, valproyl glutamate (2-propylpentanoyl glutamate); VPA-ASP, valproyl aspartate; PFB, pentafluorobenzyl.

**Address correspondence to:** Frank Abbott, Ph.D., Faculty of Pharmaceutical Sciences, 2146 East Mall, University of British Columbia, Vancouver, BC, Canada V6T 1Z3. E-mail: [fabbott@interchange.UBC.ca](mailto:fabbott@interchange.UBC.ca)

(Webster et al., 1976) have been isolated from human liver mitochondria.

In this study, we looked for evidence of the conjugation of VPA with the amino acid neurotransmitters glycine, glutamine, glutamic acid, and aspartic acid, as well as non-neurotransmitters, taurine and alanine in the biological fluids of humans by liquid-chromatography tandem mass-spectrometry (LC-MS/MS) and gas-chromatography mass-spectrometry (GC-MS). We describe here the identification and profiling of amino acid conjugates of VPA in the urine, serum, and CSF of patients on VPA therapy using LC-MS/MS and GC-MS.

### Materials and Methods

**Materials.** 1-Bromo-2,3,4,5,6-pentafluorotoluene (PFBB), diisopropylethylamine (DIPEA), VPA, L-aspartic acid, L-glutamine, L-glutamic acid, and L-glycine were purchased from Aldrich Chemical Co. (Milwaukee, WI); distilled in glass grade solvents (ethyl acetate, methanol, acetonitrile) were purchased from BDH Chemicals (Toronto, ON) or Caledon Laboratories Ltd. (Edmonton, AB). Trifluoroacetic acid (TFA) was purchased from Sigma-Aldrich (St. Louis, MO). 2-Fluoro-2-propylpentanoic acid was synthesized in our laboratory (Tang et al., 1997) and used as an internal standard (IS) in LC-MS/MS assays.

Acrodisc LC 13 PVDF (0.2  $\mu$ m) syringe filters were purchased from German Sciences Co. (Ann Arbor, MI); Bond-Elut C<sub>2</sub> solid phase extraction cartridges (500 mg/3 ml) were purchased from Varian Sample Preparation Products (Harbor City, CA); polyethylene tubing PE-10 was purchased from Clay Adams (Parsippany, NJ).

**Instrumentation and Analytical Conditions.** <sup>1</sup>H NMR spectra were obtained on a Bruker WH-200 spectrometer (Bruker, Newark, DE) or a Bruker WH-400 spectrometer (Department of Chemistry, University of British Columbia).

Gas chromatography in negative ion chemical ionization (GC-MS NICI) was performed on a Hewlett Packard (Avondale, PA) 5989 mass spectrometer (MS) engine coupled to a 5890 series II gas chromatograph (GC) fitted with a J & W Scientific (Folsom, CA) DB1701 column, (30 m  $\times$  0.32 mm, 0.25  $\mu$ m). The oven temperature was programmed as follows: 100 to 200°C at 20°C/min, held for 5 min; 150 to 300°C at 15°C/min and held for 5 min. Other GC-MS conditions were ion source at 200°C, detector temperature at 280°C, injection port temperature at 240°C, and He was set at 10 psi; the reagent gas, methane, was maintained at 1 torr; the emission current was at 300  $\mu$ A; and the ionization energy was 120 eV. For purposes of identification and characterization of the amino acid conjugates of VPA, both the scan and selected ion monitoring modes were used.

LC-MS/MS experiments were performed on a Micromass Quattro triple quadrupole mass spectrometer (Micromass, Montreal, QC) interfaced to a Hewlett Packard 1090 II liquid chromatograph. Chromatography was conducted on a Phenomenex (Torrance, CA) C<sub>8</sub> column, (100 mm  $\times$  2.1 mm, 5  $\mu$ m). Three HPLC methods were used in this study.

**Method 1.** Mobile phase A consisted of methanol/water (43:57, 0.05% TFA), and mobile phase B consisted of methanol. At time 0, mobile phase A was pumped isocratically for 30 min. Mobile phase B was gradually increased from 0 to 100% at time 30.1 min and held for 5 min, followed by a sharp gradient increase of mobile phase A to 100% at 35.1 min. This method was used initially for the identification of the amino acid conjugates of VPA in human urine. Flow rate was at 0.1 ml/min.

**Method 2.** Mobile phase A consisted of acetonitrile/water (30:70, 0.05% TFA) and mobile phase B consisted of acetonitrile. Mobile phase A was pumped for 25 min followed by a gradient increase of mobile phase B to 100% at 26 min, held for 5 min and reverted to 100% A at 30.1 min. Flow rate was at 0.1 ml/min. This method was used to detect the amino acid conjugates of VPA in urine where a longer run time program was required to separate the peaks of interest from interfering chromatographic peaks.

**Method 3.** Mobile phase A consisted of acetonitrile/water (40:60, 0.05% TFA). Mobile phase B consisted of a gradient program that allowed 100% of mobile phase A to be pumped for 14 min at a flow rate of 0.1 ml/min. Mobile phase B was then increased to 100% at a flow rate of 0.2 ml/min at 15 min and held for 5 min at 0.2 ml/min. Subsequently, mobile phase A was increased to

100% at time 20 min and at 0.1 ml/min. This method was used for the profiling of amino acid conjugates in human serum and CSF samples.

The ionization energy for LC-MS/MS experiments was in positive electrospray (ES<sup>+</sup>). The HPLC eluent was introduced to the stainless steel capillary sprayer held at 3.5 kV. The mass spectrometer was operated in either precursor ion scanning, product ion scanning or multiple reaction monitoring (MRM) mode. Multipliers 1 and 2 were set at 650 V for all three experiments; cone voltage was set at 30 V with skimmer offset by 5 V. The low and high mass resolution was set at 12.5 for both precursor ion scanning and product ion scanning and at 5 for MRM. The product ion scanning experiment determined the product ions of MH<sup>+</sup>, and the precursor ion scanning confirmed the pseudomolecular ion or the protonated molecular ion (MH<sup>+</sup>) of the metabolite of interest. Argon, the target gas, was set at a pressure of  $3.0 \times 10^{-4}$  mbar for collision-induced dissociation (CID) and with a collision energy of 40 eV. During MRM, the pseudomolecular ion for each analyte was selected in the first quadrupole and subjected to CID with the target gas and collision energy set at 50 eV. Specific fragment ions were then selected by the third quadrupole. For all the experiments, the source temperature was at 140°C. Data analyses were processed using Mass Lynx software (Micromass).

**Human Studies.** Overnight urine samples ( $n = 29$ ), 24-h urine samples ( $n = 4$ ), serum samples ( $n = 6$ ), and one CSF sample were obtained from epileptic and pediatric patients who have been on VPA therapy for more than 6 months and have reached steady state. The patients were attending the Seizure Clinic at Vancouver Children's Hospital, British Columbia, Canada. The average dose of VPA received by each patient was  $23.3 \pm 13.2$  mg/kg, the average age of the patients was  $12.6 \pm 5.9$  years. Control urine and serum samples were obtained from healthy volunteers ( $n = 4$ ). Control CSF were obtained from nonepileptic subjects ( $n = 10$ ), who were either suffering from multiple sclerosis and attending the University of British Columbia Hospital, Canada or were healthy volunteers. Approval for all human studies was issued by the U.B.C. Human Ethics Committee (U.B.C. C95-0412).

**Chemical Synthesis of Amino Acid Conjugates of VPA.** The amino acid conjugates of VPA were synthesized according to the procedure described for 2-fluoro VPA-GLN (FVPA-GLN) (Tang et al., 1997). VPA was converted to the corresponding *N*-hydroxysuccinimide ester in the presence of dicyclohexylcarbodiimide followed by coupling with L-glutamine, L-glycine, L-glutamic acid, or L-aspartic acid to form 2-propylpentanoyl glutamine (VPA-GLN), VPA-GLY, 2-propylpentanoyl glutamate (VPA-GLU), or valproyl aspartate (VPA-ASP), respectively.

**VPA-GLN.** LC-MS/MS spectrum:  $m/z$  (%) 273 (MH<sup>+</sup>, 100), 57 (75), 99 (35), 127 (25), 147 (20); <sup>1</sup>H NMR (200 MHz):  $\delta$  0.90 (t, 6H, 2xCH<sub>3</sub>), 1.15–2.50 (m, 12H, 6xCH<sub>2</sub>), 4.40 (dt, 1H, NHCH); GC-MS NICI spectrum of the PFB derivative:  $m/z$  (%) 271 ([M-181]<sup>-</sup>, 100).

**VPA-GLU.** LC-MS/MS spectrum:  $m/z$  (%) 274 (MH<sup>+</sup>, 100), 57 (100), 99 (40), 127 (35), 148 (20); <sup>1</sup>H NMR (200 MHz):  $\delta$  0.90 (2t, 6H, 2xCH<sub>3</sub>), 1.15–2.50 (m, 12H, 6xCH<sub>2</sub>), 4.45 (dd, 1H, NHCH); GC-MS spectrum of the di-PFB derivative:  $m/z$  (%) 452 ([M-181]<sup>-</sup>, 100)

**VPA-GLY.** LC-MS/MS spectrum:  $m/z$  (%) 202 (MH<sup>+</sup>, 100), 57 (70), 99 (15), 127 (15); <sup>1</sup>H NMR (400 MHz):  $\delta$  0.90 (t, 6H, 2xCH<sub>3</sub>), 1.10–1.40 (m, 8H, 4xCH<sub>2</sub>), 2.0–2.1 (m, 1H, CH<sub>2</sub>CHCH<sub>2</sub>), 4.1 (d, 2H, NHCH<sub>2</sub>); GC-MS NICI spectrum of the PFB derivative:  $m/z$  (%) 200 ([M-181]<sup>-</sup>, 100)

**VPA-ASP.** LC-MS/MS spectrum:  $m/z$  (%) 260 (MH<sup>+</sup>, 100), 57 (100), 99 (40), 127 (35), 134 (15); <sup>1</sup>H NMR (200 MHz):  $\delta$  0.90 (2t, 6H, 2xCH<sub>3</sub>), 1.10–1.40 (m, 8H, 4xCH<sub>2</sub>), 2.0–2.1 (CH<sub>2</sub>CHCH<sub>2</sub>), 4.80 (dd, 2H, NHCH<sub>2</sub>); GC-MS NICI spectrum of the di-PFB derivative:  $m/z$  (%) 438 ([M-181]<sup>-</sup>, 100)

**Derivatization of Amino Acid Conjugates of VPA for NICI Analysis.** Ten microliters of 40% PFBB in ethyl acetate and 10  $\mu$ l of DIPEA were added




Fig. 1. Structures of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP showing three proposed MS/MS fragmentation patterns commonly observed.




FIG. 2. LC-MS/MS (electrospray ionization) of  $MH^+$  of a synthetic standard of (A) VPA-GLU, (B) VPA-GLN, (C) VPA-GLY, and (D) VPA-ASP

to 10  $\mu$ l of a solution (0.01 mg/ml) of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP in ethyl acetate and allowed to react at 50°C for 1 h.

**Preparation of Urine Samples for the Identification and Profiling of Amino Acid Conjugates of VPA by Solid Phase Extraction.** One milliliter

of urine sample was adjusted to pH 3 with HCl and applied to a C<sub>2</sub> Bond-Elut cartridge that was preconditioned with water and methanol. The cartridges were washed once with water/methanol (95:5) and water. The amino acid conjugates of VPA were eluted with 6 ml of methanol. The organic layers were

dried and reconstituted with water. Ten microliters of the reconstituted solvent were then injected into the LC-MS/MS system and analyzed using LC-MS/MS methods 1 and 2.

For the identification and characterization experiments, the chromatographic and mass spectral characteristics of the conjugates from biological fluids were compared with those of the synthesized compounds using precursor ion scanning, product ion scanning or MRM. MS/MS dwell times were set at 1 s:100 amu. In MRM mode, the following transitions were monitored: VPA-GLU:  $m/z$  274  $\rightarrow$   $m/z$  57,  $m/z$  274  $\rightarrow$  99,  $m/z$  274  $\rightarrow$  127, and  $m/z$  274  $\rightarrow$  148; VPA-GLN:  $m/z$  273  $\rightarrow$  57,  $m/z$  273  $\rightarrow$  99,  $m/z$  273  $\rightarrow$  127,  $m/z$  273  $\rightarrow$  129,  $m/z$  273  $\rightarrow$  147; VPA-GLY:  $m/z$  202  $\rightarrow$  57,  $m/z$  202  $\rightarrow$  99,  $m/z$  202  $\rightarrow$  127; VPA-ASP:  $m/z$  260  $\rightarrow$  57,  $m/z$  260  $\rightarrow$  99,  $m/z$  260  $\rightarrow$  127,  $m/z$  260  $\rightarrow$  134.

For the quantitation of VPA-GLU, VPA-GLN, and VPA-GLY in urine, 1 ml of each sample was spiked with 25  $\mu$ l of 0.08 mg/ml of FVPA-GLN (IS) and VPA-ASP (IS). The samples were prepared and analyzed with a set of calibration standards ranging from 0.10 to 5.0  $\mu$ g/ml. Calibration standards and all samples of patients were extracted as described above. The LC-MS/MS MRM transitions monitored were as follows:  $m/z$  291  $\rightarrow$  130 for FVPA-GLN;  $m/z$  274  $\rightarrow$  148 for VPA-GLU;  $m/z$  273  $\rightarrow$  147 for VPA-GLN,  $m/z$  260  $\rightarrow$  127 for VPA-ASP (IS), and  $m/z$  200  $\rightarrow$  57 for VPA-GLY.

The assay for the simultaneous analysis of amino acid conjugates of VPA by LC-MS/MS was linear over a range of 0.1 to 5.0  $\mu$ g/ml for all three conjugates. The coefficients of determination ( $r^2$ ) of the calibration curves were 0.99 or better.

The interassay variation was assessed from the slopes of five calibration curves run consecutively and was less than 7% for each identified metabolite. The intra-assay variation was assessed by three replicate analyses of two samples. At 0.12  $\mu$ g/ml, the intra-assay variation was 11.4, 28.8, 18.5% for VPA-GLU, VPA-GLN, and VPA-GLY, respectively. At 1.0  $\mu$ g/ml, the intra-assay variation was less than 10% for all three metabolites. The larger variation observed at the lower limit of quantitation was acceptable in view of the fact that the metabolites were detected at concentrations well above the lowest standard (0.1  $\mu$ g/ml). At both concentrations, each metabolite was measured within 83 to 120% of their expected values.

The concentration of each amino acid conjugate of VPA measured in overnight urine samples was normalized to creatinine. The % recovery of VPA as amino acid conjugates of VPA was calculated for 24-h urine samples ( $n = 4$ ). Creatinine was measured in urine according to the routine Jaffe method (1886).

**Preparation of Serum Samples for the Identification and Quantitation of the Amino Acid Conjugates of VPA.** The amino acid conjugates of VPA were detected from the serum samples of human by liquid-liquid extraction. Two milliliters of each specimen and control serum was adjusted to pH 3 with HCl and extracted with 5 ml of ethyl acetate, twice. The organic layers were separated and evaporated, and the residue was reconstituted in 200  $\mu$ l of distilled water and filtered through a filter membrane (0.2  $\mu$ m, Acrodisc LC 13 PVDF; Gelman Sciences, Ann Arbor, MI). The filtrate was then analyzed by LC-MS/MS method 3 using the MRM conditions described for the analysis of the urine samples.

The quantizations of VPA-GLU, VPA-GLN, and VPA-GLY in human serum samples were conducted separately from each other. In each case, 1 milliliter of serum sample, calibration standards (5–100 ng/ml), and control serum samples were spiked with 12.5  $\mu$ l of FVPA-GLN (0.08  $\mu$ g/ml) and extracted by the liquid-liquid extraction procedure described above. The extracts were reconstituted in distilled water and analyzed by LC-MS/MS method 3 with MRM set for  $m/z$  291  $\rightarrow$  130,  $m/z$  274  $\rightarrow$  148,  $m/z$  273  $\rightarrow$  147,  $m/z$  202  $\rightarrow$  57 for FVPA-GLN, VPA-GLU, VPA-GLN, and VPA-GLY, respectively.

TABLE 1

GC-MS retention times ( $t_R$ ) of the PFB or di-PFB derivatives of the amino acid conjugates of VPA

PFB or di-PFB* derivatives of:	GC-MS ( $t_R = \text{min}$ )
VPA GLN	15.30
VPA GLU*	16.58
VPA GLY	8.99
VPA ASP*	15.71
FVPA GLN	14.77

TABLE 2

HPLC retention times ( $t_R$ ) of amino acid conjugates of VPA with different mobile phases.

Mobile phase for LC-MS/MS 1: methanol/water (43:57, 0.05% TFA). Mobile phase for LC-MS/MS 2: acetonitrile/water (30:70, 0.05% TFA). Mobile phase for LC-MS/MS 3: acetonitrile/water (40:60, 0.05% TFA).

	LC-MS/MS Method 1 ( $t_R = \text{min}$ )	LC-MS/MS Method 2 ( $t_R = \text{min}$ )	LC-MS/MS Method 3 ( $t_R = \text{min}$ )
VPA-GLN	22.97	14.70	7.86
VPA-GLU	30.70	23.20	10.53
VPA-GLY	30.30	24.0	11.99
VPA-ASP		23.99	9.93
FVPA-GLN			9.33

The assay was linear over a concentration range of 5 to 100 ng/ml with  $r^2$  of 0.99 or better for the calibration curves of each VPA-GLU, VPA-GLN, and VPA-GLY.

**Preparation of CSF Samples for the Identification, and Quantitation of Amino Acid Conjugates of VPA in Human CSF.** A CSF sample (100  $\mu$ l) of a patient on VPA, a control human CSF sample (100  $\mu$ l) spiked with 0.025  $\mu$ g/ml of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP, and 10 control CSF samples (100  $\mu$ l per sample) were filtered through a membrane filter as described for the serum samples and analyzed under LC-MS/MS method 3 for VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP as performed for the urine samples.

A CSF sample (100  $\mu$ l) of a patient on VPA, a control CSF sample spiked with VPA-GLU (0.025  $\mu$ g/ml), and a control CSF sample were subjected to solid phase extraction as described for the urine samples. The residues were dissolved in 500  $\mu$ l of ethyl acetate and derivatized with 10  $\mu$ l of 40% PFBBR and 10  $\mu$ l of DIPEA at 50°C, and 1  $\mu$ l of the mixture was analyzed by GC-MS NICI.

For quantitation, human control CSF samples (100  $\mu$ l per sample) were spiked with the appropriate amounts of VPA-GLU, VPA-GLN, and VPA-GLY to yield concentrations ranging from 0.01 to 0.10  $\mu$ g/ml. The calibration standards, 100  $\mu$ l of patient's CSF and control CSF were spiked with 5  $\mu$ l of FVPA-GLN (0.08  $\mu$ g/ml) and filtered through a filter membrane as described for the serum samples. Then, 60  $\mu$ l of the filtrate was injected into the LC-MS/MS system to be analyzed using LC-MS/MS method 3 as described for the serum samples.

The range of the calibration curve for each conjugate was 10 to 100 ng/ml. Standard curves afforded  $r^2$  of 0.993, 0.979, and 0.999, for VPA-GLU, VPA-GLN, and VPA-GLY, respectively.

## Results

**Synthesis of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP.** The synthesis of the amino acid conjugates of VPA was facile and involved the conversion of VPA to *N*-hydroxysuccinimide ester in the presence of dicyclohexylcarbodiimide followed by coupling with the appropriate amino acid.

Proton nuclear magnetic resonance spectroscopy of all the conjugates served as a primary tool to verify the chemico-selectivity of the conjugation reaction and to determine the structure of the four conjugates synthesized. Spectra were characterized by signals of a VPA backbone and those belonging to the amino acid moiety. The signals for the proton(s) of the  $\alpha$ -methine of the amino acid confirmed the structure of the conjugates was consistent with that of the amide of valproic acid. In the case of VPA-GLU, VPA-GLN, and VPA-ASP, this proton was coupled to the amino proton and the neighboring  $-\text{CH}_2$ -group and was split into a doublet of doublets at 4.45, 4.40, and 4.80 ppm, respectively. In VPA-GLY, the protons on the same tertiary carbon were detected as a doublet at 4.1 ppm. The structures of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP are shown in Fig. 1.

LC-MS scanning showed the  $[\text{MH}]^+$  of VPA-GLU, VPA-GLN, VPA-GLY, and VPA-ASP to be  $m/z$  274,  $m/z$  273,  $m/z$  201, and  $m/z$

TABLE 3

Summary of results for the profiling of VPA-GLU, VPA-GLN, VPA-GLY in the urine samples of patients on VPA

	VPA-GLU ( $\mu\text{g}/\text{mg}$ creatinine) <i>n</i> = 29	VPA-GLN ( $\mu\text{g}/\text{mg}$ creatinine) <i>n</i> = 29	VPA-GLY ( $\mu\text{g}/\text{mg}$ creatinine) <i>n</i> = 29
Range	0.66–13.1	0.78–9.93	Trace–1.0
Mean	4.31	4.05	0.31
S.D.	2.76	2.50	0.37

260, respectively. Upon CID of the  $[\text{MH}]^+$  of each metabolite, the resulting product ion spectra showed that the most abundant fragment at  $m/z$  57 was shared by all four conjugates as illustrated in Fig. 1. This fragment is believed to be the protonated *n*-butyl group of VPA following cleavage on two sides of the tertiary carbon ( $\text{C}_2$ ) causing the loss of the formyl amino acid group and the *n*-propyl group. The other product ions included  $m/z$  99 resulting from fragmentation at the carbonyl carbon ( $\text{C}_1$ ) causing the neutral loss of the *n*-formyl amino acid group leaving the protonated *n*-heptanyl chain of VPA. Cleavage of the amide bond gave rise to the protonated 2-propylpentanone fragment at  $m/z$  127 for each conjugate.

The product ion spectra of each compound (described in Fig. 1) following CID of their  $[\text{MH}]^+$  are shown in Fig. 2. The corresponding protonated amino acid moiety was also observed at  $m/z$  148 for VPA-GLU or  $m/z$  147 for VPA-GLN or  $m/z$  134 for VPA-ASP but not for VPA-GLY. The subsequent loss of a water molecule afforded a product ion at  $m/z$  129 from the protonated glutamine and  $m/z$  130 from the protonated glutamic acid. In contrast, the product ion spectrum of  $[\text{MH}]^+$  of FVPA-GLN had a different fragmentation pattern compared with the nonfluorinated analog and was consistent with previously reported data (Tang et al., 1997).

GC-MS NICI analysis of the PFB esters or di-esters of the conjugates served to complement the structural characterization of the compounds. The GC-MS NICI spectrum of each derivatized conjugate was characterized by a single base peak corresponding to the  $[\text{M-pentafluorobenzyl}]^-$  fragment anion of the di-PFB ester of VPA-GLU ( $m/z$  452) and VPA-ASP ( $m/z$  438) or the PFB ester of VPA-GLN ( $m/z$  271), VPA-GLY ( $m/z$  200) and FVPA-GLN ( $m/z$  289). Elution times under our GC-MS NICI experimental conditions are shown in Table 1.

**Identification of Amino Acid Conjugates of VPA in Urine Samples of Patients.** The amino acid conjugates of VPA eluted between 22.97 and 30.30 min by LC-MS/MS method 1, 14.70 and 23.99 min by method 2, and 7.86 and 11.99 min by method 3 as described in Table 2.

In all the urine samples ( $n = 33$ ) studied, all product ions of  $[\text{MH}]^+$  of VPA-GLU, VPA-GLN, and VPA-GLY were detected at the same retention times and in the same area ratio as those observed for synthetic standards of the conjugates by two mobile phase systems (water/acetonitrile and water/methanol) in MRM mode. None of the product ions were detected in the control samples. A full product ion spectrum of VPA-GLU and VPA-GLN identical to those of reference compounds shown in Fig. 2 were obtained in four patients' urine extracts. MRM experiments failed to detect any product ions of the

protonated VPA-ASP in any of the patients' samples studied. The search for VPA taurine and VPA alanine was approached by MRM of the corresponding transition  $[\text{MH}]^+$  to  $m/z$  57, the most abundant product ion shared by the identified amino acid conjugates of VPA. Neither of the two conjugates were detected in the urine extracts analyzed.

In the patients' urine samples analyzed, VPA-GLU was found at a concentration range of 0.66 to 13.1  $\mu\text{g}/\text{mg}$  creatinine; VPA-GLN was found at lower concentration range of 0.78–9.93  $\mu\text{g}/\text{mg}$  creatinine, whereas VPA-GLY was the minor metabolite at a concentration range of trace–1.0  $\mu\text{g}/\text{mg}$  creatinine (Table 3). The total recoveries of the three conjugates, VPA-GLU, VPA-GLN, and VPA-GLY in urine samples of patients on VPA therapy were calculated for a 24-h period and found to range from 0.53 to 1.45% ( $0.75 \pm 0.41$ ) of a VPA dose (Table 4). The urinary VPA concentration was available for three of the patients, and the mean concentration was found to be 420  $\mu\text{g}/\text{mg}$  of creatinine.

**Identification of Amino Acid Conjugates of VPA in Serum Samples of Patients.** Analysis of serum samples by LC-MS/MS MRM provided strong evidence for the identification of VPA-GLU, VPA-GLN, and VPA-GLY in the serum extracts of all six patients but not in any control serum. The retention times and area ratio of all product ions of  $[\text{MH}]^+$  of the metabolites were authenticated by comparison with synthetic standards. The mean serum concentration of VPA-GLU in six patients was  $8.0 \pm 2.0$  ng/ml. VPA-GLN was estimated at  $5.0 \pm 3.0$  ng/ml, and VPA-GLY was detected but was below the limit of quantitation of the assay.

**Identification of Amino Acid Conjugates of VPA in Human CSF.** The retention times and area ratio of all the product ions of the  $[\text{MH}]^+$  of VPA-GLU (Fig. 3) and VPA-GLN in a human CSF sample were the same as those observed for both synthesized amino acid conjugates of VPA by LC-MS/MS. The identification of VPA-GLU was further investigated by GC-MS under NICI by monitoring for the  $[\text{M-181}]^-$  fragment carboxylate anion of the di-PFB derivative of the conjugate at  $m/z$  452. The total ion current of the PFB-derivatized CSF extract (Fig. 4) and its corresponding PFB-derivatized serum extract showed the elution of VPA-GLU at  $t_R = 16.36$  min, the identical retention time observed for the synthetic standard. The peak was absent in the PFB-derivatized control CSF and serum extract.

The quantitation of VPA-GLU, and VPA-GLN in CSF by LC-MS/MS was relatively facile because no extraction procedure was required. The assay was designed to monitor one characteristic MRM transition for each amino acid conjugate of VPA identified in human and using VPA-ASP as the IS. A relatively high volume (60  $\mu\text{l}$ ) of sample was introduced onto the HPLC column, but the resulting slight broadening of the peaks was not chromatographically significant. The concentration of VPA-GLU in the CSF of one patient was 60 ng/ml, and that of VPA-GLN was 34 ng/ml of CSF. The corresponding serum concentrations of VPA-GLU and VPA-GLN of the same patient were 7.0 ng/ml and 6.0 ng/ml, respectively. The concentration of CSF VPA of the patient was less than 15  $\mu\text{g}/\text{ml}$  of CSF.

TABLE 4

The % of VPA dose recovered as VPA-GLU, VPA-GLN, and VPA-GLY in four patients over 24 h

Patient No.	VPA GLU	VPA GLN	VPA GLY	Total % Recovered
1	0.45	0.32	0.03	1.45
2	0.20	0.20	0	0.40
3	0.22	0.18	0.02	0.61
4	0.17	0.15	0	0.53
Mean $\pm$ S.D.				$0.75 \pm 0.41$




Fig. 3. On-line LC-MS/MS monitoring of VPA-GLU in (a) a control human CSF sample, (b) a CSF sample of a patient treated with VPA, and (c) a control human CSF sample spiked with VPA-GLU, which eluted at  $t_R = 9.66$  min.

The corresponding characteristic ion transitions monitored for MRM are shown in (d) [mobile phase, acetonitrile (40%):H<sub>2</sub>O (60%) and TFA (0.05%)].

### Discussion

We report here the identification of valproyl glutamate and valproyl glutamine, both conjugates of VPA in humans. Valproyl glycine was also identified in the biological samples of patients on VPA. Preliminary accounts of these findings were reported (Gopaul et al., 1997a,b).

The discovery of VPA-GLU, in the biological fluids of humans, is intriguing due to its extreme rarity. To the best of our knowledge, the glutamate conjugate of benzoic acid in the Indian fruit bat (Idle et al., 1975) and *p*-nitrobenzoic acid in spiders (Smith, 1962) are the only two glutamate conjugates of a xenobiotic carboxylic acid reported in the literature. Conjugates of amino acids that are normally found are species-dependent and vary according to the structure of the carboxylic acid involved.

In humans, the conjugation of glycine, glutamine, taurine, and ornithine with carboxylic acids are known (Caldwell et al., 1976; Webster et al., 1976; Hutt and Caldwell, 1990; Shirley et al., 1994; Kasuya et al., 1999). Reports in the literature indicate that glycine and to a lesser extent glutamine are the most likely amino acids to be involved in the conjugation reactions of carboxylic acids (Hutt and Caldwell, 1990). It should also be added that recently the conjugation of xenobiotic amines with glutamate has also been reported (Mutlib et al., 2000; 2002).

In this study, quantitative analysis of the conjugates in human urine, serum, and CSF samples revealed that VPA-GLU was consistently the predominant amino acid conjugate compared with VPA-GLN, and VPA-GLY was a minor conjugate. The more extensive conjugation of VPA with glutamic acid than either glutamine or glycine is also an unprecedented observation for the human species.

The higher abundance of the amino acid conjugates in urine samples relative to their serum concentrations suggests that the conjugates are readily excreted upon formation in the liver. These metabolites can also be formed at other sites, such as the kidney. For example, in vitro studies support the conjugation of benzoic acid with glycine in human liver and kidney tissues (Caldwell et al., 1976; Temellini et al., 1993). Alternatively, a portion of the conjugates formed may be excreted into the intestinal lumen (via biliary excretion or exsorption) and excreted in feces.

Amino acid conjugation of an organic acid involves the activation of the acid to its CoA derivative prior to coupling with an amino acid by the respective amino acid transferase (Hutt and Caldwell, 1990). The mean observed recovery of amino acid conjugates of VPA in the urine samples of four patients was  $0.75 \pm 0.41\%$  of the VPA dose. By comparison, the  $\beta$ -oxidation metabolites of VPA, namely 2-ene VPA and 3-keto VPA, that are also initiated by the activation of VPA to its CoA ester, account for more than 15% of a VPA dose in humans (Pollack et al., 1986). This would imply that the VPA CoA ester is more susceptible to  $\beta$ -oxidation than amino acid conjugation in mitochondria. However, whether the amino acid conjugation of VPA is a mitochondrial or microsomal-based reaction remains to be confirmed.

The identification of VPA-GLU and VPA-GLN in the CSF sample stood as the most interesting finding in this study. The availability of human CSF of patients on VPA therapy is rare, and as part of our efforts to support this human evidence, we reported the results of a preliminary study that confirmed unequivocally the identification of VPA-GLU as the only amino acid conjugate of VPA in the CSF of




Fig. 4. Selected ion chromatogram of the PFB-derivatized extract of a CSF sample of a patient on VPA monitoring the  $[M-181]$ carboxylate anion of the PFB derivative of VPA-GLU at  $m/z$  452, which eluted at  $t_R = 16.36$  min.

rabbits treated with VPA ( $n = 5$ ) (Gopaul et al., 1997b). That VPA-GLN was not detected in the CSF of rabbit suggests that VPA-GLU is probably the direct result of the conjugation of VPA with glutamic acid and is not primarily the hydrolyzed product of VPA-GLN although this needs to be confirmed.

The CSF results of the patient in this study are consistent with the levels of VPA metabolites that are usually found at concentrations  $<10\%$  of VPA in the CSF (Löscher et al., 1988) or  $<15\%$  of VPA in the human brain (Adkinson et al., 1995). The CSF concentrations of VPA-GLU and VPA-GLN in the patient in this study were  $<1\%$  of the CSF VPA level ( $15 \mu\text{g/ml}$  of CSF). Because, the concentrations of VPA and unsaturated VPA metabolites in the CSF are believed to correspond to those in the brain (Vajda et al., 1981; Löscher et al., 1988), it would be reasonable to suggest that the levels of the CSF VPA-GLU and VPA-GLN reflect their respective brain tissue concentrations. However, this would be inconsistent with a recent report from Scism et al. (2000), which suggests that VPA may be concentrated in the intracellular compartments of the brain parenchyma.

The CSF concentrations of VPA-GLU, and VPA-GLN were 9 and 5 times, respectively, their corresponding serum concentrations in the human subject, suggesting that the metabolites could be formed in the brain itself as a result of the conjugation of VPA with glutamic acid or its precursor, glutamine. If so, the detection of VPA-GLU and VPA-GLN in the CSF of human and rabbit provides some grounds to propose that the mechanism of action of VPA could in part be the result of the conjugation of VPA with the excitatory neurotransmitter glutamic acid believed to be responsible for seizure activity or its precursor glutamine. In support of this theory, significantly elevated concentrations of GLU are found in human epileptogenic foci (Perry and Hansen, 1981) and in the CSF of newly diagnosed epileptic patients untreated with anticonvulsants (Kälviäinen et al., 1993). Although the concentrations of human CSF VPA-GLU with respect to the concentration of VPA itself ( $\sim 15 \mu\text{g/ml}$ ) is low, it is consistent with the concentration of CSF glutamate that is usually found in comparatively low concentrations in the CSF of control subjects

(Kälviäinen et al., 1993; Jimenez-Jimenez et al., 1998). In light of the implication of VPA metabolism in the mechanism of action of VPA and the induction of hepatotoxicity, the significance of this new metabolic pathway, namely the glutamic acid conjugation with VPA in humans, needs to be pursued.

### Conclusion

LC-MS/MS was used for the identification and quantitation of the novel VPA-GLU and VPA-GLN conjugates in the urine, serum, and CSF samples of patients on VPA. These amino acid conjugates of VPA in urine account for about 1% of a VPA dose. The identification of VPA-GLU and VPA-GLN in the CSF of a human subject stands as the most interesting finding with the detection of amino acid conjugates of VPA in the CSF of humans. In our opinion, the significance of these conjugates in the CNS requires further investigation. A report on the studies of the amino acid conjugates of VPA in animals is to follow.

**Acknowledgments.** We thank Roland Burton for his technical assistance.

### References

- Abbott FS and Anari MR (1999) Chemistry and biotransformation, in *Valproate* (Löscher W, ed) pp 47–75, Birkhauser Verlag, Basel/Switzerland.
- Adkinson K, Ojemann GA, Rappert RL, Dills RL, and Shen DD (1995) Distribution of unsaturated metabolites of valproate in human rat brain—pharmacologic relevance? *Epilepsia* 36:772–782.
- Baillie TA and Sheffels PR (1995) Valproic acid. Chemistry and biotransformation, in *Antiepileptic Drugs*, 4th edition (Levy RH, Mattson RH, and Meldrum BS eds) pp 589–604, Raven Press, New York.
- Caldwell J, Moffatt JR, and Smith RL (1976) Post-mortem survival of hippuric acid formation in rat and human cadaver tissue samples. *Xenobiotica* 6:275–280.
- Eadie MJ, Hooper WD, and Dickinson RG (1988) Valproate-associated hepatotoxicity and its biochemical mechanisms. *Med Toxicol* 3:85–106.
- Gopaul SV (1998) *The Identification, Characterization, and Profiling of Novel Thiol and Amino Acid Conjugates of Valproic Acid in Humans and Animals*. Ph.D. Thesis, The University of British Columbia, Vancouver, BC, Canada.
- Gopaul SV, Farrell K, and Abbott FS (1997a) Amino acid conjugation of valproic acid in humans. *ISSX Proc* 11:205.
- Gopaul SV, Tabatabaei AR, and Abbott FS (1997b) The identification of novel amino acid conjugates of valproic acid in whole brain and cerebral spinal fluid of rat and rabbit using GC/MS and LC/MS/MS. *Pharma Res (NY)* 14:S-355.

- Granneman GR, Wang SI, Kesterson JW, and Machinist JM (1984a) The hepatotoxicity of valproic acid and its metabolites in rats. II. Intermediary and valproic acid metabolism. *Hepatology* **4**:1153–1158.
- Granneman GR, Wang SI, Machinist JM, and Kesterson JW (1984b) Aspects of the metabolism of valproic acid. *Xenobiotica* **14**:375–387.
- Harris RA, Swartzentruber M, Becker CM, McCune SA, Hu H, and Popov K (1991) Metabolic effects of sequestration of hepatic mitochondrial coenzyme A by valproate and other carboxylic acids, in *Idiosyncratic Reactions to Valproate: Clinical Risk Patterns and Mechanisms of Toxicity* (Levy RH and Penry JK eds) pp 97–103, Raven Press, New York.
- Hutt AJ and Caldwell J (1990) Amino acid conjugation, in *Conjugation Reactions in Drug Metabolism* (Mulder GJ ed) pp 273–305, Taylor and Francis Ltd., New York.
- Idle JR, Millburn P, and Williams RT (1975) Benzoylglutamic acid, a metabolite of benzoic acid in indian fruit bats. *FEBS Lett* **59**:234–235.
- Jaffe MZ (1886) Über den Niederschlag welchen pikrinsäure in normalen Harn erzeugt und Über eine neue Reaktion des Kreatinins. *Z Physiol Chem* **10**:391–400.
- Jimenez-Jimenez FJ, Molina JA, Gomez P, Vargas C, de Bustos F, Benito-Leon J, Tallon-Barranco A, Orti-Pareja M, Gasalla T, and Arenas J (1998) Neurotransmitter amino acids in cerebrospinal fluid of patients with Alzheimer's disease. *J Neural Transm* **105**:269–277.
- Kälviäinen R, Halonen T, Pitkänen A, and Reikkinen P (1993) Amino acids levels in the cerebrospinal fluid of newly diagnosed epileptic patients: effect of vigabatrin and carbamazepine monotherapies. *J Neurochem* **60**:1244–1250.
- Kasuya F, Igarashi K, and Fukui M (1999) Characterization of a renal medium chain acyl-CoA synthetase responsible for glycine conjugation in mouse mitochondria. *Chem Biol Interact* **118**:233–246.
- Löshner W, Nau H, and Siemes H (1988) Penetration of valproate and its active metabolites into cerebrospinal fluid of children with epilepsy. *Epilepsia* **29**:311–316.
- Mawal Y, Paradis K, and Quereshi JA (1997) Developmental profile of mitochondrial glycine N-acyltransferase in human liver. *J Pediatr* **130**:1003–1007.
- Mutlib AE, Shockcor J, Chen SY, Espina RJ, Pinto DJ, Orwat MJ, Prakash SR, and Gan LS (2002) Disposition of 1-[3-(Aminomethyl)phenyl]-N-[3-fluoro-2'-(methylsulfonyl)-[1, 1'-biphenyl]-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (DPC 423) by novel metabolic pathways. Characterization of unusual metabolites by liquid chromatography/mass spectrometry and NMR. *Chem Res Toxicol* **15**:48–62.
- Mutlib AE, Shockcor J, Espina R, Graciana N, Du A, and Gan LS (2000) Disposition of glutathione conjugates in rats by a novel glutamic pathway. Characterization of unique peptide conjugates by LC/MS and LC/NMR. *J Pharmacol Exp Ther* **294**:735–745.
- Perry TL and Hansen S (1981) Amino acid abnormalities in epileptogenic foci. *Neurology* **31**:872–876.
- Pollack GM, McHugh WB, Gengo FM, Ermer JC, and Shen DD (1986) Accumulation and washout kinetics of valproic acid and its active metabolites. *J Clin Pharmacol* **26**:668–676.
- Radatz M and Nau H (1999) Toxicity, in *Valproate* (Löscher W ed) pp 91–128, Birkhauser Verlag, Basel/Switzerland.
- Scism JL, Powers KM, Artru AA, Lewis L, and Shen DD (2000) Probenicid-inhibitable efflux transport of valproic acid in the brain parenchymal cells of rabbits: a microdialysis study. *Brain Res* **884**:77–86.
- Shirley NA, Guan X, Kaiser GW, Halstead GW, and Baillie TA (1994) Taurine conjugation of Ibuprofen in humans and rat in liver in vitro. *J Pharmacol Exp Ther* **269**:1166–1175.
- Smith JN (1962) Detoxications of aromatics acids with glutamic acid and arginine in spiders. *Nature (Lond)* **195**:399–400.
- Tang W, Palaty J, and Abbott FS (1997) Time course of  $\alpha$ -fluorinated valproic acid in mouse brain and serum and its effect on synaptosomal  $\gamma$ -aminobutyric acid levels in comparison to valproic acid. *J Pharmacol Exp Ther* **282**:1163–1172.
- Temellini A, Mogavero S, Giulianotti PC, Pietrabissa A, Mosca F, and Pacifici GM (1993) Conjugation of benzoic acid with glycine in human liver and kidney: a study on the interindividual variability. *Xenobiotica* **23**:1427–1433.
- Tishler SL and Goldman P (1970) Properties and reactions of salicyl-coenzyme A. *Biochem Pharmacol* **19**:143–150.
- Turnbull DM, Bone AJ, Barlett K, Koundakjian PP, and Sherratt HS (1983) The effects of valproate intermediary metabolism in isolated rat hepatocytes and intact rats. *Biochem Pharmacol* **32**:1887–1892.
- Vajda FJE, Donnan GA, Phillips J, and Bladin PF (1981) Human brain, plasma and cerebrospinal fluid concentration of sodium valproate after 72 hours of therapy. *Neurology* **31**:486–587.
- Webster LT, Siddiqui UA, Lucas SV, Strong JM, and Miesel JJ (1976) Identification of separate acyl-CoA:glycine and acyl-CoA:l-glutamine N-acyltransferase activities in mitochondrial fractions of rhesus monkey and man. *J Biol Chem* **251**:3352–3358.
- Williams DA (1989) Drug metabolism, in *Principles of Medicinal Chemistry* (Foye WO ed) 3rd edition, pp 100–102, Lea & Febiger, London.