

Thyroid Hormone Is Necessary for Expression of Constitutive Androstane Receptor in Rat Hepatocytes

Hidekazu Ooe, Junko Kon, Hideki Oshima, and Toshihiro Mitaka

Department of Pathophysiology, Cancer Research Institute (H.Oo., J.K., H.Os., T.M.), and Department of Surgery Section I (H.Os.), Sapporo Medical University School of Medicine, Sapporo, Japan

Received June 13, 2008; accepted June 9, 2009

ABSTRACT:

Small hepatocytes are hepatocyte progenitor cells that possess the capability of maturation and cryopreservation. When cryopreserved rat small hepatocytes were cultured in serum-free medium, the protein expression and the inducibility of CYP1A1/2, CYP2E1, and CYP3A were maintained, but those of CYP2B1 were lost. In this study we investigated the cause of the loss of CYP2B1 expression in cryopreserved small hepatocytes by reverse transcription-polymerase chain reaction, immunoblotting, and chromatin immunoprecipitation assay. Expression of mRNA and protein of the nuclear receptor, constitutive androstane receptor (CAR), which regulates the expression of CYP2B1, was inhibited in the serum-free culture of cryopreserved small hepatocytes, whereas they were expressed

in that of subcultured small hepatocytes. Serum application dramatically induced CAR expression in the culture of cryopreserved small hepatocytes. The addition of very low concentrations of thyroid hormones (THs; 3,5,3'-triiodothyronine, 5×10^{-12} M; thyroxine, 5×10^{-12} - 5×10^{-10} M) to the medium also induced the expression of CAR and CYP2B1. Moreover, CYP2B1 expression was induced by administration of phenobarbital. In rats with hypothyroidism induced by thyroidectomy and 6-propyl-2-thiouracil treatment, the expression of CAR and CYP2B1 was strongly repressed. Although THs do not directly regulate the expression of CAR, they may be important for rat hepatocytes to regulate CYP2B1 through CAR expression in the physiological condition.

Cytochromes P450 (P450s) constitute a superfamily of monooxygenases that play a key role in either the detoxification or the metabolic activation of xenobiotics (Guengerich and MacDonald, 1990; Wrighton and Stevens, 1992; Gonzalez and Gelboin, 1994). The P450s involved in xenobiotic metabolism are rich in hepatocytes. In vivo, many of the constitutive P450s are inducible by xenobiotics such as aromatic hydrocarbons, drugs, and alcohol. However, in traditional culture systems, it is very difficult to maintain the P450 expression of hepatocytes, and the activity of most P450s is rapidly lost after plating (Bissell and Guzelian, 1980). In cryopreserved hepatocytes isolated from both rodents and humans, hepatic differentiated functions, including P450 activities, also decrease immediately after plating (Jackson et al., 1985; Loretz et al., 1989; Sun et al., 1990; Chesné et al., 1993; de Sousa et al., 1996; Swales et al., 1996; Garcia et al., 2003).

Small hepatocytes (SHs) have been identified as proliferating cells with hepatic characteristics (Mitaka et al., 1992, 1995; Tateno and

Yoshizato, 1996). We showed that a single SH could clonally proliferate to form colonies (Mitaka et al., 1999). It was also reported that maturation of the proliferating SHs was induced by the application of Engelbreth-Holm-Swarm sarcoma-derived matrix (Matrigel; BD Biosciences, Bedford, MA) (Sugimoto et al., 2002). Moreover, several P450 proteins were expressed and induced in SHs even after more than 1 month of culture (Miyamoto et al., 2005). On the other hand, we reported that SHs could be cryopreserved for more than 6 months and that, even after thawing, the cells could maintain the capability of the growth and the maturation (Ikeda et al., 2002). We also examined the expression of CYP1A1/2, CYP2E1, and CYP3A in thawed SHs even after the cells were cultured in a serum-free medium for more than 1 month (Ooe et al., 2006). However, CYP2B1 was not expressed in the culture even if hepatic maturation of SHs was induced by Matrigel and induction by phenobarbital was carried out (Ooe et al., 2006).

Nuclear receptors such as aryl hydrocarbon receptor (AhR), constitutive androstane receptor (CAR), retinoid X receptor (RXR), and pregnane X receptor (PXR) are related to the transcription of P450 and other enzyme genes (Honkakoski et al., 1998; Ma, 2001; Kliewer, 2003). CAR and RXR play a central role in constitutive expression and the induction of CYP2B. The translocation of CAR from the cytoplasm to the nucleus was observed after phenobarbital treatment (Kawamoto et al., 1999). In the nucleus, CAR forms heterodimer with

This work was supported in part by the Ministry of Education, Culture, Sports, Science, and Technology, Japan [Grant 17390353]; the Ministry of Health, Labour, and Welfare; Health and Labour Sciences Research Grants; Research on Advanced Medical Technology; and the Suhara Memorial Foundation.

Article, publication date, and citation information can be found at <http://dmd.aspetjournals.org>.

doi:10.1124/dmd.108.022905.

ABBREVIATIONS: P450, cytochrome P450; SH, small hepatocyte; AhR, aryl hydrocarbon receptor; CAR, constitutive androstane receptor; RXR, retinoid X receptor; PXR, pregnane X receptor; PB, phenobarbital; cryo-SH, cryopreserved small hepatocyte; TH, thyroid hormone; MH, mature hepatocyte; PTU, 6-propyl-2-thiouracil; T₃, 3,5,3'-triiodothyronine; T₄, thyroxine; FBS, fetal bovine serum; PBS, phosphate-buffered saline; G3PDH, glycerol 3 phosphate dehydrogenase; PCR, polymerase chain reaction; PAGE, polyacrylamide gel electrophoresis; ChIP, chromatin immunoprecipitation; TR, thyroid hormone receptor; sub-SH, subcultured small hepatocyte; TRE, TH-responsive element; RT-PCR, reverse transcription-polymerase chain reaction; Dio1, 5'-deiodinase type 1; ME1, malic enzyme 1; Thrsp, thyroid hormone-responsive gene.

TABLE 1
Primer list and PCR conditions

Gene		Sequence (5'-3')	Cycle	Annealing Temperature °C	Amplicon Size base pair
CAR	Sense	ATGACAGCTACTCTAACTAGAG	27	56	1076
	Antisense	CAGCTGCAAATCTCCCAAGCAGC			
RXR	Sense	ACCCCGTGAGCAGCAGTGAGG	32	60	992
	Antisense	CCCAATGGATCGCAGTGCAGG			
AhR	Sense	CCGTCCATCCTGGAAATTCGAACC	30	60	350
	Antisense	CCTTCTTCATCCGTTAGCGGTCTC			
PXR	Sense	ATGAGACCTGAGGAGAGGTGGAAC	30	60	1007
	Antisense	TCCTCCTCACGCAGCTGTAGCTTC			
CYP2B1	Sense	GGGACACCCAAAGTCCCGTGG	27	56	867
	Antisense	GGAAACCATAAGCGAGTGTGG			
TR α	Sense	ACGTTGCTACAGAGGCCATC	32	56	700
	Antisense	AGGAAGAGTGGGGGAAGAGTT			
TR β	Sense	GAAAATCGCCTTCCAGCCTGG	32	56	619
	Antisense	GCTCTGGCTTATGCCAATGG			
Dio1	Sense	ATGGGGCTGTCCAGCTATGG	32	56	567
	Antisense	AGCAGATGTGCTGCCGCAGG			
ME1	Sense	CCTGAACCCCAAAACAAAGA	32	56	263
	Antisense	CCCATGAGTCTGTGGCTAA			
Thrsp	Sense	CCTCATCAGATCCAACAGGATG	32	56	212
	Antisense	GCAGGGACGCTGTATTACCTC			
G3PDH	Sense	ACCACAGTCCATGCCATCAC	22	56	451
	Antisense	TCCACCACCTGTTGCTGTA			
ME1 promoter	Sense	TGGACCTGTGCCCTAACAC	30	56	379
	Antisense	TCCCGCGTCAGCAGGTAGC			
CAR promoter 1	Sense	GATAGTGACAGTGTACTCATAAC	35	56	424
	Antisense	CCAGGGTGTACAAGCCTAGCC			
CAR promoter 2	Sense	GGAAGCTCCAGAATAAAGAGAG	35	56	502
	Antisense	GAGATCCCACTTTCTTACAAGG			

RXR and binds to DNA sequence called phenobarbital (PB)-responsive enhancer module (Honkakoski et al., 1998).

The aim of this study is to clarify the reason why no expression of CYP2B1 occurs in cryopreserved SHs (cryo-SHs). The results revealed that CAR expression was inhibited in the cryo-SHs when the cells were cultured in serum-free medium. The expression of CAR and CYP2B1 was recovered by the addition of serum. Furthermore, the physiological concentration of thyroid hormones (THs) could be substituted for the serum. Here, we show the importance of THs for the expression of CAR and CYP2B1 in a physiological condition.

Materials and Methods

Animals. Male F344 rats (Sankyo Lab Service, Tokyo, Japan), weighing 170 to 240 g, were used. Thyroidectomized F344 rats (male, 7 weeks old) were purchased from Sankyo Lab Service. All the animals received humane care, and the experimental protocol was approved by the Committee of Laboratory Animals according to Sapporo Medical University guidelines. They received a diet and drinking water ad libitum and were maintained in a specific pathogen free animal facility at $23 \pm 1^\circ\text{C}$ on a standard 12-h light/dark cycle. Intraperitoneal administration of PB (25 mg/kg b.wt.; Wako Pure Chemicals, Tokyo, Japan) was repeated every day for four times before preparation of mature hepatocytes (MHs). The surgically operated rats were used within 2 weeks after the operation. 6-Propyl-2-thiouracil (PTU; Sigma-Aldrich, St. Louis, MO) treatment was performed by the addition of 0.1% (w/v) PTU to drinking water for 2 weeks before preparation of MHs. PTU treatment of thyroidectomized rats was done only for 1 week. Serum were obtained before the liver perfusion and stored at -20°C until use. The concentration of 3,5,3'-triiodo-L-thyronine (T_3), thyroxine (T_4), and thyroid-stimulating hormone in serum was determined on the ADVIA Centaur immunoassay system (Bayer, Leverkusen, Germany).

Isolation and Culture of SHs. Hepatic cells were isolated by the two-step collagenase perfusion method. Cell suspension was centrifuged at 50g for 1 min. The supernatant was used for preparing SHs, and the pellet was used for MHs. Details of the isolation and culture procedure for SHs and MHs were described previously (Mitaka et al., 1999). Finally, 6×10^4 viable cells/cm² were seeded on a 100-mm dish and cultured in Dulbecco's modified Eagle's

medium (Sigma-Aldrich) supplemented with 20 mM HEPES, 25 mM NaHCO₃, 30 mg/l L-proline, 10% (v/v) fetal bovine serum (FBS; HyClone Laboratories, Logan, UT), 10 mM nicotinamide (Katayama Chemical Industries Co., Osaka, Japan), 1 mM ascorbic acid 2-phosphate (Wako Pure Chemicals), 10 ng/ml epidermal growth factor (BD Biosciences), 0.5 mg/l insulin, 10^{-7} M dexamethasone, and antibiotics. After 4 days of culture, 1% (v/v) dimethyl sulfoxide (Aldrich Chemical Co., Milwaukee, WI) was added to the medium.

Subculture and Cryopreservation of SH Colonies. As previously reported (Ooe et al., 2006), SH colonies consisting of 30 to 50 cells were observed at day 14 after plating. To collect SHs, the colonies were detached from dishes and immediately replated (subculture) or cryopreserved at -80°C until use. The cells that were cryopreserved for more than 1 month were used. SH colonies ($3\text{--}5 \times 10^3$ colonies/60-mm dish) were replated on new dishes coated with rat tail collagen. One day after replating, the medium was replaced with serum-free modified Dulbecco's modified Eagle's medium supplemented with or without TH. PB was used as inducer for CYP2B. Ten days after replating, fresh medium containing 2 mM PB was added. To enhance the CYP2B expression, the medium containing PB was renewed every day for 3 consecutive days before harvest. Some dishes were treated with Matrigel (1 mg/60-mm dish; BD Biosciences) from 7 days after replating.

RNA Preparation and Reverse Transcription-Polymerase Chain Reaction. For total RNA extraction, the cells were washed with phosphate-buffered saline (PBS) twice and treated with 1 ml of Isogen (Nippon Gene Co., Tokyo, Japan). RNA concentration was determined by measuring absorption at 260 nm. Reverse transcription was performed by OmniScript RT kit (QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instruction. cDNA was amplified with Taq polymerase (Fermentas Life Sciences, Burlington, ON, Canada). Sequences of forward and reverse primers, annealing temperature, and number of reaction cycle are listed in Table 1. Amplicon was separated by 1% (w/v) agarose gel and detected with ethidium bromide. Glycerol 3 phosphate dehydrogenase (G3PDH) was used as quantitative control for the estimation of relative expression.

Quantitative Real-Time Polymerase Chain Reaction. RNA was reverse-transcribed using an OmniScript RT kit (QIAGEN GmbH) and random hexamers. Real-time polymerase chain reaction (PCR) analyses were carried out on TaqMan Gene Expression Assays (Applied Biosystems, Foster City, CA)




FIG. 1. Expression of CYP2B1 in MHs and SHs. A, immunoblots for CYP2B1 protein in MHs and SHs. Proteins (5 $\mu\text{g}/\text{lane}$) were separated by 10% SDS-PAGE and transferred onto polyvinylidene difluoride membranes. MHs were prepared from a normal rat or a PB-treated rat liver. SH colonies were isolated at day 14, replated on a new dish, and cultured for 14 days (Sub). Some colonies were cryopreserved for more than 1 month at -80°C . Frozen colonies (F) were thawed and cultured for 14 days (Cryo) with (+PB) or without 2 mM PB. B, detection of CYP2B1 transcripts in MHs and SHs by RT-PCR analysis. Total RNA was extracted from the cells prepared for the experiment used in A. Sequences of the forward and the reverse primers, annealing temperatures, and the number of reaction cycles are listed in Table 1.

for rat CAR (Rn00576085_m1), rat CYP2B1 (Rn01457875_m1), rat hepatic nuclear factor 4 α (Rn00573309_m1), and rat G3PDH (Rn99999916_s1). Reaction solutions (50 μl) containing cDNA samples (corresponding to 100 ng of total RNA), 1 \times TaqMan Universal Master Mix (Applied Biosystems), 900 nM primers, and 200 nM probes were analyzed on an ABI Prism 7500 (Applied Biosystems). After denaturation at 94°C for 10 min, the reaction was repeated 40 times with denaturation at 94°C for 15 s, annealing at 60°C , and extension for 1 min. Relative expression was determined during the log-linear phase of amplification. G3PDH was used for the normalization of relative expression.

Western Blot Analysis. For immunoblots, the dishes were washed with PBS twice and then treated with 1 ml of lysis buffer (10 mM Tris-HCl, pH 8.0, 5 mM EDTA, 150 mM NaCl, 1% (v/v) Triton X-100, 5 $\mu\text{g}/\text{ml}$ leupeptin, 5 $\mu\text{g}/\text{ml}$ pepstatin A) for 1 h at 4°C . The cells were scraped and used for protein extraction. The concentration of proteins was measured using a bicinchoninic acid assay kit (Pierce, Rockford, IL). Samples (5 $\mu\text{g}/\text{lane}$) were separated by SDS-polyacrylamide gel electrophoresis (PAGE). A goat anti-CYP2B1 (Daiichi Pure Chemicals Co., Tokyo, Japan) antibody was used for immunoblots. The details of the method were described previously (Ooe et al., 2006).

Chromatin Immunoprecipitation Assay. Chromatin immunoprecipitation (ChIP) assay was performed using a QuikChIP kit (Imgenex, San Diego, CA) according to the manufacturer's instructions. In brief, 4-day cultured MHs or 14-day cultured cryo-SHs were fixed with 1% formaldehyde at 37°C for 15 min. Cells were washed twice with ice-cold PBS and resuspended in 1 ml of lysis buffer. Sonication was performed to reduce the DNA length to between 200 and 1000 base pairs. A rabbit anti-TH receptor (TR) α antibody (10 μg ; Santa Cruz Biotechnology, Inc., Santa Cruz, CA) and chromatin solution (500 μl) were used for ChIP assay and PCR. The sequences of the forward and reverse primers, annealing temperature, and number of reaction cycles are listed in Table 1. The amplicon was separated using 1% (w/v) agarose gel and detected with ethidium bromide.

Statistical Analysis. Statistical analysis was performed using Student's *t* test. Data acquired from three independent experiments were analyzed. A *p* value of <0.05 was considered significant.

Results

CYP2B1 Expression in Rat Cryo-SHs. Our previous report showed that the expression of CYP2B1 was repressed in cultured cryo-SHs, although it was maintained in subcultured SHs (sub-SHs). Therefore, we first confirmed whether mRNA and protein of CYP2B1 could be expressed in the various conditions of SHs. Western blot analysis showed that only a small amount of CYP2B1 protein was detected in the frozen and cultured cryo-SHs (Fig. 1A). No induction




FIG. 2. Expression of CAR and RXR α in MHs and cryo-SHs. Total RNA was extracted from cryo-SHs with or without 2 mM PB and SHs treated with both 2 mM PB and Matrigel (BD Biosciences). Total RNA of MHs was also prepared from normal rat and PB-treated rat liver. CAR and RXR α mRNAs were detected by RT-PCR analysis. Sequences of the forward and the reverse primers, annealing temperatures, and the number of reaction cycles are listed in Table 1.

was observed in the cryo-SHs even after PB treatment. The low expression was observed despite the periods of cryopreservation (data not shown). On the other hand, sub-SHs could maintain the protein expression. As shown in Fig. 1B, the loss of CYP2B1 mRNA expression was also observed in cryo-SHs, whereas the sub-SHs maintained the expression. No induction of the gene was observed in the cryo-SHs, even after PB treatment.

Expression of Nuclear Receptors. The gene expression of CYP2B1 is regulated by nuclear receptors such as CAR and RXR (Honkakoski et al., 1998). To clarify the reason why the cryo-SHs lost the ability of CYP2B1 expression, we examined the expression of the nuclear receptors. As shown in Fig. 2, CAR mRNA was scarcely expressed in cryo-SHs even after the induction by PB and maturation with Matrigel, whereas as much RXR α mRNA was expressed in cryo-SHs as in MHs. Although, unlike in MHs, PB could not induce RXR α expression in the cryo-SHs, these results showed that low expression of CAR might be involved in the poor expression of CYP2B1. Comparing the previous (Miyamoto et al., 2005; Ooe et al., 2006) and present experiments, the difference of the experimental procedure was whether cells were cryopreserved. Thus, we compared the expression of major nuclear receptors involved in the regulation of P450 expression between sub-SHs and cryo-SHs. As shown in Fig. 3, in serum-free culture, the expression of CAR was much higher in sub-SHs than in cryo-SHs. AhR, PXR, and RXR α were expressed in both cell types as much as in MHs. Next, we also examined the effect of FBS on the expression of nuclear receptors. The expression of CAR was restored when cryo-SHs were cultured in the medium supplemented with 10% (v/v) FBS (Fig. 3). The addition of FBS to the medium could prevent the loss of CYP2B1 expression in cryo-SHs, and when the addition was performed at any time of the culture, the gene was always induced soon after (data not shown). However, the addition of FBS affected neither the expression of CAR in sub-SHs nor that of AhR, PXR, and RXR α in cryo-SHs.

Effects of TH on the Expression of CAR and CYP2B1. Next, we examined which factors in serum were critical for the regulation of CAR gene. The data of GenBank show that a TH-responsive element (TRE)-like sequence exists in approximately 1.3 kilobases upstream of the gene. Therefore, we investigated whether TH could induce the CAR expression in cultured cryo-SHs. Both T_3 and T_4 possess the TH activity in vivo. Physiological activity of T_3 is stronger than that of T_4 , and T_3 concentration in serum is lower than that of T_4 (Brent, 1994). As shown in Fig. 4A, the most effective doses of T_3 and T_4 to the expression of CAR mRNA were 5×10^{-12} M and 5×10^{-12} to 5×10^{-10} M, respectively. The dose-dependent induction by THs was not observed in the experiment, and the range for the effectiveness of THs




FIG. 3. Expression of nuclear receptors in SHs. Total RNA was extracted from MHs, sub-SHs, and cryo-SHs cultured in the medium with or without 10% (v/v) FBS. CAR, AhR, PXR, and RXR α mRNAs were detected by RT-PCR analysis. Sequences of the forward and the reverse primers, annealing temperatures, and the number of reaction cycles are listed in Table 1.




FIG. 4. Induction of CAR and CYP2B1 mRNAs by THs in cryo-SHs. A, expression of CAR and CYP2B1 in cryo-SHs under various concentrations of T₃ or T₄. Total RNA was extracted from the cells cultured for 14 days and used for quantitative real-time PCR analyses. Three independent experiments were carried out. The relative expression compared with MH was calculated and standardized with G3PDH expression. Bars show S.D., and asterisks indicate significant differences between control and each concentration (*, $p < 0.05$; **, $p < 0.01$). B, time courses of CAR and CYP2B1 expression in cryo-SHs treated with or without TH. Total RNA was extracted from the cells cultured SHs with or without 5×10^{-12} M T₃ at days 0, 3, 6, 10, and 14. Quantitative real-time PCR analyses were conducted. Three independent experiments were carried out. The relative expression compared with MH was calculated and standardized with G3PDH expression. Bars show S.D., and asterisks indicate significant differences between 0 and each day (*, $p < 0.05$; **, $p < 0.01$).




FIG. 5. Expression and induction of CYP2B1 in cryo-SHs. A, immunoblots for CYP2B1 protein in MH and cryo-SHs treated with T₃. Proteins ($5 \mu\text{g}/\text{lane}$) were separated by 10% SDS-PAGE and transferred onto a polyvinylidene difluoride membrane. Cryo-SHs were cultured in the medium supplemented with 0, 5×10^{-12} , or 1×10^{-8} M T₃ for 14 days. Three independent experiments were performed. B, relative expressions of CYP2B1 compared with MH were calculated. Bars show S.D., and asterisks indicate significant differences (*, $p < 0.05$).

was very narrow. Likewise, the expression of CYP2B1 was also correlated with that of CAR. A high concentration of THs has induced neither CAR nor CYP2B1 expression.

The time courses of CAR and CYP2B1 mRNA expression were examined in the medium supplemented with or without 5×10^{-12} M T₃. As shown in Fig. 4B, in the cells cultured without T₃, the mRNA expression of CAR was very low and that did not change during culture. The addition of T₃ to the medium significantly induced the expression of CAR from 6 days after the treatment, and it increased until day 10. The maximum expression of CAR mRNA was approximately 70% as much as that in MHs. The protein expression of CYP2B1 was examined in SHs cultured in the medium containing T₃. As shown in Fig. 5, A and B, CYP2B1 protein was induced when cryo-SHs were treated with 5×10^{-12} M T₃, whereas the protein was scarcely expressed in cryo-SHs when the cells were treated both without T₃ and with 1×10^{-8} M T₃. Moreover, an administration of PB in the cells treated with 5×10^{-12} M T₃ resulted in the induction of CYP2B1. However, neither induction was observed in the cells treated without T₃ or a high concentration of T₃.

Expression of TH Receptors and Deiodinase in Cryo-SHs. To confirm the reactivity of cryo-SHs to TH, we investigated the expression of TH-related genes. Reverse transcription-polymerase chain reaction (RT-PCR) analysis indicated that TR α was expressed as strongly in cryo-SHs as in MHs (Fig. 6). TR β was also expressed in cryo-SHs, but the amount was very low. It is known that T₄ is converted into T₃ by the action of the enzyme 5'-deiodinase type 1 (Dio1) in the liver (Köhrle, 2000). Dio1 was also expressed in cryo-SHs, corresponding to the reactivity of cryo-SHs to T₃. TH-responsive genes such as malic enzyme (ME) 1, Dio1, and thyroid hormone-responsive gene spot 14 (Thrsp) (Liaw and Towle, 1984; Song et al., 1988) were clearly up-regulated in cryo-SHs treated with 1×10^{-8} M T₃. However, the addition of FBS to the medium did not induce the up-regulation of those genes in cryo-SHs. Cryo-SHs were shown to be able to respond to TH.

CAR and CYP2B1 Expression in Hypothyroid Rats. Because it is important to examine whether hypothyroid rats lose the expression of CAR and CYP2B1, we investigated the expression of both genes in




FIG. 6. Expression of thyroid hormone-related genes in cryo-SHs. Expression of TH-related genes in MHs and cryo-SHs. TRs, Dio1, ME1, and Thrsp mRNAs were detected by RT-PCR analysis. Cryo-SHs were cultured in the medium supplemented with $0, 5 \times 10^{-12}$ M T_3 (+), 1×10^{-8} M T_3 (++), or 10% (v/v) FBS for 14 days. Sequences of the forward and the reverse primers, annealing temperatures, and the number of reaction cycles are listed in Table 1.

the liver and measured the concentration of THs in serum of rats with combined thyroidectomy and PTU treatment. To dramatically decrease the T_3 concentration in the serum, MHs were prepared from rat livers treated with both thyroidectomy and PTU. All the rats treated with both treatments were dead within 2 weeks after the treatment. Therefore, we examined the expression of CAR and CYP2B1 in the liver of the rats treated with the combination at day 7, whereas either thyroidectomized or PTU-treated rats were examined at day 14. As shown in Table 2, total T_3 concentrations in the serum of thyroidectomized and PTU-treated rats were approximately 0.35 and <0.15 nM, respectively. In the rats treated with both treatments, T_3 concentration and T_4 in the serum were both less than a measurable level. As shown in Fig. 7, either thyroidectomy or PTU treatment did not significantly decrease both CAR and CYP2B1 expression. However, both expressions were dramatically reduced in hepatocytes of the rats treated with the combination of thyroidectomy and PTU, although hepatic nuclear factor 4 expression of the cells was maintained, and the low concentration of TH in the serum might result in the induction of thyroid-stimulating hormone.

ChIP Assays. To confirm whether TR could directly regulate CAR expression, we performed ChIP assays using MHs and cryo-SHs cultured with and without T_3 . The promoter region of the rat *ME1* gene, in which the existence of TRE is well known, was selected to show that the ChIP assay was accurately performed. As shown in Fig. 8A, the binding of TR α to TRE (Petty et al., 1989) in the promoter region of the gene was detected in cryo-SHs and MHs. Moreover, the binding of TR α was enhanced by T_3 treatment. These data indicated that TR could bind to TRE in the gene in cryo-SHs as in MHs. However, as shown in Fig. 8B, the binding of TR α to the TRE-like sequence that exists in the promoter region of the rat *CAR* gene was not detected in cryo-SHs even after 1×10^{-8} M T_3 treatment.

Discussion

We previously reported that primary cultured SHs maintained the protein expression of CYP1A1, CYP2B1, CYP2E1, CYP3A2, and CYP4A1 for more than 1 month and that the expression increased with the maturation of SHs (Miyamoto et al., 2005). The enzymatic activity of CYP1A, CYP2B, and CYP3A could be also measured and increased with their maturation. In addition, both expression and activity of the P450s were maintained after long-term cryopreservation (Ooe et al., 2006). However, CYP2B1 was hardly detected in the culture of cryo-SHs, and administration of PB could not induce the expression. The present study showed that nuclear receptor CAR expression was repressed when cryo-SHs were cultured in serum-free

conditions, and that the expression was restored by the addition of serum. On the other hand, AhR, PXR, and RXR α were expressed in the cells cultured with or without serum as much as in MHs. AhR and PXR act as transcription factors of CYP1A1 and CYP3A in a ligand-specific manner, respectively (Ma, 2001; Kliewer, 2003). RXR forms a heterodimer with CAR, PXR, or other nuclear receptors in nucleus and can activate the transcription of their target genes (Waxman, 1999). Considering that nuclear receptors other than CAR were expressed in cryo-SHs, it is reasonable that P450s regulated by nuclear receptors other than CAR were maintained in the cells (Ooe et al., 2006).

When a relatively large number of SHs exist in the culture dish, as SHs produce adequate amount of serum proteins, SHs can proliferate in a serum-free medium (Sugimoto et al., 2002; Miyamoto et al., 2005; Ooe et al., 2006). Therefore, whenever SH colonies are replated on new dishes, the cells are routinely cultured in a serum-free medium. In these culture conditions, although many nuclear receptors could be expressed in the cryo-SHs, only CAR disappeared. This loss was reversed by the addition of FBS. The FBS used in the present experiment contained approximately 1 nM total T_3 and 100 nM total T_4 (Table 2). The present study revealed that the effective dosage of T_3 (5×10^{-12} M) or T_4 (5×10^{-12} – 5×10^{-10} M) for the induction of CAR expression was close to the concentrations of free T_3 and free T_4 in the serum of rats (Nauman et al., 1967; Refetoff et al., 1970). We think that the recovery of CAR expression in the experiment was the result of free and protein-conjugated THs contained in FBS. TR α may have a role in CAR expression because this gene was expressed even after cryopreservation and serum did not alter the expression. On the other hand, the expression of TR β in cryo-SHs was lower than that in MHs, and the addition of serum increased its expression only a little. Because CAR expression was restored by serum, TR β might not be directly related to the regulation of the *CAR* gene. Furthermore, T_3 and T_4 possess TH activity in vivo, and T_4 is converted into T_3 by the action of Dio1 in the liver (Köhrle, 2000). Considering that T_4 also had the ability to induce CAR, Dio1 might play a role in this pathway. Induction of CAR expression by T_4 might be caused by the conversion of T_3 by Dio1 expressed in cryo-SHs.

Induction of CYP2B1 mRNA by T_3 was also observed in the cells, and the expression increased with time in culture. Because some SHs in colonies can mature and their hepatic differentiated functions also increase (Sugimoto et al., 2002), it is feasible that CAR and CYP2B1 expression increased with time in culture. A discrepancy was observed in the timing of the expression. The expression of CYP2B1 was delayed compared with that of CAR. The maximum expression of CYP2B1 was observed at day 14 and was approximately 60% of that in MHs, whereas that of CAR was at day 10 and approximately 70% of that in MHs. Our preliminary experiments suggested that the delay of CYP2B1 expression might be because of the nuclear translocation of RXR. By immunostaining, the accumulation of RXR protein was only observed in the nucleus of large-sized SHs, which were matured SHs, not small-sized SHs (data not shown). In our culture conditions, especially without Matrigel, maturation of SHs gradually progresses, and the degree of the maturation in cultured SHs depends on the number of large cells and piled-up cells (Sugimoto et al., 2002). Although T_3 supply induces CAR expression and both CAR and RXR proteins accumulate in the cytoplasm of SHs, the lesser translocation of RXR into nuclei may limit the expression of CYP2B1. Thus, CAR expression might be dependent on the existence of THs, whereas CYP2B1 might require both CAR expression and a differentiated condition of the cells.

Although it is known that CAR is involved in the metabolism of THs (Maglich et al., 2004; Qatanani et al., 2005), there is no report

TABLE 2
Thyroid hormones and TSH level in serum

Sample	Total T ₃ Concentration <i>nM</i>	Total T ₄ Concentration <i>nM</i>	TSH <i>μU/ml</i>
Normal rat	1.34 ± 0.10	53.7 ± 3.9	0.44 ± 0.22
Thyroidectomized rat	0.35 ± 0.05	<3.9	33.07 ± 8.20
PTU-treated rat	<0.15	<3.9	40.77 ± 10.19
Thyroidectomized and PTU-treated rat	<0.15	<0.39	48.16 ± 21.78
Fetal bovine serum	1.06 ± 0.05	101.8 ± 3.6	N.D.

N.D., not determined.




FIG. 7. Expression of CAR and CYP2B1 in thyroidectomized and/or PTU-treated rat livers. Total RNA was prepared from MHs isolated from rats whose thyroids were surgically resected (T) and/or treated with PTU, and quantitative real-time PCR was carried out. The relative expression compared with the normal rat was calculated and standardized with G3PDH expression. Three independent experiments were performed. Bars show S.D., and asterisks indicate significant differences between normal rats and each concentration (*, $p < 0.05$).




FIG. 8. ChIP assays. The genomic DNA-protein complex was isolated from 4-day cultured MHs or 2-week cultured cryo-SH. An anti-TR α antibody was used for immunoprecipitation (IP). A, the promoter region of the rat *ME1* gene containing the TRE sequence was amplified. Cells were cultured with 1×10^{-8} M or without T₃. B, the promoter region of the rat *CAR* gene containing the putative TRE sequence was amplified. This figure shows representative data of four experiments. Cells were cultured with 0, 5×10^{-12} , or 1×10^{-8} M T₃. To detect the sequence, four sets of primers were prepared. The sequences of the forward and reverse primers, annealing temperatures, and the number of reaction cycles are listed in Table 1. Genomic DNA samples used for ChIP assays were also amplified (In).

that THs can affect CAR expression. It is only known that a TRE-like sequence exists in the upstream of the CAR promoter region. In the present experiment, although binding of TR to the TRE sequence in

the promoter region of the *ME1* gene and amplification of the expression by T₃ were confirmed in cryo-SHs and MHs, we could not obtain proof of the direct binding of TR α to the TRE-like sequence in the promoter region of the *CAR* gene. However, it is true that a limited concentration of THs can induce CAR expression in cryo-SHs, and the regulation of CAR expression is not well understood. At present, although we do not know the exact mechanism of CAR expression by THs, unknown indirect mechanisms of regulation may exist in SHs. There is a possibility that the mechanism of *CAR* gene regulation is different between MHs and SHs. To clarify this issue and the characteristics of SHs, we need to perform further experiments.

Although sub-SHs could maintain CAR expression in serum-free culture, cryo-SHs could not. However, cryo-SHs could recover the expression of CAR in the presence of THs, and the expression of other nuclear receptors was maintained in the cryo-SHs. In addition, the duration of cryopreservation (1 week to 1 year) did not affect this phenomenon, and MHs that were cryopreserved by the same procedure as SHs possessed CYP2B1 protein (data not shown). Moreover, cryopreserved human hepatocytes were reported to maintain the activity and the expression of many metabolic enzymes (Shitara et al., 2003; Richert et al., 2006). These results indicate that the loss of the CAR expression may result from the procedure of cryopreservation of SHs. Although it is well known that cryopreservation subjects cells to intensive stress, most damaged functions may be recovered with time if suitable culture conditions are used (Richert et al., 2006). The reason why only CAR was repressed in cryo-SH is not clear. However, this phenomenon may involve the immaturity of SHs. On the other hand, THs seem to function to both maintain CAR expression in MHs and reverse CAR repression in cryo-SHs. When sub-SHs and cryo-SHs were compared, the addition of THs was necessary for re-expression of CAR in cryo-SHs, although it was not necessary for expression of CAR in sub-SHs. Otherwise, an extremely low level of THs in serum caused CAR repression in vivo. This discrepancy may also be caused by the difference between MHs and SHs. Another hypothesis is that enough THs for the expression of CAR are preserved in sub-SHs as protein conjugates and that they are lost as a result of the procedure of cryopreservation. In any case, further investigations concerning the pathway of CAR expression are necessary to clarify the mechanism of TH functions in CAR expression.

Acknowledgments. We thank Minako Kuwano and Yumiko Tsukamoto for technical assistance. We also thank K. Barrymore for help with the manuscript.

References

- Bissell DM and Guzelian PS (1980) Phenotypic stability of adult rat hepatocytes in primary monolayer culture. *Ann N Y Acad Sci* **349**:85–98.
- Brent GA (1994) The molecular basis of thyroid hormone action. *N Engl J Med* **331**:847–853.
- Chesné C, Guyonard C, Fautrel A, Poullain MG, Frémond B, De Jong H, and Guillouzo A (1993) Viability and function in primary culture of adult hepatocytes from various animal species and human beings after cryopreservation. *Hepatology* **18**:406–414.
- de Sousa G, Langouët S, Nicolas F, Lorenzon G, Placidi M, Rahmani R, and Guillouzo A (1996)

- Increase of cytochrome P4501A and glutathione transferase transcripts in cultured hepatocytes from dogs, monkeys and humans after cryopreservation. *Cell Biol Toxicol* **12**:351–358.
- Garcia M, Rager J, Wang Q, Strab R, Hidalgo JJ, Owen A, and Li J (2003) Cryopreserved human hepatocytes as alternative in vitro model for cytochrome p450 induction studies. *In Vitro Cell Dev Biol Anim* **39**:283–287.
- Gonzalez FJ and Gelboin HV (1994) Role of human cytochromes P450 in the metabolic activation of chemical carcinogens and toxins. *Drug Metab Rev* **26**:165–183.
- Guengerich FP and MacDonald TL (1990) Mechanisms of cytochrome P-450 catalysis. *FASEB J* **4**:2453–2459.
- Honkakoski P, Zelko I, Sueyoshi T, and Negishi M (1998) The nuclear orphan receptor CAR-retinoid X receptor heterodimer activates the phenobarbital-responsive enhancer module of the *CYP2B* gene. *Mol Cell Biol* **18**:5652–5658.
- Ikeda S, Mitaka T, Harada K, Sugimoto S, Hirata K, and Mochizuki Y (2002) Proliferation of rat small hepatocytes after long-term cryopreservation. *J Hepatol* **37**:7–14.
- Jackson BA, Davies JE, and Chipman JK (1985) Cytochrome P-450 activity in hepatocytes following cryopreservation and monolayer culture. *Biochem Pharmacol* **34**:3389–3391.
- Kawamoto T, Sueyoshi T, Zelko I, Moore R, Washburn K, and Negishi M (1999) Phenobarbital-responsive nuclear translocation of the receptor CAR in induction of the *CYP2B* gene. *Mol Cell Biol* **19**:6318–6322.
- Kliwer SA (2003) The nuclear pregnane X receptor regulates xenobiotic detoxification. *J Nutr* **133**:2444S–2447S.
- Köhrle J (2000) The deiodinase family: selenoenzymes regulating thyroid hormone availability and action. *Cell Mol Life Sci* **57**:1853–1863.
- Liaw CW and Towle HC (1984) Characterization of a thyroid hormone-responsive gene from rat. *J Biol Chem* **259**:7253–7260.
- Loretz LJ, Li AP, Flye MW, and Wilson AG (1989) Optimization of cryopreservation procedures for rat and human hepatocytes. *Xenobiotica* **19**:489–498.
- Ma Q (2001) Induction of *CYP1A1*. The AhR/DRE paradigm: transcription, receptor regulation, and expanding biological roles. *Curr Drug Metab* **2**:149–164.
- Maglich JM, Watson J, McMillen PJ, Goodwin B, Willson TM, and Moore JT (2004) The nuclear receptor CAR is a regulator of thyroid hormone metabolism during caloric restriction. *J Biol Chem* **279**:19832–19838.
- Mitaka T, Kojima T, Mizuguchi T, and Mochizuki Y (1995) Growth and maturation of small hepatocytes isolated from adult rat liver. *Biochem Biophys Res Commun* **214**:310–317.
- Mitaka T, Mikami M, Sattler GL, Pitot HC, and Mochizuki Y (1992) Small cell colonies appear in the primary culture of adult rat hepatocytes in the presence of nicotine and epidermal growth factor. *Hepatology* **16**:440–447.
- Mitaka T, Sato F, Mizuguchi T, Yokono T, and Mochizuki Y (1999) Reconstruction of hepatic organoid by rat small hepatocytes and hepatic nonparenchymal cells. *Hepatology* **29**:111–125.
- Miyamoto S, Hirata K, Sugimoto S, Harada K, and Mitaka T (2005) Expression of cytochrome P450 enzymes in hepatic organoid reconstructed by rat small hepatocytes. *J Gastroenterol Hepatol* **20**:865–872.
- Nauman JA, Nauman A, and Werner SC (1967) Total and free triiodothyronine in human serum. *J Clin Invest* **46**:1346–1355.
- Ooe H, Kon J, Miyamoto S, Ozone Y, Ninomiya S, and Mitaka T (2006) Cytochrome P450 expression of cultured rat small hepatocytes after long-term cryopreservation. *Drug Metab Dispos* **34**:1667–1671.
- Petty KJ, Morioka H, Mitsuhashi T, and Nikodem VM (1989) Thyroid hormone regulation of transcription factors involved in malic enzyme gene expression. *J Biol Chem* **264**:11483–11490.
- Qatanani M, Zhang J, and Moore DD (2005) Role of the constitutive androstane receptor in xenobiotic-induced thyroid hormone metabolism. *Endocrinology* **146**:995–1002.
- Refetoff S, Robin NI, and Fang VS (1970) Parameters of thyroid function in serum of 16 selected vertebrate species: a study of PBI, serum T4, free T4, and the pattern of T4 and T3 binding to serum proteins. *Endocrinology* **86**:793–805.
- Richert L, Liguori MJ, Abadie C, Heyd B, Mantion G, Halkic N, and Waring JF (2006) Gene expression in human hepatocytes in suspension after isolation is similar to the liver of origin, is not affected by hepatocyte cold storage and cryopreservation, but is strongly changed after hepatocyte plating. *Drug Metab Dispos* **34**:870–879.
- Shitara Y, Li AP, Kato Y, Lu C, Ito K, Itoh T, and Sugiyama Y (2003) Function of uptake transporters for taurocholate and estradiol 17 β -D-glucuronide in cryopreserved human hepatocytes. *Drug Metab Pharmacokinet* **18**:33–41.
- Song MK, Dozin B, Grieco D, Rall JE, and Nikodem VM (1988) Transcriptional activation and stabilization of malic enzyme mRNA precursor by thyroid hormone. *J Biol Chem* **263**:17970–17974.
- Sugimoto S, Mitaka T, Ikeda S, Harada K, Ikai I, Yamaoka Y, and Mochizuki Y (2002) Morphological changes induced by extracellular matrix are correlated with maturation of rat small hepatocytes. *J Cell Biochem* **87**:16–28.
- Sun EL, Aspar DG, Ulrich RG, and Melchior GW (1990) Cryopreservation of cynomolgus monkey (*Macaca fascicularis*) hepatocytes for subsequent culture and protein synthesis studies. *In Vitro Cell Dev Biol* **26**:147–150.
- Swales NJ, Johnson T, and Caldwell J (1996) Cryopreservation of rat and mouse hepatocytes. I. Comparative viability studies. II. Assessment of metabolic capacity using testosterone metabolism. *Drug Metab Dispos* **24**:1224–1230.
- Tateno C and Yoshizato K (1996) Long-term cultivation of adult rat hepatocytes that undergo multiple cell divisions and express normal parenchymal phenotypes. *Am J Pathol* **148**:383–392.
- Waxman DJ (1999) P450 gene induction by structurally diverse xenochemicals: central role of nuclear receptors CAR, PXR, and PPAR. *Arch Biochem Biophys* **369**:11–23.
- Wrighton SA and Stevens JC (1992) The human hepatic cytochromes P450 involved in drug metabolism. *Crit Rev Toxicol* **22**:1–21.

Address correspondence to: Toshihiro Mitaka, Department of Pathophysiology, Cancer Research Institute, Sapporo Medical University School of Medicine, South-1, West-17, Chuo-Ku, Sapporo 060-8556, Japan. E-mail: tmitaka@sapmed.ac.jp
