

ARYL HYDROCARBON RECEPTOR (AHR)-DEPENDENT INDUCTION OF FLAVIN-CONTAINING MONOOXYGENASE mRNAs IN MOUSE LIVER

Trine Celius, Steven Roblin, Patricia A. Harper, Jason Matthews, Paul C. Boutros,
Raimo Pohjanvirta and Allan B. Okey

*Department of Pharmacology and Toxicology, University of Toronto, Toronto, Canada (TC, SR,
PAH, JM, PCB, ABO)*

Research Institute, The Hospital for Sick Children, Toronto, Canada (SR, PAH)

*Department of Food and Environmental Hygiene, University of Helsinki, Helsinki, Finland and
National Public Health Institute, Laboratory of Toxicology, Kuopio, Finland (RP)*

Running title: *FMO induction by dioxin*

Number of pages: 27
Number of tables: 1
Number of figures: 8
Words in Abstract: 249
Words in Introduction: 564
Words in Discussion: 1473
Number of references 34
Supplementary materials: 2 Tables; 5 Figures

Address correspondence to:

Allan B. Okey
Department of Pharmacology and Toxicology
Medical Sciences Building, 1 King's College Circle
University of Toronto, Toronto, Ontario CANADA M5S 1A8
E-mail: allan.okey@utoronto.ca
Phone: (416) 978-7243
FAX: (416) 978-6395

Abbreviations:

AHR, aryl hydrocarbon receptor;
AHRE, aryl hydrocarbon responsive element;
ARNT, aryl hydrocarbon nuclear translocator;
ChIP, chromatin immunoprecipitation;
FMO, flavin-containing monooxygenase;
MMI, methimazole;
TCDD, 2,3,7,8-tetrachlorodibenzo-*p*-dioxin ("dioxin").

ABSTRACT

Flavin-containing monooxygenases (FMOs) are important in detoxication but generally are considered not to be inducible by xenobiotics. Our recent microarray studies revealed induction of FMO2 and FMO3 mRNAs by 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) in liver of mice with wildtype AHR but not in *Ahr*-null mice. The aim of the present study was to delineate mechanisms of FMO regulation. In adult male mice, basal FMO3 mRNA is low but was induced 6-fold at 4 h and 6000-fold at 24 h. The ED₅₀ was about 1 µg/kg for FMO2 & FMO3, similar to the classic AHR-regulated gene, *Cyp1a1*. In adult female mice basal FMO3 mRNA is high and was not induced at 4 h but was elevated 8-fold at 24 h. FMO5 mRNA was significantly downregulated by TCDD in both male and female adults. Juvenile mice show no sex difference in response to TCDD; FMO3 was induced 4-6 fold by TCDD in both sexes. Chromatin immunoprecipitation demonstrated recruitment of AHR and ARNT proteins to *Fmo3* regulatory regions, suggesting that induction by TCDD is a primary AHR-mediated event. Although FMO2 & FMO3 mRNAs were highly induced by TCDD in adult males, overall FMO catalytic activity increased only modestly. In contrast to the striking upregulation of FMO2 & FMO3 in mouse liver, TCDD has little effect on FMO mRNA in rat liver. However, FMO2 & FMO3 mRNAs were highly induced in transgenic mice that express wildtype rat AHR, indicating that lack of induction in rat is not due to an incompetent AHR in this species.

Introduction

Flavin-containing monooxygenases (FMOs) are microsomal enzymes that convert lipophilic substrates to oxygenated metabolites that are more polar and hence more readily-excreted. Substrates include a wide range of nitrogen-containing and sulfur-containing drugs and xenobiotics (reviewed in: (Krueger and Williams, 2005). FMOs are important for metabolism and detoxication of numerous therapeutic agents and environmental toxicants. However FMOs can, in some circumstances, bioactivate particular substrates into reactive metabolites that cause toxicity. There are relatively few known endogenous substrates for FMOs and their physiological functions are poorly understood (Krueger and Williams, 2005; Cashman and Zhang, 2006).

The human genome contains genes encoding five functional FMO proteins – *FMO1-5* – along with six pseudogenes, *FMO6P-11P* (Hernandez et al., 2004; Cashman and Zhang, 2006). *FMO3* is the best-studied human FMO and is expressed at high levels in human liver (Cashman and Zhang, 2006; Phillips et al., 2007). *FMO3* has been identified as the enzyme responsible for oxidation of trimethylamine; mutations in the human *FMO3* gene are responsible for the disease known as trimethylaminuria (fish odor syndrome) (Dolphin et al., 1997; Treacy et al., 1998; Motika et al., 2007; Yeung et al., 2007).

There is considerable variation in tissue-distribution and species-distribution of the different FMOs. In mice there are nine *Fmo* genes, *Fmo1-6* and *Fmo9, 12* and *13* (Hernandez et al., 2004). FMO expression is affected by exogenous factors such as nutritional status (Fu et al., 2004) and by endogenous factors such as age, gender, and hormonal status. Mouse hepatic FMO activity typically is higher in adult females than in adult males, primarily because testosterone suppresses expression of *Fmo1* and *Fmo3* genes (Falls et al., 1997). In female mice, constitutive hepatic *FMO1*, *FMO3* and *FMO5* mRNA levels peak at about 5 weeks of age, then decline somewhat

with further aging. In male mice FMO3 mRNA is maximal at 3-weeks, then is barely detectable after onset of sexual maturity (Janmohamed et al., 2004). Several hormonal and other factors have been identified that regulate developmental expression and tissue-specific expression of FMO mRNAs (Luo and Hines, 2001; Hines, 2006).

Although FMO expression is subject to extensive regulation by endogenous factors, FMOs generally have been considered not to be inducible by xenobiotic chemicals (Krueger and Williams, 2005; Cashman and Zhang, 2006; Hines, 2006). Recent microarray studies in our laboratory revealed substantial induction of FMO2 and FMO3 mRNAs by TCDD in livers of C57BL/6J mice that express wildtype AH receptor but not in *Ahr*-null mice (Tijet et al., 2006). Patel et al. (Patel et al., 2007) also detected induction of FMO3 mRNA by the AHR agonist, β -naphthoflavone, in mouse liver. In contrast to mouse, our expression array experiments in rat liver do not show significant FMO induction by TCDD (Moffat et al., in preparation).

The AHR is a ligand-activated transcription factor and a member of the basic-helix-loop-helix Per (period)-ARNT (aryl hydrocarbon nuclear translocator) SIM (single-minded) (bHLH/PAS) family. Essentially all biochemical and toxic responses to TCDD and related halogenated aromatic hydrocarbons are mediated by the AHR (reviewed in: (Okey, 2007). Upon binding ligand the AHR translocates to the nucleus where it dimerizes with ARNT. The AHR/ARNT complex binds DNA sequences known as aryl hydrocarbon response elements (AHREs) and activates expression of AHR target genes (reviewed in: (Hankinson, 2005; Okey, 2007). The goal of the present work was to further define the AHR-mediated mechanism by which TCDD induces FMOs in mice.

Materials and Methods

Chemicals and reagents. TCDD used in studies in C57BL6/N mice was from Wellington Laboratories (Guelph Ontario). TCDD used in studies in transgenic mice was purchased from the UFA-Oil Institute (Ufa, Russia) and was >99% pure as determined by gas chromatography-mass spectrometry. 5,5'-Dithiobis(2-nitrobenzoate) (DTNB), methimazole (MMI), dithiothreitol (DTT), and NADPH were from Sigma-Aldrich Inc. (St. Louis, MO, USA). Mouse anti-AHR antibody (SA-210) was from BIOMOL (Plymouth Meeting, PA); human anti-ARNT and rabbit anti-IgG antibodies were from Santa Cruz Biotechnology, (Santa Cruz, CA). Power-SYBR Green PCR Master Mix, TaqMan Universal PCR Master Mix, and 96-well real-time PCR plates were from Applied Biosystems, Foster City, CA, USA. All other reagents were of highest quality available from commercial sources.

In vivo experimental design. Juvenile (3-weeks old) or adult (8-weeks old) male and female C57BL6/N mice were obtained from Charles River Laboratories (Montreal, Quebec) and housed under a 12-hour light/dark cycle and given food and water ad libitum. Care and treatment of the mice were in compliance with the guidelines of the Canadian Council on Animal Care and the protocol was approved by the Hospital for Sick Children Animal Care Committee, Toronto, Canada. Mice were injected intraperitoneally with TCDD (30 $\mu\text{g}/\text{kg}$ body weight) dissolved in corn oil vehicle (Tijet et al., 2006) or corn oil alone, and exposed for 2, 4, or 24 h. In the dose-response experiment adult male mice were injected with 0.1, 1.0, or 10 $\mu\text{g}/\text{kg}$ TCDD and exposed for 24 h. Animals were sacrificed by cervical dislocation and livers were snap-frozen and stored at -80°C for ChIP assays and enzyme activity assays, or liver samples were cut into smaller pieces and stored in *RNAlater* for RNA isolation and real-time PCR analysis.

Development of transgenic mice. Constructs for wildtype rat AHR ligated to an EF-1 α 1 promoter were isolated from plasmids and microinjected into fertilized mouse oocytes with the C57BL/6J01aHsd strain as recipient. The endogenous mouse AHR was then eliminated by selective crossing with *Ahr*-null mice. Expression of the transgene was confirmed by measurement of its mRNA (by real-time RT-PCR) and protein (by immunoblotting). The rat AHR transgene fully supports induction, by TCDD, of the classic AHR-regulated mouse genes, *Cyp1a1* and *Cyp1a2* in liver and is expressed there at levels some 5 times higher than the endogenous wildtype AHR in C57BL/6 mice. The procedure will be described in detail in a later paper (Pohjanvirta, in preparation).

Real-time reverse transcription (RT) quantitative Polymerase Chain Reaction (PCR).

Probes and primers for mouse FMO1 (NM010231), FMO2 (NM018881), FMO3 (NM008030), FMO4 (NM144878), and FMO5 (NM010232) were designed using the Primer3 program (http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi) (Rozen, 2000). Liver total RNA was isolated using GE Healthcare Spin Mini kit. Real-time RT-PCR was performed using 1 μ g of total RNA and random hexamer primers by a standard procedure. Two μ L of RT-reaction was used for each real-time qPCR reaction. Each reaction mixture (25 μ L) contained optimized probe and primer concentrations as well as the TaqMan universal mastermix. Expression levels of target mRNA were normalized to levels of 18S rRNA and analyzed using the comparative C_T ($\Delta\Delta C_T$) method. Primer sequences are given in Supplementary Table 1.

In Vivo Chromatin Immunoprecipitation (ChIP) assay. ChIP assays were performed according to Matthews et al. (Matthews et al., 2005) except that 100 to 120 mg of snap-frozen liver was homogenized and cross-linked in 1% formaldehyde and the DNA was isolated using a PCR purification kit (BioBasic) and eluted in 50 μ L. ChIP DNA (1 μ L) was amplified by PCR

with primers 5'AAGCCAAGCCAGAAAATCAA3' and 5' TCGAGGAACAGAGTGCAATG3' for the mouse *Fmo3* AHRE (-3183 to -3260 relative to the transcriptional start site). For real-time PCR, Power SYBR green PCR master mix was used to amplify the DNA fragments.

Preparation of hepatic microsomes. Liver samples (100 mg) were homogenized with five volumes of ice-cold homogenization buffer containing 0.1 mM potassium phosphate, pH 7.5, with 0.1 mM EDTA and 1.15% KCl. The homogenate was centrifuged at $10,000 \times g$ for 15 min at 4° C in a refrigerated centrifuge. The supernatant was centrifuged at $100,000 \times g$ for 60 min at 4° C. The firmly packed microsomal pellets were resuspended by homogenization in 0.1 mM potassium phosphate, pH 7.5, with 0.1mM EDTA and 0.25 M sucrose. Microsomes were stored at -80° until use. Microsomal protein concentration was measured using Bradford's method (BioRad kit) with BSA as the standard protein.

Enzyme activity assay. FMO activity was measured by monitoring methimazole oxidation as described by Dixit and Roche (Dixit and Roche, 1984). Sample and reference cuvettes contained a 1 mL mixture of 0.1 M Tricine/KOH, pH 8.5, 0.1 mM EDTA, 0.06 mM DTNB (in 0.1 M calcium phosphate buffer, pH 8.0), 0.025 mM DTT, 0.1 mM NADPH, 100 µg microsomal proteins, and 2.0 mM MMI. Briefly, tricine/EDTA buffer, NADPH and microsomes were mixed and incubated at room temperature for 40 sec. DTT, DTNB, and MMI (in sample cuvette only) were added and the reaction was incubated at 37°C in the chamber of a recording dual-beam spectrophotometer (Beckman DU-65). Methimazole was added to the sample cuvette and absorbance was recorded for 10 min at 412 nm. The reaction mixture excluding methimazole was used as reference.

Statistics. Statistical significance was tested using two-tailed unpaired t-tests with Welch's adjustment for heteroscedasticity (Microsoft Excel 2002).

Results

Recent microarray studies from our laboratories showed induction of FMO2 and FMO3 mRNAs in livers of C57BL/6J mice exposed to TCDD but not in livers from AHR knock-out (*Ahr*^{-/-}) mice (Tijet et al., 2006). To determine the breadth of the FMO response to TCDD, in the current project we measured mRNA levels for the five most prominent FMOs in mouse liver. As recently reported by Siddens et al. (Siddens et al., 2008), the most abundant hepatic mRNA under basal (untreated) conditions in adult C57BL/6J mice is FMO5. At a dose of 30 µg/kg for 24 h, TCDD significantly suppressed FMO5 mRNA levels both in male (Fig. 1) and in female mice (Fig. 2). In contrast, mRNA levels for FMO1, FMO2 and FMO3 are low in untreated male mice but each of these FMOs was highly induced by TCDD; indeed, the magnitude of mRNA induction for FMO3 in adult males was similar to that of the prototypical TCDD-inducible/AHR-regulated gene, *Cyp1a1* (Fig. 1). In female mice FMO1 and FMO2 mRNA levels were unaffected by 24 h exposure to TCDD whereas FMO3 mRNA was significantly induced and FMO4 was significantly suppressed by TCDD (Fig. 2).

To further characterize induction of FMOs we obtained dose-response curves for TCDD-induction of FMO2 and FMO3 mRNAs and found that the approximate ED₅₀ values for induction of FMO2 and FMO3 are similar to that for CYP1A1 in male mice (Fig. 3). In livers of male mice, induction of FMO2 and FMO3 mRNAs is significant within 2 h after TCDD injection and the magnitude of induction increases further at 4 h (Fig. 4). In male mice the magnitude of FMO3 induction is dramatically elevated at 24 h, achieving mRNA levels about 6000-fold higher than in control mice (Fig. 5A).

In adult mice there is a notable sex difference in constitutive expression of FMO3 mRNA in liver where mRNA levels in untreated animals are about 500-fold to 2300-fold higher in females than in males (Figs. 5A & 5B). The reason why the constitutive level is 2300-fold higher in females in the experiment shown in Fig. 5A but only 500-fold higher in the experiment shown in Fig. 5B is not known. However, the experiment shown in Fig. 5B was conducted several months after the experiment shown in Fig. 5A; the difference may reflect the sensitivity of basal FMO expression to many factors such as season and nutritional factors. The difference is not due to diurnal variation in basal expression because tissues in all experiments were harvested at a similar time of day – late morning. At 4 h exposure to TCDD there is no significant induction of FMO3 mRNA in females (Fig. 5B) but FMO3 mRNA is induced about 6-fold in males (Fig. 5A). After 24 h exposure to TCDD, FMO3 mRNA in female mice does respond with an induction about 8-fold (Fig. 5B). In juvenile mice (3-weeks old) basal levels of FMO3 mRNA are 15-fold higher in females than in males but males and females exhibit a similar magnitude of FMO3 induction, about 4-fold, at this age (Fig. 6). Constitutive levels of FMO3 mRNA are higher in juveniles than in adults for both male (36-fold) and female (26-fold) (data not shown).

As described above, our previous experiments in *Ahr*-null mice indicated that induction of FMO2 and FMO3 was AHR-dependent (Tijet et al., 2006). To gain further insight into the mechanism of induction we performed ChIP assays to determine if TCDD could evoke binding of the AHR and its dimerization partner, ARNT, to an enhancer region in the *Fmo3* gene. Within 2 h after TCDD injection, both the AHR protein and the ARNT protein were recruited to the region -3260 to -3184 (relative to the transcription start site) that contains an AH responsive element (AHRE); this TCDD-induced signal persisted at least up to 4 h (Fig. 7). There were no obvious differences in recruitment of AHR or ARNT between male and female mice (Fig. 7).

Although mRNAs for FMO1, FMO2 and FMO3 are highly upregulated by TCDD in mice, as shown in our current study and previous study (Tijet et al., 2006), we have found that in liver of TCDD-sensitive male rats carrying wildtype AHR, TCDD has negligible effects on FMO2 and FMO3 (Moffat et al. in preparation). There are substantial differences in AH receptor structure between C57BL/6J mice and wildtype rat AHR (Moffat et al., 2007b; Okey, 2007). In order to determine if these differences underlie the lack of FMO induction in rats we measured FMO2 and FMO3 mRNA levels in TCDD-treated transgenic mice that express wildtype rat AHR in a C57BL/6J background. As shown in Figure 8, the wildtype rat AHR fully supports induction of the prototypical AHR-regulated gene, *Cyp1a1* as well as FMO2 and FMO3 mRNAs in these transgenic animals. This demonstrates that the failure of TCDD to upregulate FMOs in rats is not the result of an inherent inability of wildtype rat AHR to regulate FMO genes.

We attempted to determine if the large elevation of FMO mRNAs in TCDD-treated mice led to a corresponding increase in FMO proteins by performing immunoblotting with antibodies that had been raised against human FMO3. Due to lack of specificity and high cross-reactivity with other proteins it was not possible to distinguish FMO3 from other protein bands (data not shown). Unfortunately, suitable antibodies are currently not available for immunologic characterization of mouse FMO proteins (Siddens et al., 2008). Thus, as an alternative index to the effect of TCDD on FMO protein, we measured FMO catalytic activity with methimazole, a substrate that is metabolized by multiple species of FMO (Zhang et al., 2007). Metabolism of methimazole was modestly but significantly increased in liver microsomes from TCDD-treated male mice but not in microsomes from female mice (Table 1).

One possible explanation for the modest increase in methimazole metabolism despite very high mRNA induction is that TCDD might inhibit catalytic function of FMO enzymes by

occupying active sites on the enzyme proteins. There is precedent for such a mechanism in the cytochrome P450 field where TCDD has been shown to bind extensively to CYP1A2 protein and inhibit CYP1A2-mediated enzyme activity (Diliberto et al., 1997; Chen et al., 2003; Staskal et al., 2005). Thus we tested the possibility that TCDD might inhibit FMO activity by adding TCDD directly to microsomal preparations in the methimazole-metabolism experiments. At a concentration of 1 μ M, TCDD had no effect on metabolism of methimazole (data not shown). Therefore the low magnitude of the increase in FMO activity in TCDD-treated animals cannot be attributed to an inhibitory effect of TCDD on FMO enzyme function.

In addition to liver, we also assessed possible induction by TCDD (30 μ g/kg for 24 h) of FMO2 and FMO3 mRNA in lung, kidney, small intestine, brain and thymus of adult male mice and did not observe significant induction in any of these tissues (data not shown).

Discussion

FMOs are subject to regulation by hormones (Falls et al., 1997; Janmohamed et al., 2004) and nutritional status (Fu et al., 2004) but usually are considered not to be inducible by xenobiotic chemicals (Krueger and Williams, 2005; Cashman and Zhang, 2006; Hines, 2006; Phillips et al., 2007). However, we found that in mouse liver, especially in males, TCDD causes very large increases in mRNA levels for FMO1, FMO2 and FMO3. The AHR plays an essential role in this upregulation as evidenced by our current ChIP assays and our previous experiments in *Ahr*-null versus wildtype mice which showed that the AHR is required for induction of FMO2 and FMO3 mRNAs (Tijet et al., 2006).

Prolonged feed restriction has been reported to increase FMO3 mRNA levels in mice (Fu et al., 2004). Since TCDD is well-known to reduce food intake in dioxin-susceptible rodents (Pohjanvirta and Tuomisto, 1994) it might be argued that FMO upregulation is a secondary response to reduced food intake rather than a direct, AHR-mediated event. Our current study indicates that upregulation of FMO3 mRNA is direct, via a rapid TCDD-induced recruitment of the AHR and ARNT proteins to an AHRE-containing enhancer region between -3260 to -3184 in the *Fmo3* gene. Induction of FMO3 mRNA follows a pattern very similar to that of the classic AHR-regulated gene, *Cyp1a1* in that: (1) the dose-response curves for their induction by TCDD are similar; (2) their induction is abolished in *Ahr*-null mice; (3) their induction is accompanied by recruitment of AHR and ARNT proteins to AHREs in upstream enhancer regions; (4) their induction occurs within a few hours after TCDD administration *in vivo*.

The dramatic elevation of FMO mRNA, particularly of FMO3 in male mice, is accompanied by only a modest increase in FMO catalytic activity measured with methimazole. Methimazole was reported to be metabolized by FMO1 and FMO3 but not FMO5 when mouse FMOs are

expressed as fusion proteins in *E. coli* (Zhang et al., 2007). However, selectivity of substrates for different forms of FMO in various animal species in vivo remains poorly defined (Yeung and Rettie, 2006). This creates difficulties in the FMO field because, in contrast to P450 enzymes, there is a general lack of "probe" substrates that are reliable indicators of catalytic function for specific forms of FMO.

It is possible that the abundant FMO mRNAs in TCDD-treated mice are not able to be effectively translated into their respective proteins. Numerous microRNAs are predicted to target mouse FMO2 and mouse FMO3 (Supplementary Table 2). We previously tested the effect of TCDD on expression of microRNAs in rodent livers (Moffat et al., 2007a). Out of the list of microRNAs that are predicted to target mouse FMO3 mRNA, we found that TCDD treatment caused about a 1.2-fold increase in levels of hsa-miR-519 in livers of C57BL/J mice. In addition, mmu-miR203 and mmu-miR-148b each are increased about 1.5-fold by TCDD in an AHR-dependent manner (Moffat et al., 2007a); these latter two miRNAs are closely related in sequence to miRNAs (mmu-miR-203* and mmu-148a*) that are predicted to target mouse FMO3 mRNA. It is conceivable that the concerted increase in levels of these three microRNAs in TCDD-treated mice interferes with translation of FMO3 mRNA into protein. Our previous microRNA study (Moffat et al., 2007a) also showed a 1.2-fold increase in hepatic levels of mmu-miR-203 in TCDD-treated mice; miR-203 is predicted to target mouse FMO2 mRNA, again providing a potential explanation for the relatively low increase in FMO catalytic activity when compared with the large increase in FMO2 mRNA in TCDD-treated mice.

Mouse and rat are the two most extensively-studied laboratory species in biochemistry, pharmacology and toxicology. In regard to FMOs, there are notable differences in regulation between mouse and rat. As mentioned earlier, our initial expression array studies (Tijet et al.,

2006) and the results of the current study show that FMO2 and FMO3 mRNAs are strongly upregulated by TCDD in mouse liver but this is not the case in rat liver (Moffat et al. in preparation).

Moreover, for FMO1 the effect of TCDD is opposite in the two rodent species: FMO1 is significantly upregulated in mouse liver (this study and (Boutros et al., 2008) and significantly downregulated in rat liver (Boutros et al., 2008). Our experiments in mice that express transgenic wildtype AHR show that the rat AHR is fully competent to drive induction of FMO2 and FMO3 in mice, indicating that the lack of FMO induction in rat resides in differences in non-AHR regulatory factors between rat and mouse, perhaps in the *FMO* gene structures themselves. Our current ChIP studies in mice revealed binding of AHR/ARNT to an AHRE in an upstream enhancer region of the mouse *Fmo3* gene. While neither the rat nor the mouse *Fmo3* gene contain full-length AHRE-I motifs in their up-stream regulatory regions, both contain extended AHRE-I motifs, as well AHRE-II and ARE motifs (Supplementary Figures 1-4). Very recently Klick et al. (Klick et al., 2008) showed that structural differences in responsive elements in the promoter region can account for some differences among mammalian species in hepatic constitutive FMO3 expression. Although structural differences in AH response elements potentially could explain the fact that that FMO3 mRNA levels are highly upregulated by TCDD in mouse but not in rat, our analysis of AHREs in the two species indicates that the difference in FMO inducibility cannot be explained solely by differences in regulatory motifs between their *FMO3* genes. (Supp Figs. 1-4). Instead, epigenetic factors such as DNA methylation patterns, nucleosome placements, histone modifications, or the presence/absence of interfering or obligate partner proteins may be involved.

It is not known whether TCDD can induce hepatic FMO3 in humans. We aligned the mouse AHR-associated region with human and rat genomes by a BLAT analysis (Kent, 2002); see Suppl Fig. 5). The mouse AHR-associated region showed minimal overlap with genomic sequences in human (26 of 78 base-pairs aligning) or in rat (20 of 78 base-pairs aligning). Further, the aligned regions are on separate chromosomes from the *FMO3* gene in each species. Taken together these results suggest that the AHR-binding-element is part of a mouse-specific regulatory module and, at present, it cannot be inferred from promoter analysis in silico whether human FMO3 is likely to be inducible by AHR agonists.

In rodents there are substantial differences between male and female in FMO expression. In mice, previous studies (Falls et al., 1997; Cherrington et al., 1998; Janmohamed et al., 2004) as well as our current experiments show that constitutive FMO3 is highly expressed in liver of adult female mice but is virtually absent in livers of adult male mice. However, in juvenile animals, males express significant basal levels of FMO3 mRNA and TCDD upregulates FMO3 to a similar extent in both sexes (Fig. 6). In mice, testosterone suppresses hepatic FMO3 and FMO1 (Falls et al., 1997). Surprisingly, the sex differences and response to sex steroids in rats are opposite to those in mice. In rats FMO3 expression is independent of the sex (Cherrington et al., 1998) and treatment with testosterone elevates FMO levels rather than suppressing them (Dannan et al., 1986). One way to view the effect of TCDD on FMO3 expression in mice is that TCDD overcomes the suppressive action of testosterone. Female mouse livers express high constitutive levels of FMO3 that could not be further elevated by TCDD after 2-4 h whereas FMO3 mRNA levels in male livers were induced 6-fold upon exposure to TCDD for 4 h. After 24 h exposure male liver FMO3 expression levels were increased by 6000-fold and achieved female-like levels of FMO3 mRNA. Female constitutive FMO3 expression levels were high and increased only by 8-fold upon TCDD exposure for 24 h. The mechanism by which testosterone negatively regulates FMO3 in mice is not known. The suppression in FMO3 mRNA levels seen

in adult male mouse liver were not observed in kidney and lung, indicating that FMO3 is subject to tissue-specific regulation (Janmohamed et al., 2004) in addition to apparent species-specific regulation. FMO enzyme activity was higher in female liver microsomes than in microsomes from male mice in our study, both in the constitutive state and after TCDD treatment.

In summary, our present study shows that multiple FMO mRNAs in mice can, indeed, be upregulated by a xenobiotic chemical, TCDD. This induction is mediated by the AH receptor in a fashion similar to that for the classic AHR-regulated gene, *Cyp1a1*. Further analytical tools, including antibodies that are selective to individual mouse FMO isoforms and substrates that can discriminate among the isoforms will be necessary in order to fully reveal the downstream consequences of the profound upregulation at the mRNA level.

Acknowledgements

We thank Sharon Choi, Laura MacPherson and Agnes Forgacs for help with harvesting tissues and Ulla Naukkarinen, Janne Korkalainen and Arja Tamminen for technical assistance, Dr. Ivy Moffat for microRNA data and Dr. Sharon Krueger & Dr. David Williams (Oregon State University) for advice regarding FMO enzyme assays.

References

- Boutros PC, Yan R, Pohjanvirta R and Okey AB (2008) Transcriptomic responses to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in liver: comparison of rat and mouse. *BMC Genomics* **in press**.
- Cashman JR and Zhang J (2006) Human flavin-containing monooxygenases. *Annu Rev Pharmacol Toxicol* **46**:65-100.
- Chen JJ, Chen GS and Bunce NJ (2003) Inhibition of CYP 1A2-dependent MROD activity in rat liver microsomes: an explanation of the hepatic sequestration of a limited subset of halogenated aromatic hydrocarbons. *Environ Toxicol* **18**:115-119.
- Cherrington NJ, Cao Y, Cherrington JW, Rose RL and Hodgson E (1998) Physiological factors affecting protein expression of flavin-containing monooxygenases 1, 3 and 5. *Xenobiotica* **28**:673-682.
- Dannan GA, Guengerich FP and Waxman DJ (1986) Hormonal regulation of rat liver microsomal enzymes. Role of gonadal steroids in programming, maintenance, and suppression of delta 4-steroid 5 alpha-reductase, flavin-containing monooxygenase, and sex-specific cytochromes P-450. *J Biol Chem* **261**:10728-10735.
- Diliberto JJ, Burgin D and Birnbaum LS (1997) Role of CYP1A2 in hepatic sequestration of dioxin: studies using CYP1A2 knock-out mice. *Biochem Biophys Res Commun* **236**:431-433.
- Dixit A and Roche TE (1984) Spectrophotometric assay of the flavin-containing monooxygenase and changes in its activity in female mouse liver with nutritional and diurnal conditions. *Arch Biochem Biophys* **233**:50-63.
- Dolphin CT, Janmohamed A, Smith RL, Shephard EA and Phillips IR (1997) Missense mutation in flavin-containing mono-oxygenase 3 gene, FMO3, underlies fish-odour syndrome. *Nat Genet* **17**:491-494.
- Falls JG, Ryu DY, Cao Y, Levi PE and Hodgson E (1997) Regulation of mouse liver flavin-containing monooxygenases 1 and 3 by sex steroids. *Arch Biochem Biophys* **342**:212-223.
- Fu C, Xi L, Wu Y, McCarter R, Richardson A, Hickey M and Han ES (2004) Hepatic genes altered in expression by food restriction are not influenced by the low plasma glucose level in young male GLUT4 transgenic mice. *J Nutr* **134**:2965-2974.
- Hankinson O (2005) Role of coactivators in transcriptional activation by the aryl hydrocarbon receptor. *Arch Biochem Biophys* **433**:379-386.
- Hernandez D, Janmohamed A, Chandan P, Phillips IR and Shephard EA (2004) Organization and evolution of the flavin-containing monooxygenase genes of human and mouse: identification of novel gene and pseudogene clusters. *Pharmacogenetics* **14**:117-130.
- Hines RN (2006) Developmental and tissue-specific expression of human flavin-containing monooxygenases 1 and 3. *Expert Opin Drug Metab Toxicol* **2**:41-49.
- Janmohamed A, Hernandez D, Phillips IR and Shephard EA (2004) Cell-, tissue-, sex- and developmental stage-specific expression of mouse flavin-containing monooxygenases (Fmos). *Biochem Pharmacol* **68**:73-83.
- Kent WJ (2002) BLAT--the BLAST-like alignment tool. *Genome Res* **12**:656-664.
- Klick DE, Shadley JD and Hines RN (2008) Differential regulation of human hepatic flavin containing monooxygenase 3 (FMO3) by CCAAT/enhancer-binding protein beta

- (C/EBPbeta) liver inhibitory and liver activating proteins. *Biochem Pharmacol* **76**:268-278.
- Krueger SK and Williams DE (2005) Mammalian flavin-containing monooxygenases: structure/function, genetic polymorphisms and role in drug metabolism. *Pharmacol Ther* **106**:357-387.
- Luo Z and Hines RN (2001) Regulation of flavin-containing monooxygenase 1 expression by ying yang 1 and hepatic nuclear factors 1 and 4. *Mol Pharmacol* **60**:1421-1430.
- Matthews J, Wihlen B, Thomsen J and Gustafsson JA (2005) Aryl hydrocarbon receptor-mediated transcription: ligand-dependent recruitment of estrogen receptor alpha to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-responsive promoters. *Mol Cell Biol* **25**:5317-5328.
- Moffat ID, Boutros PC, Celius T, Linden J, Pohjanvirta R and Okey AB (2007a) microRNAs in Adult Rodent Liver Are Refractory to Dioxin Treatment. *Toxicol Sci* **99**:470-487.
- Moffat ID, Roblin S, Harper PA, Okey AB and Pohjanvirta R (2007b) Aryl hydrocarbon receptor splice variants in the dioxin-resistant rat: tissue expression and transactivational activity. *Mol Pharmacol* **72**:956-966.
- Motika MS, Zhang J and Cashman JR (2007) Flavin-containing monooxygenase 3 and human disease. *Expert Opin Drug Metab Toxicol* **3**:831-845.
- Okey AB (2007) An Aryl Hydrocarbon Receptor Odyssey to the Shores of Toxicology: The Deichmann Lecture, International Congress of Toxicology-XI. *Toxicol Sci* **98**:5-38.
- Patel RD, Hollingshead BD, Omiecinski CJ and Perdew GH (2007) Aryl-hydrocarbon receptor activation regulates constitutive androstane receptor levels in murine and human liver. *Hepatology* **46**:209-218.
- Phillips IR, Francois AA and Shephard EA (2007) The flavin-containing monooxygenases (FMOs): genetic variation and its consequences for the metabolism of therapeutic drugs. *Current Pharmacogenomics* **5**:292-313.
- Pohjanvirta R and Tuomisto J (1994) Short-term toxicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in laboratory animals: effects, mechanisms, and animal models. *Pharmacol Rev* **46**:483-549.
- Rozen S, Skaletsky, H. (2000) *Primer3 on the WWW for general users and for biologist programmers*. Humana Press, Totawa, NJ.
- Siddens LK, Henderson MC, Vandyke JE, Williams DE and Krueger SK (2008) Characterization of mouse flavin-containing monooxygenase transcript levels in lung and liver, and activity of expressed isoforms. *Biochem Pharmacol* **75**:570-579.
- Staskal D, Diliberto J, Devito M and Birnbaum L (2005) Inhibition of Human and Rat CYP1A2 by TCDD and Dioxin-like Chemicals. *Toxicol Sci*.
- Tijet N, Boutros PC, Moffat ID, Okey AB, Tuomisto J and Pohjanvirta R (2006) The aryl hydrocarbon receptor regulates distinct dioxin-dependent and dioxin-independent gene batteries. *Mol Pharmacol* **69**:140-153.
- Treacy EP, Akerman BR, Chow LM, Youil R, Bibeau C, Lin J, Bruce AG, Knight M, Danks DM, Cashman JR and Forrest SM (1998) Mutations of the flavin-containing monooxygenase gene (FMO3) cause trimethylaminuria, a defect in detoxication. *Hum Mol Genet* **7**:839-845.
- Yeung CK, Adman ET and Rettie AE (2007) Functional characterization of genetic variants of human FMO3 associated with trimethylaminuria. *Arch Biochem Biophys* **464**:251-259.
- Yeung CK and Rettie AE (2006) Prochiral sulfoxidation as a probe for flavin-containing monooxygenases. *Methods Mol Biol* **320**:163-172.

Zhang J, Cerny MA, Lawson M, Mosadeghi R and Cashman JR (2007) Functional activity of the mouse flavin-containing monooxygenase forms 1, 3, and 5. *J Biochem Mol Toxicol* **21**:206-215.

Footnotes

This work was supported by grant MOP-57903 from the Canadian Institutes of Health to A.B.O. and grant 123345 from the Academy of Finland to R.P.

Address correspondence to:

Allan B. Okey

Department of Pharmacology & Toxicology

Medical Sciences Building, 1 King's College Circle

University of Toronto

Toronto, Ontario CANADA M5S 1A8

Figure Legends

FIGURE 1. **Male adult mice: FMO mRNA levels in livers of TCDD-treated versus control.**

Mice were injected with TCDD (30 $\mu\text{g}/\text{kg}$) and livers removed 24 h later. mRNA levels for FMO1-5 were measured by real-time RT-PCR as described in Materials and Methods. mRNA levels are expressed relative to the level of FMO3 mRNA in male control mice set at 1.0. ** $p < 0.01$ compared with vehicle control. Plotted bars in all figures represent mean \pm SD; $n = 4$ mice per group.

FIGURE 2. **Female adult mice: FMO mRNA levels in livers of TCDD-treated versus control.**

Female mice were injected with TCDD (30 $\mu\text{g}/\text{kg}$) and livers removed 24 h later. mRNA levels were measured as described for male mice in the legend to Fig. 1 and are expressed relative to the level of FMO3 mRNA in female control mice set at 1.0. ** $p < 0.01$ compared with vehicle control; $n = 4$ mice per group.

FIGURE 3. **Dose-dependent induction of mRNAs for FMO2, FMO3 and CYP1A1.**

Adult male mice were injected with TCDD at doses of 0.1, 1.0 or 10 $\mu\text{g}/\text{kg}$. Liver was harvested 24 h later and mRNA levels were measured by real-time RT-PCR. Results are plotted as percent of the maximal induced response for that particular mRNA. Error bars represent SD; $n = 3$ mice per group. Curves were fitted with GraphPad Prism 4.

FIGURE 4. **Time-dependent induction of FMO2 and FMO3 mRNAs in male mice.**

Mice were exposed to TCDD (30 $\mu\text{g}/\text{kg}$) for 2 h or 4 h. mRNA levels were measured by real-time RT-PCR and are expressed relative to the mRNA level in control mice set at 1.0. ** $p < 0.01$; $n = 3$ mice per group.

FIGURE 5. Sex-dependent induction of FMO3 mRNA. Adult male and female mice were exposed to TCDD (30 $\mu\text{g}/\text{kg}$) for 4 h (A) or for 24 h (B). FMO3 mRNA was measured by real-time RT-PCR and levels are expressed relative to the level in control male mice set at 1.0. Note the difference in scales of the y-axes for male vs. female mice and that the induction of FMO3 after TCDD exposure in females at 4h is not significant (n.s.) relative to the female control set at 1.0. ** $p < 0.01$ compared with vehicle control (n = 4).

FIGURE 6. Lack of sex difference in magnitude of FMO3 induction in juvenile mice. Three-week-old male and female mice were exposed to TCDD (30 $\mu\text{g}/\text{kg}$) for 4 h. FMO3 mRNA was measured by real-time RT-PCR and levels are expressed relative to the level in control male mice set at 1.0. Note the difference in scales of the y-axes for male vs. female mice. * $p < 0.05$; ** $p < 0.01$ compared with vehicle control (n = 4).

FIGURE 7. Recruitment of AHR and ARNT to *Fmo3* enhancer region (AHRE) after TCDD treatment. Adult male and female mice were exposed to TCDD (30 $\mu\text{g}/\text{kg}$) or the corn oil vehicle control (CO) for 2 h or 4 h prior to harvesting of liver. ChIP assays were performed as described in Materials and Methods with antibodies to AHR and ARNT as well as IgG (as a negative control). Primer pairs were designed to encompass the AHRE-containing region (-3183 to -3260) shown in the diagram. Results are reported as percent enrichment of AHR and ARNT binding to this region out of the total input. * $p < 0.05$; ** $p < 0.01$ compared with vehicle control (n = 4).

FIGURE 8. FMO induction in mice that express a rat wildtype AHR transgene. Mice with the rat wildtype AHR transgene were treated with TCDD (5 $\mu\text{g}/\text{kg}$) and liver was harvested 24 h later. mRNA levels for FMO2, FMO3 and CYP1A1 are given relative to the level in control

mice for that particular gene set at 1.0. Bars for control FMO3 and control CYP1A1 are not visible when plotted on this scale. ** $p < 0.01$ compared with vehicle control (n = 4).

TABLE 1

FMO enzyme activity in TCDD-treated versus control mice.

FMO catalytic activity was measured in liver microsomes from TCDD-treated (30 $\mu\text{g}/\text{kg}$; 24 h) or control mice using methimazole as substrate as described in Experimental Procedures. Results are reported as $\mu\text{mol}/\text{min}/\text{mg} \pm \text{SD}$ * $p < 0.05$ compared to with control.

	Control	TCDD
Male	3.75 \pm 0.93	5.13 \pm 0.99*
Female	11.42 \pm 2.34	11.14 \pm 4.06




Fig 1




Fig 2




Fig 3




Fig 4

Adult mice




Fig 5






Fig 6

A**B****Fig 7**

Transgenic mice that express wildtype rat AHR




Fig 8