

DMD # 80952

Hepatic Abundance and Activity of Androgen and Drug Metabolizing Enzyme, UGT2B17, are Associated with Genotype, Age, and Sex

Deepak Kumar Bhatt, Abdul Basit, Haeyoung Zhang, Andrea Gaedigk, Seung-been Lee, Katrina G. Claw, Aanchal Mehrotra, Amarjit Singh Chaudhry, Robin E. Pearce, Roger Gaedigk, Ulrich Broeckel, Timothy A. Thornton, Deborah A. Nickerson, Erin G. Schuetz, John K. Amory, J. Steven Leeder, and Bhagwat Prasad

Department of Pharmaceutics, University of Washington, Seattle, WA, USA (D.K.B., A.B., H.Z., K.G.C., A.M., B.P.)

Division of Pediatric Pharmacology and Medical Toxicology, Department of Pediatrics, Children's Mercy Hospitals and Clinics, Kansas City, MO (A.G., R.E.P., R.G., J.S.L.)

Department of Genome Sciences, University of Washington, Seattle, WA (S.L., D.A.N.)

Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN (A.S.C., E.G.S.)

Section of Genomic Pediatrics, Department of Pediatrics, and Human and Molecular Genetics Center, Medical College of Wisconsin, Milwaukee, WI (U.B.)

Department of Biostatistics, University of Washington, Seattle, WA (T.A.T.).

Department of Medicine, University of Washington, Seattle, WA (J.K.A.)

DMD # 80952

Running title: Interindividual variability in hepatic UGT2B17

Corresponding author: Bhagwat Prasad, Ph.D.; Department of Pharmaceutics, University of Washington, 1959 NE Pacific Street, Seattle, WA 98195, Phone: (206) 221-2295, Fax: (206) 543-3204; E-mail: bhagwat@uw.edu

Word limit

Number of text pages: 15

Number of tables: 2

Number of figures: 3

Number of references: 49

Number of words in Abstract: 209

Number of words in Introduction: 624

Number of words in Discussion: 1356

Abbreviations: ABB, ammonium bicarbonate buffer; BSA, bovine serum albumin; CYPs, cytochrome P450 enzymes; DTT, dithiothreitol; FPKM, fragments per kilobase per million reads; HLM, human liver microsomes; HSA, human serum albumin; IAA, iodoacetamide; LC-MS/MS, liquid chromatography-tandem mass spectrometry; LD, linkage disequilibrium; LOD, limit of detection; LLOQ, lower limit of quantification; PGRN, Pharmacogenomics Research Network; SNPs, single nucleotide polymorphisms; T, testosterone; DHT, dihydrotestosterone.

DMD # 80952

ABSTRACT

The major objective of this study was to investigate association of genetic and non-genetic factors with variability in protein abundance and in vitro activity of the androgen metabolizing enzyme, UGT2B17, in human liver microsomes (n=455). UGT2B17 abundance was quantified by LC-MS/MS proteomics and enzyme activity was determined by using testosterone and dihydrotestosterone as in vitro probe substrates. Genotyping or gene resequencing and mRNA expression were also evaluated. Multivariate analysis was used to test association of UGT2B17 copy number variation, single nucleotide polymorphisms (SNPs), age and sex with its mRNA expression, abundance and activity. UGT2B17 gene copy number and SNPs (rs7436962, rs9996186, rs28374627 and rs4860305) were associated with gene expression, protein levels and androgen glucuronidation rates in a gene-dose dependent manner. UGT2B17 protein (mean \pm SD pmol per mg microsomal protein) is sparsely expressed in children below 9 years (0.12 ± 0.24), but profoundly increases from age 9 years to adults (~10 fold) with ~2.6-fold higher abundance in males than females (1.2 vs. 0.47). Association of androgen glucuronidation with UGT2B15 abundance was only observed in the low UGT2B17 expressers. These data can be used to predict variability in the metabolism of UGT2B17 substrates. Drug companies should include UGT2B17 in early phenotyping assays during drug discovery to avoid late clinical failures.

DMD # 80952

INTRODUCTION

Uridine 5'-diphospho-glucuronosyltransferases (UGTs; EC 2.4.1.17) facilitate excretion of a wide variety of lipophilic drugs, environmental chemicals, and endogenous substrates containing hydroxyl, carboxyl, amino, and sulfur-containing functional groups by catalyzing conjugation of these substrates with glucuronic acid to increase hydrophilicity. UGT enzymes belong to distinct subfamilies of more than 26 genes with 19 well-characterized functional proteins. The functional isoforms belong to the UGT1 and UGT2 superfamilies, which are further divided into three subfamilies based on their sequence similarities into UGT1As (UGT1A1, UGT1A3-UGT1A10), UGT2As (UGT2A1-UGT2A3) and UGT2Bs (UGT2B4, UGT2B7, UGT2B10, UGT2B11, UGT2B15, UGT2B17, and UGT2B28) (Guillemette, 2003; Oda et al., 2015; Yuan et al., 2016).

Although all hepatic UGT isoforms are generally variable, UGT2B17 shows extensively higher interindividual variability in its protein abundance and activity (Fallon et al., 2013; Neumann et al., 2016), and it is expressed in a variety of tissues such as liver, intestine, kidney, testis, uterus, placenta, mammary gland, adrenal gland, skin, and prostate (Beaulieu et al., 1996; Ekstrom et al., 2013). A number of endogenous steroids including testosterone, dihydrotestosterone (DHT), androstane-3- α , 17- β -diol (3- α -diol), androsterone and estradiol and xenobiotics, e.g., 17-dihydroexemestane, vorinostat and lorcaserin have been identified as substrates of UGT2B17 (Beaulieu et al., 1996; Wong et al., 2011; Sadeque et al., 2012; Chen et al., 2016b; Neumann et al., 2016). The expression of UGT2B17 in sex hormone-sensitive organs also indicates its role in sex hormone homeostasis. For example, although controversial, gene deletion in UGT2B17 is associated with higher risk of developing androgen-sensitive prostate diseases (Barbier and Belanger, 2008; Paquet et al., 2012; Kpoghomou et al., 2013; Gauthier-Landry et al., 2015). The *UGT2B17* gene deletion allele has been shown to be associated with a number of other pathophysiological conditions such as obesity (Zhu et al., 2015), chronic lymphocytic leukemia (CLL) (Gruber et al., 2013), and endometrial cancer (Hirata et al., 2010). UGT2B17 also appears to play a critical role in the metabolism of tobacco-specific carcinogens and the risk of lung cancer

DMD # 80952

(Lazarus et al., 2005; Chen et al., 2016a). In addition, high intratumoral UGT2B17 expression levels correlate with better survival outcomes in patients with breast cancer (Hu et al., 2016). Besides its role in disease pathophysiology, the *UGT2B17* gene deletion is associated with false-negative doping test results, which in turn is linked to variable testosterone metabolism (Schulze et al., 2008). An investigational drug developed by Merck, MK-7246, a selective CRTH2 (prostaglandin D2 receptor 2) antagonist, was discontinued from development after unpredicted variability observed in its pharmacokinetics (PK). MK-7246 was later characterized as a selective UGT2B17 substrate (Wang et al., 2012). Similarly, PK and anticancer effectiveness of UGT2B17 substrates, 17-dihydroexemestane and vorinostat, are highly variable (Wong et al., 2011; Chen et al., 2016b). Particularly, the normalized 17-dihydroexemestane and vorinostat levels were 28% and 26% higher, respectively, in subjects carrying UGT2B17 gene deletion as compared to those carrying the reference allele (Wong et al., 2011; Luo et al., 2017). In vitro glucuronidation rate of 17-dihydroexemestane is significantly decreased (14-fold) in human liver microsomes (HLMs) exhibiting the UGT2B17 deletion genotype versus wild-type UGT2B17 HLMs (Sun et al., 2010). Taken together, high variability in UGT2B17 abundance significantly contributes to an unpredictable fate of its substrates that may lead to adverse pathophysiological consequences and drug toxicity or lack of efficacy.

To better understand the underlying causes of UGT2B17 variability, we investigated association of genetic and non-genetic factors with variability in protein abundance and *in vitro* activity of UGT2B17 in human liver microsomes (HLMs). The knowledge of individual contribution of population factors in UGT2B17 variability can be applied to predict metabolism of androgens and other UGT2B17 substrates. Further, because androgens are also metabolized by UGT2B15 (minor pathway), we studied the effect of its genetic polymorphism (UGT2B15*2) on metabolism of testosterone and DHT, particularly in the poor expressers of UGT2B17.

DMD # 80952

MATERIALS AND METHODS

Chemicals and Reagents

Iodoacetamide (IAA), dithiothreitol (DTT), and Pierce trypsin protease (MS grade) were purchased from Thermo Fisher Scientific (Rockford, IL). Ammonium bicarbonate buffer (ABB, 98% purity) was purchased from Acros Organics (Geel, Belgium). Chloroform, MS-grade acetonitrile, methanol and formic acid were purchased from Fischer Scientific (Fair Lawn, NJ). Human serum albumin (HSA) and bovine serum albumin (BSA) were obtained from Calbiochem (Billerica, MA) and Thermo Fisher Scientific (Rockford, IL), respectively. The purified light peptides for UGT2B17 (FSVGYTVEK) and UGT2B15 (SVINDPVYK) were purchased from New England Peptides (Cambridge, MA). Synthetic isotopically pure heavy stable isotope-labeled peptides were produced by Thermo Fisher Scientific (Rockford, IL). UDPGA and $MgCl_2$ were purchased from Sigma-Aldrich (St. Louis, MO). T (1 mg/mL in 100 % acetonitrile) and testosterone-glucuronide were purchased from Cerilliant (Round Rock, TX). Testosterone -glucuronide was dissolved in 100% methanol (1 mg/mL). DHT (1 mg/mL in methanol) and DHT-glucuronide were purchased from Sigma-Aldrich (St. Louis, MO) and Cerilliant (Round Rock, TX), respectively. Testosterone-glucuronide-d3 and DHT-glucuronide were procured from Cerilliant (Round Rock, TX).

Human Liver Tissue and Preparation of Microsomes

Previously prepared HLM samples (Pearce et al., 2016; Shirasaka et al., 2016) were used in this study. The liver tissue samples for HLM preparation were received from three liver tissue banks: i) the University of Washington Human Liver Bank (Seattle, WA) (n=56), ii) Children's Mercy Kansas City (Kansas City, MO) (n=128), and iii) the Liver Bank at the St. Jude Children's Research Hospital (Memphis, TN) (n=271). The Children's Mercy Kansas City samples were originally obtained from the University of Maryland Brain and Tissue Bank for Developmental Disorders and the Liver Tissue Cell Distribution System. Additional details on the selection, procurement and

DMD # 80952

storage information of the livers and investigator blinding for sample analyses have been described previously (Prasad et al., 2014; Shirasaka et al., 2016; Tanner et al., 2017). Age, sex, and ethnicity were known for 96%, 98% and 88% of the liver donors, respectively (Table S1). The age range for donors was from 0-87 years (median age, 24 years). Of the 455 samples analyzed for UGT2B abundance, demographic association analyses were conducted on 423 samples for which *UGT2B17* copy number variation (CNV) information was available. Sex distribution of these 423 samples was- 252 males, 163 females, and 8 were unknown. The ethnicity distribution was 333 Caucasian, 26 African-American, 4 Hispanic, 1 Native American, 1 Asian, 1 Pacific Islander and 56 unknown. Cause of death, medications used, and liver pathology were known for less than 50% of the donors. The collection and use of these tissues for research purposes was approved by the human subjects Institutional Review Boards of the University of Washington (Seattle, WA), the St. Jude Children's Research Hospital (Memphis, TN), and the Pediatric Institutional Review Board of Children's Mercy Kansas City (Kansas City, MO).

UGT2B17 and UGT2B15 Protein Quantification in HLM Samples

Total protein quantification in HLM samples was performed using a BCA assay kit (Pierce™ BCA Protein Assay Kit). HLMs (80 μ L, 2 mg/mL total protein) were digested as described in the Supplementary Materials. The surrogate peptides of UGT2B17 (FSVGYTVEK and SVINDPIYK) and UGT2B15 (SVINDPVYK) were quantified in the digested samples using a validated LC-MS/MS method (Vrana et al., 2017), described in the Supplementary Materials.

***UGT2B17* mRNA Quantification**

A subset of liver tissue samples (n=230) were available for *UGT2B17* mRNA expression analysis (Table S1). Details of the RNA-seq procedures including RNA isolation, TruSeq stranded mRNA preparation, and read processing and analysis pipeline have been described previously (Tanner et al., 2017). mRNA transcript levels are presented in FPKM values.

DMD # 80952

***UGT2B17* Sequencing, Genotyping, Haplotype and Copy Number Variation Analysis**

Because liver samples were obtained from different sources, two approaches, gene sequencing and genotyping, were used for genetic characterization of the liver tissue samples as discussed in the Supplementary Materials. The University of Washington and St. Jude Liver Bank samples were sequenced using the PGRN-Seq platform, a targeted sequencing approach, as described elsewhere (Gordon et al., 2016) whereas the samples provided by Children's Mercy were genotyped on DMET or PharmacoScan arrays (Affymetrix, Santa Clara, CA, USA). Linkage disequilibrium (LD) analysis of *UGT2B17* variants and inferred haplotypes were determined using Haploview 4.2 (Cambridge, MA, USA).

***UGT2B17* and *UGT2B15* Enzyme Activity Assay**

For activity assays, 346 HLM samples (donor age ranges from 0 to 87 years; median age = 18 years, Table S1) were available. Glucuronidation activity was determined by quantifying the rates of testosterone- and DHT-glucuronide formation (pmol/min/mg of microsomal protein), in triplicate. The assay reactions contained 0.1 mg/mL HLM protein, 0.1 M phosphate buffer (pH 7.4), a mix of 1 μ M testosterone and 1 μ M DHT, bovine serum albumin (BSA) (0.01 %) and alamethicin (0.1 mg/mL) (final volume of 95 μ L). Final vehicle (methanol or ethanol) concentration was less than 1 %. Reactions were pre-incubated for 15 minutes on ice. UDPGA (5 μ l; final concentration, 2.5 mM) was added to initiate reactions and mixtures were gently agitated for 30 min at 37°C before being quenched with ice-cold of acetonitrile containing 50 ng/mL progesterone (200 μ l, internal standard) and subjected to centrifugation for 5 min at \sim 1,300 x g. Supernatants were analyzed by an optimized LC-MS/MS method provided in the Supplementary Materials.

Data Analysis

We used a robust strategy to ensure optimum reproducibility of *UGT2B17* and *UGT2B15* protein

DMD # 80952

quantification (Bhatt and Prasad, 2017). For example, ion suppression was addressed by using heavy peptide internal standards. BSA or human serum albumin (HSA) was used as an exogenous protein internal standard, which was added to each sample in a fixed quantity before desalting by methanol-chloroform-water extraction and trypsin digestion. The addition of BSA or HSA addresses the variability introduced during pre-digestion processing such as i) protein loss during methanol-chloroform-water extraction and ii) sample-to-sample trypsin digestion artifacts. To address inter-batch variability, two to three sets of pooled representative HLM samples were processed each day, which served as quality controls across the entire study. In total, a three-step data normalization approach was used; first, average light peak areas for specific peptide daughter fragments were divided by corresponding average heavy peak areas. Next, this ratio was further divided by the BSA or HSA light/heavy area ratio. Finally, for each day, these data were further normalized by mean values of the quality control values run with each individual batch to adjust for any inter-day variability.

Ontogeny was measured by categorical and continuous analyses of age vs. UGT2B17 protein abundance data. For categorical analysis, the samples were grouped based on the following age-categories: neonatal (0 to 27 days), infancy (28 to 364 days), toddler/early childhood (1 year to <6 years), middle childhood (6 to <12 years), adolescence (12 to 18 years) and adulthood (>18 years).

Statistical analyses were performed using GraphPad Prism 5 (La Jolla, CA, USA) and Microsoft Excel (365 ProPlus, Redmond, WA, USA). Non-parametric tests were used to test age-, sex-, and genotype-dependence. To compare two groups (e.g., male vs. female), the Mann-Whitney test was employed. The Kruskal-Wallis test followed by Dunn's multiple comparison tests were used to perform the age-dependent data analyses and determine associations between the genotype and mRNA expression, protein abundance, and the enzyme activity. For correlation analysis, the non-parametric Spearman regression test was used. Additionally, Jonckheere-Terpstra test and multivariate analysis were performed by using RStudio (Version 1.0.136, R version 3.3.2).

DMD # 80952

A nonlinear regression equation (Equation 1) was used to fit the ontogeny data, as described previously (Bhatt et al., 2017), where $Adult_{max}$ is the maximum average relative protein abundance, Age is the age in years of the subject at the time of sample collection, Age_{50} is the age in years at which half-maximum adult protein abundance is obtained, E is protein abundance in adult samples, E_{birth} is protein abundance at birth, and n is the exponential factor.

$$E = \left(\frac{Adult_{max} - E_{birth}}{Age_{50}^n + Age^n} \right) \times Age e^n + E_{birth} \quad (\text{Equation 1})$$

For haplotype analysis, the number of variants was directly counted. Hardy-Weinberg equilibrium (HWE) was determined by comparing the variant frequencies with the expected values using a contingency table chi-square statistic with Yates' correction. The numbers of haplotype, statistics D , D' and LD were estimated by Haploview 4.2 (Cambridge, MA, USA) software. Relationships were considered significant at $P < 0.05$.

RESULTS

Hierarchical Clustering of Major Hepatic Drug Metabolizing Enzymes and Correlation Analysis of UGT2B17 Protein, mRNA and Activity

Hierarchical clustering analysis of quantitative proteomics results of major hepatic drug metabolizing enzymes from a preliminary study conducted in first 165 samples (out of $n=455$) suggested unique and highly variable protein abundance for UGT2B17 (Figure S1). The correlation between *UGT2B17* mRNA and protein abundance or activity was moderate ($r^2 = 0.17$ and 0.19 , respectively) but statistically significant ($P < 0.0001$) (Figure S2-A). Consistent with the literature (Ohtsuki et al., 2012), the correlation between mRNA and protein expression in tissues was weak; in contrast, a strong correlation between protein abundance and activity was observed (Figure S2-B and C). Although our quantitative proteomics method was very sensitive for UGT2B17 detection (lower limit of quantification, LLOQ=0.17 pmol/mg of microsomal protein), the

DMD # 80952

protein was not detected in 48% of the samples indicating a rather large range of variability in protein expression. For statistical analysis, samples below the limit of detection (<LOD) were assigned a value of 0.06 pmol/mg of microsomal protein (i.e., 1/3rd of LLOQ) instead of zero. In low UGT2B17 expressers, rates of testosterone- and DHT-glucuronide formation were also consistently low. The average human liver UGT2B17 protein abundance in liver microsomal samples (n=370, excluding zero copy number samples) was 0.92 ± 1.6 pmol/mg microsomal protein with 162-fold interindividual variability (0.06-9.7 pmol/mg microsomal protein). Mean rates of testosterone and DHT glucuronide formation (range; fold difference) were 15.4 (0.3-184; 558) and 41.8 (1.0-233; 233) pmol/min/mg microsomal protein, respectively (Table 1).

Association of UGT2B17 Protein Abundance, Testosterone/DHT Glucuronide Formation and Genetic Variation

UGT2B17 protein was detected in 38% and 52% of the samples carrying one and two gene copies, respectively. UGT2B17 protein was undetectable in samples homozygous for the *UGT2B17* gene deletion (CNV=0). This variability was also reflected by mRNA data demonstrating that fragments per kilobase per million reads (FPKM) values were significantly higher ($P < 0.0001$) in samples with one or two *UGT2B17* gene copies; however, FPKM values were not different among samples with one and two gene copies due to the high variability (>205-fold) within each group. Samples carrying two *UGT2B17* gene copies showed a 1.7-fold higher mean UGT2B17 protein abundance compared to single gene copy samples (Figure 1 and Table 1). The *UGT2B17* gene-dose dependent effects on rates of testosterone- and DHT-glucuronide formation rates were consistent with the abundance data (Figure 1 and Table 1). Of the 11 *UGT2B17* SNPs detected in our samples, only four variants (rs7436962, rs9996186, rs28374627, and rs7668258) were associated with mRNA expression, protein abundance, and activity (Table S2). Haplotype analyses suggested significant linkage disequilibrium (LD) between three intronic SNPs

DMD # 80952

(rs7436962, rs9996186, and rs4860305) and a missense SNP rs28374627 (Figure 1). We identified four haplotypes (H1-H4; Figure 1) representing combinations of these SNPs with frequency >10% in our sample set. When the diplotypes were compared, samples harboring H3/H4 and H2/H2 haplotypes showed higher UGT2B17 mRNA expression, protein abundance, and activity compared to the reference H1/H1 diplotype (Figure 1 and Table 1). The gene-dose effect was also verified by using multivariate analysis (Table 2) and Jonckheere-Terpstra test (Table S5)

Association of UGT2B17 Abundance and Testosterone/DHT Glucuronide Formation with Age and Sex

UGT2B17 protein abundance was significantly higher in adults compared to infancy and early/middle childhood (Figure 2A and Figure S3, P values are marked in the figures). Noticeably, the age at which protein expression reaches 50% of that observed in adults (Age_{50}) was >10 years in both male and female samples (Figure S3). Trend analysis (Jonckheere-Terpstra test) showed that there was a statistically significant higher median UGT2B17 abundance with increasing age category (neonatal - infant - early childhood- middle childhood- adolescence- adulthood; Table S5). Male liver donors have a 2.8-fold higher mean UGT2B17 protein level ($P<0.0001$) compared to females in samples from donors ≥ 12 years of age (Figure 2D) (Table 1). Consistent with the protein abundance data, rates of testosterone- and DHT-glucuronide formation were significantly higher in adulthood compared to neonatal, infancy, early childhood and middle childhood age groups (Figure 2; E and I). Age-dependent increase in UGT2B17 abundance and activity was greater in male vs. female livers (Figure 2; B, F, J vs. C, G, K). Overall, male liver showed 2- and 1.4-fold higher rates of testosterone- and DHT-glucuronide formation compared to females in samples ≥ 12 years of age (Figure 2; H and L).

Association of UGT2B15 Abundance and Testosterone/DHT-Glucuronide Formation with

DMD # 80952

UGT2B15*2 in Poor UGT2B17 Expressers

An association of rates of testosterone- or DHT-glucuronide formation with UGT2B15 protein abundance was only observed in samples with low UGT2B17 levels (i.e., <LOD) (Figure S4). Importantly, in these low UGT2B17 expressers, rs1902023, a non-synonymous *UGT2B15* SNP (Court et al., 2004), was not associated with UGT2B15 protein abundance, however there was a significant gene-dose dependent association with this SNP and rates of testosterone- and DHT-glucuronide formation (Figure 3), respectively. These data suggest that this non-synonymous UGT2B15 genotype likely affects substrate affinity (km) or catalytic activity (kcat) due to change in amino acid residue in the active or cofactor binding site. Furthermore, the correlation between protein level and activity improved in samples that did not carry heterozygous or homozygous variants of rs1902023 (Figure S4).

Multivariate Linear Regression Analyses

The results were further verified using multivariate linear regression analyses to evaluate associations with UGT2B17 protein abundance, and rates of testosterone-, or DHT-glucuronide formation as the outcomes, whereas CNV, diplotype, age category, and sex as the predictors. Base-line parameters were set as females, children, reference diplotype (H1/H1), and a copy number of 1 (0 copy number samples were excluded from the analyses). The results of the multivariate linear regression analyses, with missing values imputed are presented in Table 2. For UGT protein abundance, the coefficient of determination (R^2 value) for the multivariate linear regression was 0.26, indicating that 26% of the variability in UGT2B17 abundance is explained by the predictors in the model. Sex, age categories and diplotypes were significant predictors of UGT2B17 protein abundance. For testosterone- and DHT-glucuronide formation, the R^2 value for the multivariate linear regression were 0.21 and 0.24, indicating that 21% and 24% of the variability in testosterone- and DHT-glucuronide formation is explained by the predictors in the model, respectively. All other multivariate analysis output parameters are presented in Table 2.

DMD # 80952

DISCUSSION

Distinct from other major drug metabolizing enzymes, unusually high interindividual variability was observed for UGT2B17 in our large cohort of human liver tissue samples. The major factors impacting the observed variability include CNVs, SNPs, age and sex. Developmental *UGT2B17* gene expression, and association of SNPs located in the *UGT2B17* gene with its mRNA expression have been previously reported (Burgess et al., 2015; Neumann et al., 2016). Similarly, highly variable protein abundance of UGTs in adult liver is known (Fallon et al., 2013). However, our data are novel in respect to measuring protein abundance by selective LC-MS/MS proteomics, enzyme activity using two probe substrates (testosterone and DHT), and comprehensive CNV and diplotype analyses in the same set of samples. The large cohort of samples allowed comprehensive multivariate analyses, which revealed the individual contributions of many factors impacting UGT2B17 protein abundance and androgen glucuronidase activities. These protein abundance and activity data are important to predict variability in the metabolism of UGT2B17 xenobiotic substrates and sex steroids.

With respect to drug metabolism, UGT2B17 is a less studied enzyme and no regulatory guidance for industry currently exists for this enzyme. The US Food and Drug Administration (USFDA) and European Medicines Agency (EMA) recommend *in vitro* testing for the likelihood of a new chemical entity to be a substrate or inhibitor of other UGT isoforms such as UGT1A1, UGT1A3, UGT1A4, UGT1A6, UGT1A9, UGT2B7 and UGT2B15. Our data predicts that overlooking UGT2B17 could lead to clinical failure of a UGT2B17 substrate drug due to high PK variability. Indeed, the UGT2B17 substrate MK-7246 was discontinued from clinical trials due to high PK variability (Wang et al., 2012). Consistent with the literature (Gallagher et al., 2010), females have lower UGT2B17 expression levels compared to males. Likewise, distinct ontogeny of UGT2B17 as compared to the common drug metabolizing enzymes, cytochrome P450s and other UGTs, is

DMD # 80952

an important finding of this study. These data predict that the use of UGT2B17 substrate drugs, e.g., vorinostat and lorcaserin, in women and children below 12 years could lead to supratherapeutic drug levels. We therefore recommend that UGT2B17 should be included in the in vitro UGT screening panel during early drug discovery and caution should be taken when designing clinical studies of UGT2B17 substrate drugs in females and children along with consideration of genetic polymorphisms. UGT2B17 is expressed in other organs such as intestine, appendix, bone-marrow and prostate; however, liver is considered as an effective elimination organ for UGT2B17 substrates because of its larger size (resulting high total abundance) and high blood flow. The liver microsomes used in this study were isolated in two different laboratories, and other factors such as medication use and storage condition could affect protein abundance in the microsomes. However, the lack of correlation between UGT2B17 and other proteins (e.g., UGT2B15, Figure S1) in the same samples indicates that the observed UGT2B17 data primarily reflect biological or inter-individual variability.

By regulating testosterone metabolism, UGT2B17 is linked to multiple pathophysiological conditions such as obesity (Zhu et al., 2015) and prostate cancer (Barbier and Belanger, 2008; Paquet et al., 2012; Kpoghomou et al., 2013; Gauthier-Landry et al., 2015). For example, the *UGT2B17* gene deletion (homozygous) is shown to be associated with decrease in fat mass ($P < 0.01$) and insulin sensitivity ($P < 0.05$) (Swanson et al., 2007), and the males with lower testosterone levels are 2.4 times more likely to be obese than males with higher testosterone levels (Mulligan et al., 2006). On the other hand, high UGT2B17 protein levels have been identified as the strongest independent molecular prognostic marker of overall survival in mutated chronic lymphocytic leukemia patients (Bhoi et al., 2016). The *UGT2B17* deletion (homozygous) genotype is associated with a decreased risk of colorectal cancer in men, but was non-predictive in women (Angstadt et al., 2013). It is also noteworthy that association studies on UGT2B17 gene deletion and disease risks, e.g., prostate cancer, are controversial. These contradictions in

DMD # 80952

literature could be explained as the published association studies do not acknowledge the effect of confounding factors other than gene deletion (e.g., SNPs and non-genetic factors) on UGT2B17 variability. Further, UGT2B15 which affects testosterone glucuronidation in UGT2B17 poor expressers has not been considered in these association studies. Therefore, the data presented here will help designing better clinical studies to investigate association of individual factors affecting UGT2B17 with disease risk.

Although testosterone and its glucuronides are believed to be transported by organic anion polypeptide transporters (OATPs) and multidrug resistance-associated protein (MRPs) (Hamada et al., 2008), the UGT2B17 interindividual variability is significantly higher as compared to the variability in transporter abundance previously reported by us in a subset of these samples (Prasad et al., 2014; Prasad et al., 2016). Nevertheless, genetic polymorphism in transporters should also be considered when designing clinical studies to investigate association of UGT2B17 variability with testosterone related clinical outcomes.

UGT2B17 interindividual variability data could also be used to develop a better doping test approach to avoid false negative or positive test results. The urinary testosterone (T) to epitestosterone (E) ratio (T/E) has a cut-off limit of 4 and is used to detect testosterone-doping in all cases. Testosterone is metabolized by UGT2B17 while epitestosterone is metabolized by UGT2B7. Individuals homozygous for the *UGT2B17* deletion allele excrete negligible amounts of testosterone in urine compared to subjects with one or two gene copies (Bao et al., 2008) and rarely reach the T/E cut-off value of 4 after testosterone-doping, indicating that genetic testing for the *UGT2B17* deletion allele may increase the chances of identifying atypical findings, especially in dissecting false-negative test results. Moreover, based on our study, there should be different cut-off values for males vs. females, adolescents vs. adults, and for different haplotypes.

DMD # 80952

The sex- or age-dependent expression of UGT2B17 may be explained by its regulation by androgens and estradiol. For example, Bao et al. demonstrated that *UGT2B15* and *2B17* are androgen-regulated genes and that androgen receptor (AR) is required for both their basal and androgen-regulated expression (Bao et al., 2008). Similarly, UGT2B17 is 5-fold more abundant in metastatic versus benign prostate cancer samples (Paquet et al., 2012). *UGT2B17* and myeloid cell leukemia-1 (Mcl-1) expression is upregulated in endometrial cancer (EC) tissues and UGT2B17 depletion induces inhibition of cell growth and apoptosis in EC cells through Mcl-1 downregulation (Hirata et al., 2010). However, UGT2B17 variability within a single group (e.g., adult males) indicates the involvement of multiple other epigenetic and transcriptional mechanisms. For example, UGT2B15 and UGT2B17 are both negatively regulated by the miR-376c microRNA that binds to the 3'-UTRs of UGT2B15 and UGT2B17 mRNA in prostate cancer cells (Wijayakumara et al., 2015). It has also been shown that Forkhead Box Protein A1 (FOXA1) regulates UGT2B17 gene transcription in LNCaP prostate cancer cells (Hu et al., 2010). Similarly, polymorphic PXR and CAR are associated with altered expression of UGT2Bs, respectively (Verreault et al., 2010). Therefore, SNPs in these polymorphic transcriptional factors, e.g., -298G/G and 11193C/C in PXR (Du et al., 2013) and IVS2-99C>T in CAR (Urano et al., 2009), can also indirectly influence UGT2B17 expression. Clearly, further research characterizing the underlying mechanisms contributing to interindividual variability in UGT2B17 abundance and activity is warranted.

Despite the significant physiological role of testosterone, testosterone replacement therapy (TRT) is controversial. For example, a meta-analysis suggested an association of TRT with prostate cancer; however, many independent studies failed to reproduce this finding (Barbier and Belanger, 2008; Paquet et al., 2012; Kpoghomou et al., 2013; Gauthier-Landry et al., 2015). The FDA recently re-issued a black-box warning on TRT because of its association with cardiovascular side effects. Our data suggest that a UGT2B17 and UGT2B15-based precision medicine strategy

DMD # 80952

for TRT, may ensure safe and effective testosterone use. A similar strategy may also be considered to improve high-testosterone therapy (also referred as bipolar androgen therapy, BAT) in prostate cancer patients (Schweizer et al., 2015).

Taken together, the findings of this study are of clinical importance and can be directly translated to individualize drug therapy, by stratifying patients based on *UGT2B17* and *UGT2B15* genotype and predicted phenotype. Moreover, physiologically-based PK (PBPK) models can be developed based on these data to accurately predict *UGT2B17* mediated glucuronidation of endobiotics and xenobiotics and translate such data to predict *in vivo* disposition of these substrates.

DMD # 80952

ACKNOWLEDGMENTS

We would like to thank Prachi Choudhari and Mikael Boberg for assistance with LC-MS/MS sample analysis. We sincerely thank Dr. Kenneth Thummel and Dr. Allan Rettie, UW School of Pharmacy for their valuable suggestions throughout this project. We would also like to thank Amy Turner, Praful Aggarwal and Rachel Lorier from Dr. Ulrich Broeckel's lab to generate a part of the genotyping data.

DMD # 80952

AUTHORSHIP CONTRIBUTIONS

Participated in research design: D.K.B., A.G., J.S.L., B.P.

Conducted experiments: D.K.B., A.B., H.Z., K.G.C., S.L., A.M., B.P., A.G., R.E.P., R.G., U.B.

Contributed new reagents or analytic tools: A.S.C., E.G.S., J.S.L., B.P.

Performed data analysis: D.K.B., A.B., H.Z., K.G.C., S.L., A.M., A.G., U.B., T.A.T., B.P., D.A.N.

Wrote or contributed to the writing of the manuscript: D.K.B., A.B., H.Z., K.G.C., A.M., A.G., S.L.,
R.E.P., R.G., U.B., D.A.N., T.A.T., J.K.A., J.S.L., B.P.

DMD # 80952

REFERENCES

- Angstadt AY, Berg A, Zhu J, Miller P, Hartman TJ, Lesko SM, Muscat JE, Lazarus P, and Gallagher CJ (2013) The effect of copy number variation in the phase II detoxification genes UGT2B17 and UGT2B28 on colorectal cancer risk. *Cancer* **119**:2477-2485.
- Bao BY, Chuang BF, Wang Q, Sartor O, Balk SP, Brown M, Kantoff PW, and Lee GSM (2008) Androgen Receptor Mediates the Expression of UDP-Glucuronosyltransferase 2 B15 and B17 Genes. *Prostate* **68**:839-848.
- Barbier O and Belanger A (2008) Inactivation of androgens by UDP-glucuronosyltransferases in the human prostate. *Best Pract Res Clin Endocrinol Metab* **22**:259-270.
- Beaulieu M, Levesque E, Hum DW, and Belanger A (1996) Isolation and characterization of a novel cDNA encoding a human UDP-glucuronosyltransferase active on C19 steroids. *J Biol Chem* **271**:22855-22862.
- Bhatt DK, Gaedigk A, Pearce RE, Leeder JS, and Prasad B (2017) Age-dependent Protein Abundance of Cytosolic Alcohol and Aldehyde Dehydrogenases in Human Liver. *Drug Metab Dispos* **45**:1044-1048.
- Bhatt DK and Prasad B (2017) Critical issues and optimized practices in quantification of protein abundance level to determine inter-individual variability in DMET proteins by LC-MS/MS proteomics. *Clin Pharmacol Ther* **10.1002/cpt.819**.
- Bhoi S, Baliakas P, Cortese D, Mattsson M, Engvall M, Smedby KE, Juliusson G, Sutton LA, and Mansouri L (2016) UGT2B17 expression: a novel prognostic marker within IGHV-mutated chronic lymphocytic leukemia?, in: *Haematologica*, pp e63-65.
- Burgess KS, Philips S, Benson EA, Desta Z, Gaedigk A, Gaedigk R, Segar MW, Liu Y, and Skaar TC (2015) Age-Related Changes in MicroRNA Expression and Pharmacogenes in Human Liver. *Clin Pharmacol Ther* **98**:205-215.
- Chen G, Luo S, Kozlovich S, and Lazarus P (2016a) Association between Glucuronidation Genotypes and Urinary NNAL Metabolic Phenotypes in Smokers. *Cancer Epidemiol Biomarkers Prev* **25**:1175-1184.
- Chen SM, Atchley DH, Murphy MA, Gurley BJ, and Kamdem LK (2016b) Impact of UGT2B17 Gene Deletion on the Pharmacokinetics of 17-Hydroxemestane in Healthy Volunteers. *J Clin Pharmacol* **56**:875-884.
- Court MH, Hao Q, Krishnaswamy S, Bekaii-Saab T, Al-Rohaimi A, von Moltke LL, and Greenblatt DJ (2004) UDP-glucuronosyltransferase (UGT) 2B15 pharmacogenetics: UGT2B15 D85Y genotype and gender are major determinants of oxazepam glucuronidation by human liver. *J Pharmacol Exp Ther* **310**:656-665.
- Du QQ, Wang ZJ, He L, Jiang XH, and Wang L (2013) PXR polymorphisms and their impact on pharmacokinetics/pharmacodynamics of repaglinide in healthy Chinese volunteers. *Eur J Clin Pharmacol* **69**:1917-1925.
- Ekstrom L, Johansson M, and Rane A (2013) Tissue distribution and relative gene expression of UDP-glucuronosyltransferases (2B7, 2B15, 2B17) in the human fetus. *Drug Metab Dispos* **41**:291-295.
- Fallon JK, Neubert H, Hyland R, Goosen TC, and Smith PC (2013) Targeted quantitative proteomics for the analysis of 14 UGT1As and -2Bs in human liver using NanoUPLC-MS/MS with selected reaction monitoring. *J Proteome Res* **12**:4402-4413.
- Gallagher CJ, Balliet RM, Sun D, Chen G, and Lazarus P (2010) Sex Differences in UDP-Glucuronosyltransferase 2B17 Expression and Activity, in: *Drug Metab Dispos*, pp 2204-2209.
- Gauthier-Landry L, Belanger A, and Barbier O (2015) Multiple roles for UDP-glucuronosyltransferase (UGT)2B15 and UGT2B17 enzymes in androgen metabolism and prostate cancer evolution. *J Steroid Biochem Mol Biol* **145**:187-192.
- Gordon AS, Fulton RS, Qin X, Mardis ER, Nickerson DA, and Scherer S (2016) PGRNseq: a targeted capture sequencing panel for pharmacogenetic research and implementation. *Pharmacogenetics and genomics* **10.1097/FPC.000000000000202**.
- Gruber M, Bellemare J, Hoermann G, Gleiss A, Porpacz E, Bilban M, Le T, Zehetmayer S, Mannhalter C, Gaiger A, Shehata M, Fleiss K, Skrabs C, Levesque E, Vanura K, Guillemette C, and Jaeger U (2013) Overexpression of uridine diphospho glucuronosyltransferase 2B17 in high-risk chronic lymphocytic leukemia. *Blood* **121**:1175-1183.
- Guillemette C (2003) Pharmacogenomics of human UDP-glucuronosyltransferase enzymes. *Pharmacogenomics J* **3**:136-158.
- Hamada A, Sissung T, Price DK, Danesi R, Chau CH, Sharifi N, Venzon D, Maeda K, Nagao K, Sparreboom A, Mitsuya H, Dahut WL, and Figg WD (2008) Effect of SLCO1B3 haplotype on testosterone transport and clinical outcome in Caucasian patients with androgen-independent prostatic cancer. *Clin Cancer Res* **14**:3312-3318.

DMD # 80952

- Hirata H, Hinoda Y, Zaman MS, Chen Y, Ueno K, Majid S, Tripsas C, Rubin M, Chen LM, and Dahiya R (2010) Function of UDP-glucuronosyltransferase 2B17 (UGT2B17) is involved in endometrial cancer. *Carcinogenesis* **31**:1620-1626.
- Hu DG, Gardner-Stephen D, Severi G, Gregory PA, Treloar J, Giles GG, English DR, Hopper JL, Tilley WD, and Mackenzie PI (2010) A novel polymorphism in a forkhead box A1 (FOXA1) binding site of the human UDP glucuronosyltransferase 2B17 gene modulates promoter activity and is associated with altered levels of circulating androstane-3 α ,17 β -diol glucuronide. *Mol Pharmacol* **78**:714-722.
- Hu DG, Selth LA, Tarulli GA, Meech R, Wijayakumara D, Chanawong A, Russell R, Caldas C, Robinson JL, Carroll JS, Tilley WD, Mackenzie PI, and Hickey TE (2016) Androgen and Estrogen Receptors in Breast Cancer Coregulate Human UDP-Glucuronosyltransferases 2B15 and 2B17. *Cancer Res* **76**:5881-5893.
- Kpoghomou MA, Soatiana JE, Kalembo FW, Bishwajit G, and Sheng W (2013) UGT2B17 Polymorphism and Risk of Prostate Cancer: A Meta-Analysis. *ISRN Oncol* **2013**:465916.
- Lazarus P, Zheng Y, Runkle EA, Muscat JE, and Wiener D (2005) Genotype-phenotype correlation between the polymorphic UGT2B17 gene deletion and NNAL glucuronidation activities in human liver microsomes. *Pharmacogenet Genomics* **15**:769-778.
- Luo S, Chen G, Truica C, Baird CC, Leitzel K, and Lazarus P (2017) Role of the UGT2B17 deletion in exemestane pharmacogenetics. *Pharmacogenomics J*.
- Mulligan T, Frick M, Zuraw Q, Stemhagen A, and McWhirter C (2006) Prevalence of hypogonadism in males aged at least 45 years: the HIM study. *Int J Clin Pract* **60**:762-769.
- Neumann E, Mehboob H, Ramirez J, Mirkov S, Zhang M, and Liu W (2016) Age-Dependent Hepatic UDP-Glucuronosyltransferase Gene Expression and Activity in Children. *Front Pharmacol* **7**:437.
- Oda S, Fukami T, Yokoi T, and Nakajima M (2015) A comprehensive review of UDP-glucuronosyltransferase and esterases for drug development. *Drug Metab Pharmacokinet* **30**:30-51.
- Ohtsuki S, Schaefer O, Kawakami H, Inoue T, Liehner S, Saito A, Ishiguro N, Kishimoto W, Ludwig-Schwellinger E, Ebner T, and Terasaki T (2012) Simultaneous absolute protein quantification of transporters, cytochromes P450, and UDP-glucuronosyltransferases as a novel approach for the characterization of individual human liver: comparison with mRNA levels and activities. *Drug Metab Dispos* **40**:83-92.
- Paquet S, Fazli L, Grosse L, Verreault M, Tetu B, Rennie PS, Belanger A, and Barbier O (2012) Differential expression of the androgen-conjugating UGT2B15 and UGT2B17 enzymes in prostate tumor cells during cancer progression. *The Journal of clinical endocrinology and metabolism* **97**:E428-432.
- Pearce RE, Gaedigk R, Twist GP, Dai H, Riffel AK, Leeder JS, and Gaedigk A (2016) Developmental Expression of CYP2B6: A Comprehensive Analysis of mRNA Expression, Protein Content and Bupropion Hydroxylase Activity and the Impact of Genetic Variation. *Drug Metab Dispos* **44**:948-958.
- Prasad B, Evers R, Gupta A, Hop CE, Salphati L, Shukla S, Ambudkar SV, and Unadkat JD (2014) Interindividual variability in hepatic organic anion-transporting polypeptides and P-glycoprotein (ABCB1) protein expression: quantification by liquid chromatography tandem mass spectroscopy and influence of genotype, age, and sex. *Drug Metab Dispos* **42**:78-88.
- Prasad B, Gaedigk A, Vrana M, Gaedigk R, Leeder JS, Salphati L, Chu X, Xiao G, Hop C, Evers R, Gan L, and Unadkat JD (2016) Ontogeny of Hepatic Drug Transporters as Quantified by LC-MS/MS Proteomics. *Clin Pharmacol Ther* **100**:362-370.
- Sadeque AJ, Usmani KA, Palamar S, Cerny MA, and Chen WG (2012) Identification of human UDP-glucuronosyltransferases involved in N-carbamoyl glucuronidation of lorcaserin. *Drug Metab Dispos* **40**:772-778.
- Schulze JJ, Lundmark J, Garle M, Skilving I, Ekstrom L, and Rane A (2008) Doping test results dependent on genotype of uridine diphospho-glucuronosyl transferase 2B17, the major enzyme for testosterone glucuronidation. *J Clin Endocrinol Metab* **93**:2500-2506.
- Schweizer MT, Antonarakis ES, Wang H, Ajiboye AS, Spitz A, Cao H, Luo J, Haffner MC, Yegnasubramanian S, Carducci MA, Eisenberger MA, Isaacs JT, and Denmeade SR (2015) Effect of bipolar androgen therapy for asymptomatic men with castration-resistant prostate cancer: results from a pilot clinical study. *Science translational medicine* **7**:269ra262.
- Shirasaka Y, Chaudhry AS, McDonald M, Prasad B, Wong T, Calamia JC, Fohner A, Thornton TA, Isoherranen N, Unadkat JD, Rettie AE, Schuetz EG, and Thummel KE (2016) Interindividual variability of CYP2C19-catalyzed drug metabolism due to differences in gene diplotypes and cytochrome P450 oxidoreductase content.

DMD # 80952

- Pharmacogenomics J* **16**:375-387.
- Sun D, Chen G, Dellinger RW, Sharma AK, and Lazarus P (2010) Characterization of 17-dihydroexemestane glucuronidation: potential role of the UGT2B17 deletion in exemestane pharmacogenetics. *Pharmacogenet Genomics* **20**:575-585.
- Swanson C, Mellstrom D, Lorentzon M, Vandenput L, Jakobsson J, Rane A, Karlsson M, Ljunggren O, Smith U, Eriksson AL, Belanger A, Labrie F, and Ohlsson C (2007) The uridine diphosphate glucuronosyltransferase 2B15 D85Y and 2B17 deletion polymorphisms predict the glucuronidation pattern of androgens and fat mass in men. *J Clin Endocrinol Metab* **92**:4878-4882.
- Tanner JA, Prasad B, Claw KG, Stapleton P, Chaudhry A, Schuetz EG, Thummel KE, and Tyndale RF (2017) Predictors of Variation in CYP2A6 mRNA, Protein, and Enzyme Activity in a Human Liver Bank: Influence of Genetic and Nongenetic Factors. *J Pharmacol Exp Ther* **360**:129-139.
- Urano T, Usui T, Shiraki M, Ouchi Y, and Inoue S (2009) Association of a single nucleotide polymorphism in the constitutive androstane receptor gene with bone mineral density. *Geriatrics & gerontology international* **9**:235-241.
- Verreault M, Kaeding J, Caron P, Trottier J, Grosse L, Houssin E, Paquet S, Perreault M, and Barbier O (2010) Regulation of endobiotics glucuronidation by ligand-activated transcription factors: physiological function and therapeutic potential. *Drug Metab Rev* **42**:110-122.
- Vrana M, USA UoWDoPSW, Whittington D, Medicinal Chemistry UoWSWU, Nautiyal V, India IloTKKWB, Prasad B, and USA UoWDoPSW (2017) Database of Optimized Proteomic Quantitative Methods for Human Drug Disposition - Related Proteins for Applications in Physiologically Based Pharmacokinetic Modeling. *CPT: Pharmacometrics & Systems Pharmacology* **6**:267-276.
- Wang YH, Trucksis M, McElwee JJ, Wong PH, Maciolek C, Thompson CD, Prueksaritanont T, Garrett GC, Declercq R, Vets E, Willson KJ, Smith RC, Klappenbach JA, Opitck GJ, Tsou JA, Gibson C, Laethem T, Panorchan P, Iwamoto M, Shaw PM, Wagner JA, and Harrelson JC (2012) UGT2B17 genetic polymorphisms dramatically affect the pharmacokinetics of MK-7246 in healthy subjects in a first-in-human study. *Clin Pharmacol Ther* **92**:96-102.
- Wijayakumara DD, Hu DG, Meech R, McKinnon RA, and Mackenzie PI (2015) Regulation of Human UGT2B15 and UGT2B17 by miR-376c in Prostate Cancer Cell Lines. *J Pharmacol Exp Ther* **354**:417-425.
- Wong NS, Seah E, Wang LZ, Yeo WL, Yap HL, Chuah B, Lim YW, Ang PC, Tai BC, Lim R, Goh BC, and Lee SC (2011) Impact of UDP-gluconosyltransferase 2B17 genotype on vorinostat metabolism and clinical outcomes in Asian women with breast cancer. *Pharmacogenetics and genomics* **21**:760-768.
- Yuan LM, Gao ZZ, Sun HY, Qian SN, Xiao YS, Sun LL, and Zeng S (2016) Inter-isoform Hetero-dimerization of Human UDP-Glucuronosyltransferases (UGTs) 1A1, 1A9, and 2B7 and Impacts on Glucuronidation Activity. *Sci Rep* **6**:34450.
- Zhu AZ, Cox LS, Ahluwalia JS, Renner CC, Hatsukami DK, Benowitz NL, and Tyndale RF (2015) Genetic and phenotypic variation in UGT2B17, a testosterone-metabolizing enzyme, is associated with BMI in males. *Pharmacogenet Genomics* **25**:263-269.

DMD # 80952

FOOTNOTES

A majority of this work was supported by the National Institute of Child Health and Human Development (NICHD), National Institutes of Health (NIH) grant [R01 HD081299]. The data collection was also supported by NIH grants, [U01 GM092676] and [P01 GM116691]. The NICHD Brain and Tissue Bank for Developmental Disorders at the University of Maryland is funded by the NIH contract, [N01-HD-9-0011/HHSN275200900011C] and the Liver Tissue Cell Distribution System is funded by NIH contract number, [N01-DK-7-0004/ HHSN267200700004C].

DMD # 80952

LEGENDS FOR FIGURES

Figure 1. The *UGT2B17* gene deletion is associated with its protein abundance (A), rates of testosterone- and DHT-glucuronide formation (B and C, respectively). *UGT2B17* diplotypes (haplotype pairs on homologous chromosomes) are associated with *UGT2B17* mRNA expression (D), protein abundance (E), testosterone-glucuronide formation (F) and DHT-glucuronide formation (G). Confounding factor, age below 12 years samples were excluded from the sub-analysis. *, ** and *** represent p values <0.05, <0.01 and 0.0001, respectively. Sample number in each group is shown in parentheses in the x-axis.

Figure 2. Categorical age-dependent *UGT2B17* protein abundance (A-D), testosterone (T)-glucuronide formation (E-H) and DHT-glucuronide formation (I-L) data in all (A, E and I), male (B, F and J) and female (C, G and K) livers. X-axis labels identifying data categories in the bottom panel (I-L) are also applicable to the corresponding top two panels (A-H). Number of samples is presented either as main label in x-axis (A-H) or x-axis parenthesis (I-L). Donors with zero *UGT2B17* gene copy were excluded from this analysis. 205 out of 375 of these samples (male plus female) were below LOD of *UGT2B17* protein measurement. For statistical analysis, samples <LOD (excluding zero copy number) were assigned a value of 0.06 pmol/mg of microsomal protein, which was 1/3rd of LLOQ (0.17 pmol/mg of microsomal protein). *UGT2B17* was sparsely (12 out of 92 samples) detected in children below the age of 9 years. Association of age with *UGT2B17* abundance or testosterone and DHT-glucuronide formation was more prominent in male vs. female. Mean *UGT2B17* protein abundance and testosterone and DHT-glucuronide formation in these samples was 2.8-, 1.9- and 1.4-fold higher in male vs. female donors with age ≥12 years, respectively (D). *, ** and *** represent p values <0.05, <0.01 and <0.0001, respectively.

DMD # 80952

Figure 3: The *UGT2B15* SNP, rs1902023, is not associated with *UGT2B15* protein abundance in low *UGT2B17* expressers (A), but is significantly associated with increased rates of testosterone-glucuronide (B) and DHT-glucuronide (C) formation (pmol/mg/min microsomal protein). Only samples with *UGT2B17* abundance <LOD (0.06 pmol/mg of microsomal protein) samples were included in this analysis.

Table 1: Effect of genetic variations, age and gender on UGT2B17 protein abundance and activity. 32 samples (out of 450) were excluded from these analyses because copy number (CN) variation data were not available on these samples.

	Protein abundance (pmol/mg of microsomal protein)						Glucuronide formation activity (pmol/min/mg microsomal protein)									
	n	Mean	Median	Range (min- max, difference)	fold BLOQ	Number of samples	n	Mean	Median	Range (min- max, difference)	fold	n	Mean	Median	Range (min- max, difference)	fold
• All samples*	370	0.92	0.06	0.06-9.7, 162		202	325	15.4	6.4	0.3-184, 558		325	47.8	26.2	1.0-233, 233	
• CN=0	53	0	0	-		-	46	5.8	3.7	0.6-20.5, 34		46	27.8	21.2	2.6-90.2, 35	
• CN=1	172	0.67	0.06	0.06-6.5, 108		107	133	13.8	6.2	0.3-73.6, 223		133	40.8	27.4	1.0-165, 165	
• CN=2	198	1.14	0.19	0.06-9.7, 162		95	146	19.9	7.8	0.5-184, 368		146	47.7	29.5	1.5-233, 155	
Diplotypes																
• Reference CAGA/CAGA	131	0.57	0.06	0.06-7.9, 132		94	91	13.8	5.7	0.5-123, 246		91	38.7	25.3	2.1-181.4, 86.4	
• Heterozygous TCAA/CAGG	62	1.59	1.04	0.06-6.02, 100		15	35	27.5	21.4	0.5-98.2, 196		35	63.4	58.8	1.5-160, 107	
• Homozygous TCAG/TCAG	58	1.63	1.03	0.06-9.7, 162		11	31	30.3	23.2	1.9-184, 96.8		31	66.9	57	8.3-233, 28	
Age Categories																
• Neonatal	3	0.06	0.06	0.06-0.06		3	3	1.1	0.94	0.4-2.0, 5		3	4.0	3.8	1.6-9.0, 5.6	
• Infancy	23	0.11	0.06	0.06-0.86, 14.3		20	19	3.3	1.3	0.4-11.5, 29		19	14.1	6.2	1.0-55.1, 55.1	
• Early childhood	38	0.13	0.06	0.06-1.19, 21		34	29	6.6	3.7	0.3-33.1, 110		29	24.8	17.0	1.0-101, 101	
• Middle childhood	44	0.15	0.06	0.06-1.49, 24.8		34	35	6.0	3.7	0.7-38.9, 56		35	25.0	16.9	2.6-116, 44.6	
• Adolescence	61	0.81	0.06	0.06-6.9, 115		33	54	18.8	7.9	0.5-131.3, 263		54	48.5	32.1	2.1-194, 92	
• Adulthood	185	1.33	0.46	0.06-9.7, 162		76	127	23.2	12.6	0.8-185.8, 232		127	56.3	47.4	3.0-235, 78.3	
Gender (age ≥ 12 years)																
• Male	149	1.60	0.91	0.06-9.7, 162		60	108	26.9	18.1	0.5-184, 368		108	60.3	47.6	2.1-232, 110	
• Female	96	0.57	0.06	0.06-5.5, 92		49	73	13.9	9.2	0.79-98, 124		73	43.3	36.2	2.9-160, 55	

*The samples with zero copy number were excluded from the analysis in all categories except for the second row.

Downloaded from https://academic.oup.com/ajph/advance-article-abstract/doi/10.1093/ajph/2019.09.1600/5340011 at ASST Journals on April 19, 2019

Table 2. Multivariate linear regression analysis of predictors associated with interindividual variability of UGT2B17 protein abundance, and UGT2B17 mediated testosterone and DHT-glucuronide formation.

Dependent variable	Independent variable	Effect size (Coefficient)	β	Standard error (SE)	t-Stat	P-value
Protein abundance (pmol/mg of microsomal protein)	Intercept	-1.0		0.28	-3.5	5.8 x 10⁻⁴
	Male	0.84		0.18	4.5	9.7 x 10⁻⁶
	Adolescence	0.48		0.38	1.3	0.21
	Adulthood	1.67		0.25	4.7	4.2 x 10⁻⁶
	Diplotype H3/H4	0.95		0.25	3.8	1.8 x 10⁻⁴
	Diplotype H2/H2	1.1		0.23	4.7	4.2 x 10⁻⁶
Testosterone glucuronide formation (pmol/min/mg microsomal protein)	Copy number-2	0.22		0.22	1.1	0.31
	Intercept	-13.9		7.0	-2.0	0.05
	Male	11.7		4.0	2.9	0.004
	Adolescence	15.2		8.8	1.7	0.09
	Adulthood	21.0		6.2	3.4	9.8 x 10⁻⁴
	Diplotype H3/H4	12.7		5.5	2.3	0.02
DHT -glucuronide formation (pmol/min/mg microsomal protein)	Diplotype H2/H2	20.1		5.3	3.8	2.2 x 10⁻⁴
	Copy number-2	5.5		4.8	1.1	0.25
	Intercept	-8.5		11	-0.8	0.44
	Male	15.6		6.4	2.45	0.015
	Adolescence	27.5		14	2.0	0.05
	Adulthood	42.8		9.9	4.3	2.7 x 10⁻⁵
	Diplotype H3/H4	26.5		8.7	3.1	0.003
	Diplotype H2/H2	31.5		8.4	3.8	0.0002
	Copy number-2	2.8		7.6	0.4	0.71

Statistically significant correlations are shown in bold.

DMD # 80952




Figure 1.

DMD # 80952




Figure 2

DMD # 80952




Figure 3