

DMD # 81307

When does the rate-determining step in the hepatic clearance of a drug switch from sinusoidal uptake to all hepatobiliary clearances? Implications for predicting drug-drug interactions

Gabriela I. Patilea-Vrana and Jashvant D. Unadkat

Department of Pharmaceutics, University of Washington, Seattle, WA, USA

DMD # 81307

a) Running title: RDS switch from uptake to all hepatobiliary clearances

b) Corresponding author: Jashvant D. Unadkat, Ph.D.

Department of Pharmaceutics

University of Washington

Box 357610

Seattle, WA 98195

Phone: +1-206-685-2869

Fax: +1-206-543-3204

Email: jash@uw.edu

c) Number of:

Text Pages = 18

Tables = 1

Figures = 7

Supplemental Tables = 1

Supplemental Figures = 4

References = 43

Word in Abstract = 250/250

Words in Introduction = 522/750

Words in Discussion = 1587/1500

d) Nonstandard abbreviations: AUCR = area under the curve ratio; CL_{bile} = biliary (canalicular) efflux clearance; CL_{int} = intrinsic clearance; CL_{met} = metabolic clearance; $CL_{\text{in}}^{\text{s}}$ = sinusoidal influx clearance; $CL_{\text{ef}}^{\text{s}}$ = sinusoidal efflux clearance; DDI = drug – drug interaction; ECM = extended clearance model; IVIVE = *in vitro* to *in vivo* extrapolation; HPGL = 10^6 hepatocytes per gram

DMD # 81307

liver; NDA = new drug application; NME = new molecular entity; OATP = organic anion transporting polypeptide; $PI_{\text{met+bile}}$ = percent inhibition of $CL_{\text{met+bile}}$ necessary for RDS_{uptake} to switch to RDS_{all} ; PBPK = physiologically based pharmacokinetics; MPPGL = microsomal protein per gram liver; SCRH = sandwich cultured rat hepatocytes; SCHH = sandwich cultured human hepatocytes; RDS = rate-determining step

DMD # 81307

ABSTRACT

For dual transporter/enzyme substrate drugs, the extended clearance model (ECM) can be used to predict the rate-determining step(s) (RDS) of a drug and hence predict its drug-drug interaction (DDI) liabilities (i.e. transport, metabolism, or both). If the RDS of the hepatic clearance of the drug is sinusoidal uptake clearance (CL_{in}^s), even if the drug is mainly eliminated by hepatic metabolism, its DDI liability (as viewed from changes to systemic drug concentrations) is expected to be inhibition or induction of uptake transporters but not hepatic enzymes. However, this is true only if the condition required to maintain CL_{in}^s as the RDS is maintained. Here, we illustrate through theoretical simulations that the RDS condition may be violated in the presence of a DDI. That is, the RDS of a drug can switch from CL_{in}^s to all hepatobiliary clearances (i.e. metabolic/biliary clearance [$CL_{met+bile}$] and CL_{in}^s) leading to unexpected systemic DDI's, such as metabolic DDI's when only transporter DDI's are anticipated. As expected, these analyses revealed that the RDS switch depends on the ratio of $CL_{met+bile}$ to sinusoidal efflux clearance (CL_{ef}^s). Additional analyses revealed that for intravenously administered drugs, the RDS switch also depends on the magnitude of CL_{in}^s . We analyzed published *in vitro* quantified hepatobiliary clearances and observed that most drugs have $CL_{met+bile}/CL_{ef}^s$ ratio < 4, and hence in practice, the magnitude of CL_{in}^s must be considered when establishing the RDS. These analyses provide insights, previously not appreciated, and a theoretical framework to predict DDI liabilities for drugs that are dual transporter/enzyme substrates.

DMD # 81307

INTRODUCTION

Identifying liabilities with respect to drug – drug interactions (DDI) is important in drug development. In 2015, 25 out of the 33 new drug applications (NDA's) contained *in vitro* transporter data and out of 20 clinical trials using the new molecular entities (NME's) as victim drugs, only 9 resulted in a significant area under the curve (AUC) change (Yu et al., 2017). These data acknowledge that drug transporters are important in determining drug disposition (Giacomini et al., 2010; Hillgren et al., 2013; Patel et al., 2016).

As shown by the hepatic extended clearance model (ECM), when a drug is both transported into and metabolized/biliary excreted by the liver, the rate-determining step (RDS) in the systemic clearance of the drug can be its hepatic uptake clearance, metabolic clearance, biliary (canalicular efflux) clearance, or all hepatobiliary clearances (Miyachi et al., 1987; Sirianni and Pang, 1997; Shitara et al., 2006; Kushara and Sugiyama, 2009; Li et al., 2014; Patilea-Vrana and Unadkat, 2016). The RDS of a drug can be identified using models such as the Extended Clearance Concept Classification System (ECCCS) and the Extended Clearance Classification System (ECCS) that use the drug's *in vitro* quantified hepatobiliary clearance values or the drug's physicochemical properties, respectively (Camenisch and Umehara, 2012; Varma et al., 2015). Using such models is advantageous since the RDS of a drug helps identify where the DDI liabilities lie. Of note, unless indicated otherwise, all subsequent reference to DDI should be interpreted as those DDI that can be observed from measurement of the systemic concentrations of the victim drug. For example, if the RDS of a drug is its hepatic uptake clearance (RDS_{uptake}), then the focus of the DDI studies should be transporter-based (e.g. hepatic organic anion-transporting polypeptide (OATP) – mediated uptake of atorvastatin, (Maeda et al., 2011)) or if the RDS is both hepatic uptake and metabolic/biliary clearance (RDS_{all}), the focus of DDI studies should be all hepatobiliary pathways (e.g. OATP and CYP-mediated clearance of cerivastatin, (Mück et al., 1999; Backman et al., 2002)).

DMD # 81307

Here, we asked if knowledge of the RDS of a drug is enough to predict DDI liabilities for drugs that are dual transporter/enzyme substrates? If it is not, the focus of DDI studies will be misdirected and will result either in a negative or unexpected DDI and therefore toxicity. Under the worst-case scenario, the latter will lead to discontinuation of drug development. The end result is that both outcomes will increase drug development cost (Paul et al., 2010). For these reasons, it is important to ask: can the RDS switch from hepatic uptake clearance to all hepatobiliary clearance pathways, thus resulting in unexpected systemic DDIs? Using the ECM theory and simulations, we aimed to: i) provide a theoretical framework of when the RDS_{uptake} switches to RDS_{all} in the presence of a DDI, and ii) apply the RDS framework to predict DDI liabilities through theoretical and practical examples. The resulting analyses and simulations provide novel insights, hitherto not appreciated, into factors that determine when a victim drug experiences the RDS_{uptake} switch to RDS_{all} and elucidate important considerations when predicting DDI liabilities for drugs that are substrates of both hepatic transporters and enzymes.

DMD # 81307

MATERIALS AND METHODS

Theoretical background

The ECM describes complex hepatobiliary clearance in terms of transport at the sinusoidal membrane via sinusoidal influx (CL_{in}^s) and efflux (CL_{ef}^s), transport at the canalicular membrane via biliary efflux (CL_{bile}), metabolism (CL_{met}), hepatic blood flow (Q_h), and fraction unbound in blood (f_{ub}) (Eq. 1). CL_{in}^s and CL_{ef}^s terms incorporate both transport-mediated plus passive diffusion clearance while CL_{bile} describes active transport only. The interrelationships between the hepatobiliary clearances defined by the ECM create the RDS in the hepatic clearance of a drug. As described by us and others (Miyachi et al., 1987; Sirianni and Pang, 1997; Shitara et al., 2006; Patilea-Vrana and Unadkat, 2016), these can be: i) $RDS_{met+bile}$ when the metabolic and biliary efflux clearances of the drug are much less than sinusoidal efflux clearance ($CL_{met+bile} \ll CL_{ef}^s$) and the drug is highly permeable (passive diffusion \gg active transport, $CL_{in}^s \approx CL_{ef}^s$) and can thus rapidly distribute across the sinusoidal membrane, ii) RDS_{uptake} when the metabolic plus biliary efflux clearances are much greater than the sinusoidal efflux clearance ($CL_{met+bile} \gg CL_{ef}^s$), or iii) RDS_{all} when a drug has both active transport and metabolism but the two extreme scenario from above do not apply ($CL_{in}^s \neq CL_{ef}^s$).

$$CL_h = \frac{Q_h f_{ub} CL_{in}^s (CL_{met} + CL_{bile})}{Q_h (CL_{ef}^s + CL_{met} + CL_{bile}) + f_{ub} CL_{in}^s (CL_{met} + CL_{bile})} \quad (1)$$

Identifying the RDS of a drug can be used to predict the liability of transporter versus metabolic DDI's (see Patilea-Vrana and Unadkat, 2016 for simulations of systemic and hepatic AUC when hepatobiliary clearances are inhibited). For example, while a victim drug has RDS_{uptake} , inhibition of $CL_{met+bile}$ will not result in a significant increase in the systemic AUC even though such DDI could result in significant drug accumulation in the liver and hence potentially enhanced hepatic efficacy or toxicity of the drug. That is, from the point of view of a systemic (e.g. victim plasma concentrations) measurements, inhibition of $CL_{met+bile}$ will be incorrectly

DMD # 81307

interpreted as negative because there will be no change in systemic concentrations of the drug. On the other hand, inhibition of CL_{in}^s will result in an increase in the drug's systemic AUC (and therefore potentially non-hepatic efficacy and toxicity of the drug) but will result in no changes in the hepatic AUC provided the liver is the primary eliminating organ (see Patilea-Vrana and Unadkat, 2016 for examples). However, less appreciated is the fact that in the presence of metabolic/biliary efflux DDI, the RDS of a drug can switch from RDS_{uptake} to RDS_{all} and hence switch the DDI liability from uptake transporters to both metabolic/biliary and uptake pathways. Consequently, the drug's systemic AUC will significantly change due to metabolic and biliary efflux DDI's even though uptake was the RDS of the drug in the absence of a DDI. This would lead to unexpected DDIs as viewed from the systemic concentrations of the victim drug. Therefore, through MATLAB simulations (R2016a; MathWorks, Natick, MA), we illustrated when the RDS_{uptake} to RDS_{all} switch occurs for a victim drug in the presence of a DDI. We then applied our proposed RDS framework to published *in vitro* hepatobiliary clearances to determine if *in vivo* observed DDI liabilities can be correctly predicted. While the insights illustrated can be derived from analytical solutions of the ECM equation (Eq. 1), for clarity, we chose to use simulations to illustrate the principles of these DDI liabilities within the RDS framework.

Simulation assumptions

The hepatic ECM was simulated using the governing differential equations as previously described (Endres et al., 2009; Patilea-Vrana and Unadkat, 2016) and for simplicity, the following assumptions about the victim drug were made: i) it was administered intravenously (IV); ii) fraction unbound (f_u) in blood and tissue (liver) was set to 1; iii) liver was the only eliminating organ; iv) Q_h was set to 1 L/min. All references to systemic AUC are derived from drug concentrations in blood. Our conclusions regarding the RDS switch are generalizable to when victim drugs are administered orally but our conclusions of the RDS dependence on CL_{in}^s only apply for IV administered drugs (see text below). Furthermore, for oral drug administration,

DMD # 81307

our findings apply only to changes to the hepatic clearance/bioavailability of the victim drug and do not address the intestinal availability of the victim drug. If there is significant non-hepatic clearance, our conclusions will stand except that the magnitude of the change observed in the systemic and/or hepatic AUC of the drug will differ (Patilea-Vrana and Unadkat, 2016).

Identifying when the RDS_{uptake} switches to RDS_{all} and factors that influence this switch

First, we determined when the RDS of a drug switches from uptake clearance to all hepatobiliary clearance pathways. This requires violating the condition $CL_{\text{met+bile}} \gg CL^{\text{S}}_{\text{ef}}$, the condition necessary for uptake clearance to be the RDS in the hepatic clearance of drug. To illustrate this effect, for three theoretical victim drugs where $CL_{\text{met+bile}} \gg CL^{\text{S}}_{\text{ef}}$, ($CL_{\text{met+bile}} = 1, 10, 100$ L/min, $CL^{\text{S}}_{\text{ef}} = 0.1$ L/min, and $CL^{\text{S}}_{\text{in}} = 1 \times Q_h$), the systemic AUC ratio (AUCR) of the victim drug in the absence and presence of 10-99% inhibition of $CL_{\text{met+bile}}$ was simulated. Following the FDA guidelines, an AUCR of 1.25 was considered to be significant.

To illustrate that the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio and not the absolute magnitude of $CL_{\text{met+bile}}$ and/or $CL^{\text{S}}_{\text{ef}}$ determines when the RDS_{uptake} switches to RDS_{all} we conducted the following simulations: the systemic AUC of the drug was simulated for $CL^{\text{S}}_{\text{ef}}$ values ranging from 0.1 to 10 L/min (representing 0.1x to 10x Q_h) with $CL_{\text{met+bile}}$ set to 1-20 fold the value of the corresponding $CL^{\text{S}}_{\text{ef}}$ value. The simulated systemic AUC's when $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio was held constant were compared to the simulated systemic AUC's when $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio varied.

Next, we defined the tipping point as the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio at which RDS_{uptake} switches to RDS_{all} . Following the same strategy as above, we simulated the AUCR for various $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratios for victim drugs that originally had RDS_{uptake} to illustrate the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio at which $AUCR = 1.25$, thus signifying that RDS_{uptake} switched to RDS_{all} . The systemic AUC where the RDS is uptake was simulated such as $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio = 1000 ($AUC_{\text{ratio = control}}$, $CL_{\text{met+bile}} = 100$ L/min, $CL^{\text{S}}_{\text{ef}} = 0.1$ L/min). Then, systemic AUC was simulated for $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$

DMD # 81307

“test” ratios ranging from 0.1 – 10 ($CL_{\text{met+bile}} = 0.01 - 1 \text{ L/min}$, $CL_{\text{ef}}^s = 0.1 \text{ L/min}$) and the resulting AUC ($AUC_{\text{ratio = test}}$) was normalized to the control simulation ($AUCR = AUC_{\text{ratio = test}}/AUC_{\text{ratio = control}}$). The decrease in $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratio is akin to inhibition of $CL_{\text{met+bile}}$ since CL_{ef}^s is held constant. The $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratio which resulted in a significant change to the systemic AUC ($AUCR = 1.25$) compared to control was identified as the tipping point.

To illustrate that the magnitude of CL_{in}^s contributes to the tipping point, we simulated the tipping point for CL_{in}^s values ranging from $0.01 \times Q_h - 4 \times Q_h$ (henceforth, for simplicity, CL_{in}^s notation will be used instead of $f_u CL_{\text{in}}^s$ since $f_u = 1$). The tipping point can be explicitly derived from the ECM (Eq. 1) by defining the RDS switch for any chosen AUCR as $AUCR = RDS_{\text{uptake}}/RDS_{\text{all}}$ and solving for the $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratio (Eq. 2). This relationship (Eq. 2 with $AUCR = 1.25$) was used later to identify DDI liabilities when considering CL_{in}^s magnitude and $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratio of a drug.

$$\text{Tipping point} = \frac{1}{(AUCR - 1) \left(1 + \frac{CL_{\text{in}}^s}{Q_h}\right)} \quad (2)$$

Quantifying when a drug with RDS_{uptake} will switch to RDS_{all} due to metabolic/biliary efflux DDI's

We defined $PI_{\text{met+bile}}$ as the percent inhibition of $CL_{\text{met+bile}}$ required for RDS_{uptake} to switch to RDS_{all} . This quantifies when a significant DDI ($AUCR \geq 1.25$) will occur due to inhibition of $CL_{\text{met+bile}}$ even when uptake was the RDS in the absence of DDI. For $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratios ranging from 1-100, $CL_{\text{met+bile}}$ was inhibited 10-99%. Simulations were conducted for CL_{in}^s values = 0.25x, 1x, 4x Q_h . CL_{in}^s values were chosen to represent ER = 0.2, 0.5, and 0.8 (low, mid, and high extraction ratio (ER), respectively) and were back calculated from Eq. 3-4. The percent inhibition of $CL_{\text{met+bile}}$ at which the $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ ratio reaches the tipping point (i.e. $PI_{\text{met+bile}}$) and thus causes the RDS_{uptake} to switch to RDS_{all} was calculated as shown in Eq. 5.

$$CL_h = Q_h * ER \quad (3)$$

DMD # 81307

$$CL_h = \frac{Q_h f_{ub} CL_{in}^s}{Q_h + f_{ub} CL_{in}^s} \quad (4)$$

$$PI_{met+bile} (\%) = \frac{CL_{met+bile}/CL_{ef}^s - \text{tipping point}}{CL_{met+bile}/CL_{ef}^s} \times 100 \quad (5)$$

Applying the RDS framework to *in vitro* and *in vivo* examples

Published data sets where all hepatobiliary clearance pathways (CL_{in}^s , CL_{ef}^s , CL_{bile} , CL_{met}) were quantified *in vitro* were collected. The *in vivo* hepatobiliary clearances must be used to identify the RDS of a drug. As such, the provided *in vitro* to *in vivo* extrapolated (IVIVE) clearances were utilized; otherwise, *in vitro* hepatobiliary clearance values were scaled to *in vivo* using IVIVE scaling factors (i.e. MPPGL, HPGL, liver weight) as provided by the authors. For all drugs, $f_{ub} CL_{in}^s / Q_h$ was used to calculate the tipping point using Eq. 2 (see Results section below). RDS was labeled as RDS_{uptake} and RDS_{all} if the $CL_{met+bile} / CL_{ef}^s$ ratio was above and below the tipping point, respectively. For drugs with RDS_{uptake} , the $PI_{met+bile}$ was calculated using Eq. 5. Finally, for selected drugs, the predicted DDI liabilities using the RDS and $PI_{met+bile}$ were compared to the observed *in vivo* data. To ensure that only the systemic clearance, and not bioavailability of the victim drug was affected, clinical DDI studies were included if the victim was a dual transporter/enzyme substrate and co-administered with a selective enzyme inhibitor administered IV. It should be noted that the availability of such studies was limited.

DMD # 81307

RESULTS

Identifying the tipping point (i.e. when RDS_{uptake} switches to RDS_{all}) and factors that influence this switch

As described under theoretical background, RDS_{uptake} occurs when $CL_{\text{met+bile}} \gg CL_{\text{ef}}^{\text{s}}$, and as such, inhibition of $CL_{\text{met+bile}}$ will not manifest in the systemic AUC of a victim drug. However, when the above condition is violated due to extensive inhibition of $CL_{\text{met+bile}}$, there will be a significant increase in the systemic AUC of the victim drug when $CL_{\text{met+bile}}$ is inhibited further. In other words, when $CL_{\text{met+bile}}$ is no longer $\gg CL_{\text{ef}}^{\text{s}}$, then RDS_{uptake} switches to RDS_{all} . In Fig.1A, 84%, 98%, 99.8% inhibition of $CL_{\text{met+bile}}$ led to a clinically significant increase in the systemic AUC of the three theoretical victim drugs shown ($AUCR \geq 1.25$). Even though the victim drugs had different pre-inhibition $CL_{\text{met+bile}}$ values (1, 10, 100 L/min), the post-inhibition $CL_{\text{met+bile}}$ values were all the same (0.2 L/min). Since $CL_{\text{ef}}^{\text{s}}$ was kept constant (0.1 L/min), an $AUCR$ of 1.25 was observed when $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}} = 2$ for all three victim drugs. This simulation illustrates that the RDS_{uptake} switch to RDS_{all} depends on the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio and not the extent of $CL_{\text{met+bile}}$ inhibition.

To further emphasize the dependence on the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio, we simulated the systemic AUC of the victim drug (in the absence of DDI) for different $CL_{\text{met+bile}}$ and $CL_{\text{ef}}^{\text{s}}$ values while holding $CL_{\text{in}}^{\text{s}}$ constant. The systemic AUC remained unchanged when the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio remained fixed even though the $CL_{\text{met+bile}}$ and $CL_{\text{ef}}^{\text{s}}$ values varied, demonstrating that the RDS in the hepatic clearance of a drug is dependent on the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio and not on the absolute value of these clearances (Fig. 1B). This was true for both when $CL_{\text{met+bile}}$ was higher and lower than $CL_{\text{ef}}^{\text{s}}$ (also see Supplementary Fig. 1). Since the systemic AUC decreased as the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio increased, only the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio needs to be considered when determining when the RDS_{uptake} switches to RDS_{all} for a victim drug.

DMD # 81307

Next, we identified the tipping point, defined here as the $CL_{\text{met+bile}}/CL^{\text{s}}_{\text{ef}}$ ratio when RDS_{uptake} switches to RDS_{all} . The RDS_{uptake} switch to RDS_{all} signifies when DDI's due to inhibition of $CL_{\text{met+bile}}$ start to become significant for a victim drug that has RDS_{uptake} . As demonstrated above, the RDS_{uptake} switch to RDS_{all} depends on the $CL_{\text{met+bile}}/CL^{\text{s}}_{\text{ef}}$ ratio. As such, we identified the tipping point as the $CL_{\text{met+bile}}/CL^{\text{s}}_{\text{ef}}$ ratio at which the systemic AUC increases significantly (AUCR = 1.25) due to decrease in the $CL_{\text{met+bile}}/CL^{\text{s}}_{\text{ef}}$ ratio for a victim drug that has RDS_{uptake} (Fig. 1C). As demonstrated in Fig. 1C, the tipping point for a low, mid, and high ER drug was 3.2, 2, and 0.8, respectively.

Since the tipping point varied for a low, mid, and high ER, the magnitude of $CL^{\text{s}}_{\text{in}}$ is also an important factor in determining when the RDS_{uptake} switches to RDS_{all} (Fig. 1C). Extending the simulations to identify the tipping point across a range of $CL^{\text{s}}_{\text{in}}$ values, we established a theoretical (Eq. 2) and practical (Fig 2.) relationship between $CL^{\text{s}}_{\text{in}}/Q_h$ and the tipping point. The tipping point decreases as $CL^{\text{s}}_{\text{in}}$ increases. In other words, as a drug's $CL^{\text{s}}_{\text{in}}$ (and therefore its ER) increases, the drug is more likely to have RDS_{uptake} and a larger $PI_{\text{met+bile}}$, therefore making the drug more resistant to switching its RDS. In addition, as the influx across the sinusoidal membrane becomes large, hepatic clearance becomes limited by blood flow and therefore less likely to result in a change in AUCR when either $CL^{\text{s}}_{\text{in}}$ (or for that matter $CL_{\text{met+bile}}$) is inhibited. On the other hand, when $CL^{\text{s}}_{\text{in}}$ (or ER) is small and the hepatic clearance becomes proportional to $CL^{\text{s}}_{\text{in}}$, the victim drug becomes more susceptible to a change in RDS. This demonstrates that low ER drugs are more susceptible to RDS_{uptake} switching to RDS_{all} whereas high ER drugs are more resistant to the RDS switch.

It should be noted that the relationship between $CL^{\text{s}}_{\text{in}}/Q_h$ and the tipping point (Eq. 2 and Fig. 2) depends on the chosen AUCR cutoff. Here, an AUCR of 1.25 was chosen based off FDA guidelines of what constitutes a positive DDI. If a higher AUCR cutoff were to be selected (Supplementary Fig. 2), this would lead to estimation of lower tipping points, thus making it more

DMD # 81307

likely that drugs are labeled with RDS_{uptake} . Labeling a drug with RDS_{uptake} when in fact it has RDS_{all} can lead to underpredictions of DDI liabilities due to metabolic enzymes and biliary transporters.

By understanding the relationship between $CL^{\text{s}_{\text{in}}}$ and the tipping point, the RDS can be identified for any combination of a drug's hepatobiliary clearance values (Fig. 2). For example, a high ER drug with a $CL_{\text{met+bile}}/CL^{\text{s}_{\text{ef}}}$ ratio of 3 will have RDS_{uptake} but a low ER drug with the same $CL_{\text{met+bile}}/CL^{\text{s}_{\text{ef}}}$ ratio will have RDS_{all} . Furthermore, a drug will always have RDS_{uptake} if the $CL_{\text{met+bile}}/CL^{\text{s}_{\text{ef}}}$ ratio is greater than 4, irrespective of the magnitude of $CL^{\text{s}_{\text{in}}}$. It should be noted that for orally administered drugs, the tipping point will no longer depend on the magnitude of $CL^{\text{s}_{\text{in}}}$, and therefore will always be 4, because blood flow limitations from systemic clearance are cancelled out by blood flow limitations of hepatic bioavailability.

Quantifying the $PI_{\text{met+bile}}$ for drugs with RDS_{uptake}

Identifying the RDS of a drug as well as when the RDS_{uptake} to RDS_{all} switch will happen identifies the drug's DDI liabilities. We quantified the $PI_{\text{met+bile}}$, defined here as the percent inhibition of $CL_{\text{met+bile}}$ needed to cause the RDS_{uptake} switch to RDS_{all} , to understand when inhibition of $CL_{\text{met+bile}}$ starts to become a DDI liability for victim drugs that have RDS_{uptake} . As the $CL_{\text{met+bile}}/CL^{\text{s}_{\text{ef}}}$ ratio of the victim drug (prior to inhibition) increases, the $PI_{\text{met+bile}}$ increases (Fig. 3A). This is because as $CL_{\text{met+bile}}$ becomes \gg than $CL^{\text{s}_{\text{ef}}}$, the victim drug become resistant to the RDS_{uptake} switch to RDS_{all} . High ER drugs have a higher $PI_{\text{met+bile}}$ than low ER drugs, demonstrating again that high ER drugs are resistant to the RDS switch while low ER drugs are sensitive (Fig. 3A). Fig. 3B illustrates that while a low, mid, and high ER victim drug with $CL_{\text{met+bile}}/CL^{\text{s}_{\text{ef}}}$ ratio of 6 have RDS_{uptake} (prior to inhibition), inhibition of $CL_{\text{met+bile}}$ greater than 46%, 66%, and 87%, respectively, will cause the RDS_{uptake} to switch to RDS_{all} . This translates to observing a positive DDI due to inhibition of $CL_{\text{met+bile}}$ for a victim drug that has been identified to

DMD # 81307

have RDS_{uptake} (prior to inhibition). Without knowledge of the $PI_{\text{met+bile}}$, such a DDI may not be expected.

The purpose and conclusions of the simulations that have been used to establish the RDS framework up to this point are summarized in Fig.4. As discussed, identifying the drug's RDS is not enough to correctly predict the drug's DDI liabilities. The tipping point concept is an important consideration when identifying DDI's for victim drugs that are dual substrates of enzymes and transporters.

The flowchart in Fig. 5 can be used as a guide to identify the DDI liabilities for dual transporter/enzyme substrates. All drugs with $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio > 4 will have RDS_{uptake} whereas drugs with $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio < 4 will have RDS_{uptake} as long as this ratio is greater than the tipping point. Drugs with $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio less than the tipping point will have RDS_{all} . If the drug has RDS_{uptake} , then uptake transporters will become a DDI liability, whereas if the drug has RDS_{all} , then transporters and enzymes will be a DDI liability. However, even for drugs that have RDS_{uptake} , $CL_{\text{met+bile}}$ can become a DDI liability if inhibition of $CL_{\text{met+bile}}$ is greater than the predicted $PI_{\text{met+bile}}$ and thus causes the RDS_{uptake} switch to RDS_{all} . The flowchart identifies the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratios at which 25%, 50%, 75%, and 95% expected inhibition of $CL_{\text{met+bile}}$ is going to result in the RDS switch. This information can be used to assess when $CL_{\text{met+bile}}$ starts to become a DDI liability for drugs with RDS_{uptake} . It also helps answer the question of how much larger does $CL_{\text{met+bile}}$ need to be compared to $CL_{\text{ef}}^{\text{s}}$ in order for sinusoidal uptake clearance to become and maintain as the RDS in the hepatic clearance of any drug. Such information may be used during drug development to select drug candidates if a certain RDS is desired.

Applying the RDS framework to *in vitro* and *in vivo* examples

To provide context to the theoretical framework presented, examples from literature, where available, were utilized. For drugs with *in vitro* quantified hepatobiliary clearances that

DMD # 81307

were extrapolated to *in vivo* via IVIVE, the tipping point and $PI_{\text{met+bile}}$ were calculated using Eq. 2 and 5, and a subset of the analyzed data set, which includes primarily statin drugs, is shown in Fig. 6 (also see Supplementary Table 1) (Camenisch and Umehara, 2012; Jones et al., 2012; Varma et al., 2014; Kunze et al., 2015; Riede et al., 2017). If no empirical scaling factors (such as for active uptake clearance to match observed *in vivo* clearance) are included in the IVIVE process, then almost all drugs have RDS_{all} , except for valsartan and pravastatin (Fig. 6A). This is because most $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratios < 4 and since the IVIVE $CL^{\text{S}}_{\text{in}}$ magnitudes were small, most $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratios were less than the tipping point. Because many statins have been identified to have RDS_{uptake} , the trend in Fig. 6A suggests that $CL^{\text{S}}_{\text{in}}$ was underestimated *in vitro*. When hepatobiliary clearances were adjusted by empirical scaling factors (Varma et al., 2014) or parameters were fitted from *in vivo* IV concentration-time profiles using a PBPK model (Jones et al., 2012), the distribution of drugs is altered such as low drugs ($ER < 0.2$) tended to have RDS_{all} whereas mid and high ER drugs ($ER > 0.2$) were more likely to have RDS_{uptake} (Fig. 6B). This analysis of the published *in vitro* hepatobiliary clearances provides insight that drugs with RDS_{uptake} exist within the moderate RDS framework space, meaning that in general their $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio is < 4 and are quite susceptible to the RDS switch (Supplementary Table 1). It further elucidates that current *in vitro* quantification techniques may underestimate $CL^{\text{S}}_{\text{in}}$ which can lead to erroneous labeling of the RDS and thus incorrect DDI liability predictions (Fig 7, Supplementary Fig. 3).

To further illustrate the applicability of the RDS framework, predicted DDI liabilities using the RDS framework were compared to *in vivo* DDI examples. As indicated in Table 1, when empirical scaling factors are utilized during the IVIVE process or hepatobiliary clearances were estimated from *in vivo* via PBPK, atorvastatin and repaglinide have RDS_{uptake} and $PI_{\text{met+bile}}$ of 10-51% and 15-40%, respectively, while bosentan has RDS_{all} . For atorvastatin and repaglinide, the *in vitro* data predicted that uptake transporters (OATPs) are the primary DDI liability with the

DMD # 81307

drugs' major metabolic enzymes (CYP3A and CYP2C8, respectively) becoming a potential liability only if the *in vivo* hepatic metabolic inhibition is greater than the $PI_{\text{met+bile}}$. For bosentan, the *in vitro* data predicted that both OATPs and CYP3A4 are a potential DDI liability. Clinically, for atorvastatin, co-administration of rifampin (an OATP inhibitor) lead to AUCR of 12 whereas 33% inhibition of CYP3A4 due to IV itraconazole (as measured using CYP3A4 probe midazolam) did not change atorvastatin systemic AUC even though inhibition of atorvastatin metabolism was observed via a decrease in the 2-hydroxyatorvastatin concentrations (Maeda et al., 2011). In a similarly conducted experiment, co-administration of rifampin resulted in AUCR of 3.2 and 1.9 for bosentan or repaglinide, respectively, whereas 73% inhibition of CYP3A4 due to IV itraconazole did not significantly change the systemic AUC of these drugs (Yoshikado et al., 2017). Furthermore, repaglinide co-administered with PO rifampin and trimethoprim (CYP2C8 selective inhibitor) resulted in AUCR was 2.6 and 1.8, respectively (Kim et al., 2016). The *in vivo* DDI liability for OATPs was well predicted for all three victim drugs. The *in vivo* DDI liability for CYP3A4 was well predicted for atorvastatin. Since a probe was not used to assess the degree of CYP2C8 inhibition, it is difficult to interpret if the significant DDI when repaglinide was co-administered with trimethoprim was because RDS_{uptake} switched to RDS_{all} , or because repaglinide truly has RDS_{all} . The *in vitro* metrics as well as a whole-body PBPK DDI model suggests that repaglinide has RDS_{uptake} (Varma et al., 2013) and thus the repaglinide-trimethoprim DDI is likely due to the RDS switch. Lastly, since bosentan was predicted to have RDS_{all} , a DDI was expected due to CYP3A4 inhibition but none was observed. It should be noted that the metabolic DDI liability prediction is assuming one main drug metabolizing enzyme and no significant biliary efflux (e.g. $CL_{\text{met+bile}} = CL_{\text{CYP3A4}}$ for atorvastatin and bosentan). This assumption predicts the highest DDI risk due to inhibition of $CL_{\text{met+bile}}$ and has a higher chance of predicting false positive DDI results.

DMD # 81307

In the published *in vitro* datasets, discrepancies in the *in vitro* quantified values, particularly for CL_{in}^s , can be observed (Table 1 and Supplementary Table 1). For example, in one report the authors used empirical scaling factors for active sinusoidal uptake clearance in order to match hepatic clearance with clinical observed data that ranged from 1.1 – 101.8 with geometric mean of 10.6 (Varma et al., 2014). However, the scaling factor used severely impacted the labeling of the RDS (e.g. fluvastatin, glyburide, and pravastatin) or impacted the predicted $PI_{met+bile}$ of drugs (e.g. atorvastatin, rosuvastatin, fluvastatin, and repaglinide) (Supplementary Table 1). Assumptions regarding CL_{ef}^s also caused discrepancies. In all reports, CL_{ef}^s was assumed to be equal to passive diffusion across the sinusoidal membrane, except in one report where CL_{ef}^s was back-calculated from total SCHH CL_{int} (Camenisch and Umehara, 2012). The assumptions surrounding CL_{ef}^s impacted the $CL_{met+bile}/CL_{ef}^s$ ratio which either changed how the RDS was labeled or the magnitude of the $PI_{met+bile}$ (e.g. aliskerin, ciprofloxacin, and digoxin) (Supplementary Table 1). All in all, mispredictions of any of the hepatobiliary clearances impact the RDS labeling, magnitude of the $PI_{met+bile}$, and DDI liability predictions.

Errors from *in vitro* quantification of hepatobiliary clearances can propagate when establishing the RDS and the predicted DDI liabilities. Underprediction of both CL_{in}^s and $CL_{met+bile}$ may erroneously label a drug with RDS_{all} when it is truly RDS_{uptake} (Fig. 7). $CL_{met+bile}$ is the more sensitive parameter when determining the RDS because underpredictions of CL_{in}^s may mislabel the RDS only for drugs with $CL_{met+bile}/CL_{ef}^s$ ratio < 4 (Supplementary Fig. 4). For such drugs, even moderate (e.g. 2-5 fold) underpredictions of either clearance pathway will lead to RDS mislabeling (Supplementary Fig. 4). Furthermore, underpredictions of both CL_{in}^s and $CL_{met+bile}$ leads to underprediction of $PI_{met+bile}$, resulting in predicting a larger DDI liability due to $CL_{met+bile}$ inhibition for a drug with RDS_{uptake} (Fig. 7; Supplementary Fig. 4). While underpredictions of hepatobiliary clearances will result in conservative DDI decisions, they also increase the chances of negative DDI studies.

DMD # 81307

DISCUSSION

We built a theoretical RDS framework and identified important considerations when predicting DDI liabilities for dual transporter/enzymes substrate drugs. First, inhibition of $CL_{\text{met+bile}}$ can cause the RDS of a victim drug to switch from RDS_{uptake} to RDS_{all} and hence result in an unexpected systemic DDI. Two metrics have been developed to identify when the RDS switch occurs: the tipping point, defined as the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio at which RDS_{uptake} will switch to RDS_{all} and the $PI_{\text{met+bile}}$, defined as the percent inhibition of $CL_{\text{met+bile}}$ at which a significant AUC change ($AUCR > 1.25$) for a drug with RDS_{uptake} will start to be observed. The tipping point depends on the drug's $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio and on the magnitude of $CL^{\text{S}}_{\text{in}}$. The former but not latter condition is relevant when victim drugs are administered orally. Second, we showed that the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio must be > 4 in order for any drug to have RDS_{uptake} . Third, we applied the RDS framework to *in vitro* quantified hepatobiliary clearances and observed that most drugs have $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio < 4 , and hence in practice, the magnitude of $CL^{\text{S}}_{\text{in}}$ must be considered when establishing the RDS.

Our theoretical analysis demonstrates that the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio and not the absolute magnitudes of the clearances determines the RDS in the hepatic clearance of a drug. Previous publications allude to this relationship. The authors of the ECCCS observed through experimental data that when $CL_{\text{met+bile}}$ is $2 \times CL^{\text{S}}_{\text{ef}}$, drugs that have RDS_{uptake} can be separated from those that do not (Riede et al., 2016). Furthermore, β value ($\beta = CL_{\text{met+bile}} / (CL_{\text{met+bile}} + CL^{\text{S}}_{\text{ef}})$) introduced by Yoshikado et al., can be used to differentiate the RDS, such as when β approaches unity (i.e. $CL_{\text{met+bile}} \gg CL^{\text{S}}_{\text{ef}}$), a drug has RDS_{uptake} (Yoshikado et al., 2016). Our analyses corroborate and expand upon these results to provide a quantitative definition of the demarcation point between RDS_{uptake} and RDS_{all} , i.e. the tipping point, and illustrate that the magnitude of $CL^{\text{S}}_{\text{in}}$ in addition to the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio is an important factor in determining the

DMD # 81307

RDS of a drug. That is, as a drug's CL_{in}^s value increases, the drug is more likely to have RDS_{uptake} and to become resistant to the RDS_{uptake} switch to RDS_{all} .

We found good agreement for atorvastatin *in vivo* predicted DDI liabilities (Table 1). For bosentan, the overprediction of expected DDI due to inhibition of $CL_{met+bile}$ may be due to errors in the quantification of the hepatobiliary clearances. Indeed, a study in cynomolgus monkey where bosentan plasma and liver drug concentrations were quantified found that the *in vitro* scaled CL_{in}^s and CL_{met} were 28 and 13-fold underpredicted while CL_{ef}^s (assumed equal to passive diffusion) was overpredicted by 2-fold when compared to the *in vivo* fitted values (Morse et al., 2017). Combining the *in vitro* metrics that identify RDS_{uptake} for repaglinide with *in vivo* repaglinide DDI's, it appears CYP2C8 but not CYP3A4 inhibition may lead to RDS_{uptake} switch to RDS_{all} . Indeed, inhibition of repaglinide with gemfibrozil (CYP2C8 and OATP1B1 inhibitor) led to an 8-fold increase in systemic AUC while co-administration of itraconazole or cyclosporine (OATP1B1 and CYP3A4 inhibitor) led to much more modest 1.4 and 2.4-fold increase in systemic AUC (Niemi et al., 2003; Kajosaari et al., 2005).

The DDI liabilities discussed so far are relevant for systemic drug exposure but not necessarily for hepatic drug exposure and thus efficacy/toxicity if the site of action is in the liver. For example, the LDL cholesterol lowering effect mediated by atorvastatin does not change for subjects with OATP1B1 polymorphism c.521T>C even though there is a significant increase in atorvastatin systemic AUC (Maeda, 2015). This is because if the liver is the main eliminating organ, changes to sinusoidal uptake alters the hepatic concentration-time profile but not the hepatic AUC. However, systemic increase of atorvastatin may lead to off-target toxicity, such as muscle myopathy. We refer the readers to our previous publication (Patilea-Vrana and Unadkat, 2016), where simulations demonstrate the impact of inhibition of uptake or metabolism on both systemic and hepatic AUC when the liver is and is not the main eliminating organ.

DMD # 81307

The contrast between *in vitro* quantified CL_{in}^s with and without empirical scaling factors in Fig. 6 demonstrates that IVIVE of accurate transporter mediated clearance remains challenging (Chu et al., 2013; Feng et al., 2014). The system used for *in vitro* quantification may be crucial, since CL_{in}^s for statins quantified in SCHH appeared to be lower in magnitude than when quantified in suspended hepatocytes (Table 1 and Supplementary Table 1). This may be mediated by significant intracellular localization of plasma membrane transporters (Kumar et al., 2017), or high interindividual variability when using individual donors (Vildhede et al., 2014). These reasons may also cause underpredictions of CL_{in}^s or CL_{bile} . For transporter IVIVE, we have previously recommended using a bottom-up proteomic approach and adjusting for *in vitro* activity via *in vitro* to *in vivo* transporter expression based scaling factors (Prasad and Unadkat, 2014). We have recently demonstrated the successful prediction of hepatobiliary clearance of rosuvastatin in rat using the aforementioned approach (Ishida et al., 2017).

Special emphasis needs to be put on quantifying CL_{ef}^s along with $CL_{met+bile}$ since the $CL_{met+bile}/CL_{ef}^s$ ratio is one of the anchor points when establishing the RDS. Because CL_{ef}^s is a difficult parameter to quantify *in vitro*, it is typically assumed to be equal to passive diffusion across the sinusoidal membrane. However, there are examples of active sinusoidal efflux transport, such as MRP3 efflux of rosuvastatin (Pfeifer et al., 2013). Active sinusoidal efflux would increase the magnitude of CL_{ef}^s and decrease the $CL_{met+bile}/CL_{ef}^s$ ratio, making a drug more likely to have RDS_{all} . One approach to measuring CL_{ef}^s is to use an integrative temporal modeling approach in SCRH (Pfeifer et al., 2013; Ishida et al., 2017).

Errors in the quantification of CL_{in}^s and/or the $CL_{met+bile}/CL_{ef}^s$ ratio can impact DDI liability predictions. For example, patients with OATP1B1 polymorphism c.521T>C have a ~2-fold higher atorvastatin AUC compared to the wild type allele (Maeda, 2015). Because of the lower CL_{in}^s and therefore higher susceptibility to the RDS_{uptake} to RDS_{all} switch, patients with OATP1B1 polymorphism may experience a DDI due to inhibition of CYP3A whereas patients with the wild-

DMD # 81307

type allele may not. The same trend would be true for patients with polymorphic enzymes that result in lower $CL_{\text{met+bile}}$ and thus lower $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratios. Polypharmacy use can also impact DDI liability predictions. For example, highly active antiretroviral therapy (HAART) typically includes potent CYP3A4 and moderate OATP inhibitor ritonavir among other drugs, which can impact the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio more severely than if only one drug is administered. Indeed, the systemic AUC of atorvastatin increased 3.9 and 9.4-fold when co-administered with saquinavir/ritonavir and tipranavir/ritonavir, respectively (Fichtenbaum et al., 2002; Pham et al., 2009). Lastly, saturation of enzymes, leading to a lower CL_{met} with increased dose, may lower the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio and cause DDI's due to the RDS_{uptake} switch to RDS_{all} .

If a victim drug has RDS_{all} but it has been mislabeled as RDS_{uptake} , then the DDI liability due to inhibition of both transporter and metabolic activity could be underestimated. Considering potential DDI risks, it would be most conservative to assume a drug has RDS_{all} ; however, making such an assumption would lead to an increase in negative DDI studies, particularly when conducting metabolic/biliary efflux DDI studies if the drug has RDS_{uptake} . An analysis of the DDI's performed for a cohort of NME's in 2013 showed a modest return on investment because 57% (n=141) of all *in vivo* DDI's were negative (Lesko and Lagishetty, 2016). Given the high prevalence of negative DDI's, it may be more appropriate to make mechanistic-based rather than conservative decisions with regard to DDI liabilities.

The RDS framework presented here should be used as a guide when identifying the DDI liabilities whereas PBPK models should be used to predict the direction and magnitude of complex transporter-enzyme DDI's. Several examples of such models (e.g. repaglinide, simvastatin, rosuvastatin) exist that predict complex interactions due to chemical inhibition or genetic polymorphism (Varma et al., 2013; Rose et al., 2014; Tsamandouras et al., 2015). Even with PBPK models, there are limitations. For example, when a drug has RDS_{uptake} , the $CL_{\text{met+bile}}$ is unidentifiable from plasma concentrations data since only $CL_{\text{in}}^{\text{s}}$ plays a significant role in

DMD # 81307

determining hepatic clearance. Focusing on capturing the correct $CL_{\text{met+bile}}$ magnitude and not the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio can be misleading and will impact PBPK predictions. For instance, in an atorvastatin PBPK model, when cyclosporine CYP3A4 K_i was modulated 100-fold, a maximum 1.6-fold AUCR was achieved (Duan et al., 2017). While the tendency is to run sensitivity analysis on the active components (transport and metabolism), a sensitivity analysis on $CL_{\text{ef}}^{\text{s}}$ value (in the model it was assumed to be equal to passive diffusion) should also be run as, for the specific example provided, it would likely have revealed a larger impact of cyclosporine on atorvastatin systemic AUC. Such an analysis may be helpful in consolidating *in vitro* K_i data with observed *in vivo* DDI data.

In summary, we introduced a theoretical RDS framework to better predict DDI liabilities for drugs that are dual transporter/enzyme substrates. We provide useful insights, such as: i) the RDS_{uptake} switch to RDS_{all} depends on the ratio of $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ and the magnitude of $CL_{\text{in}}^{\text{s}}$, ii) $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio > 4 ensures RDS_{uptake} independent of $CL_{\text{in}}^{\text{s}}$ magnitude or administration route iii) existing drugs exist within a moderate space within the RDS framework and are susceptible to the RDS_{uptake} switch to RDS_{all} . While the above insights were obtained from the hepatic ECM, they can be equally applied to other organs such as the kidneys where vectorial (basal to apical) transport of drugs is possible.

DMD # 81307

ACKNOWLEDGEMENTS

We would like to thank Bhagwat Prasad and Neha Maharao for insightful and constructive comments/suggestions. We would also like to thank the anonymous reviewers, who gave excellent feedback to help improve the clarity and significance of this work and whose suggestions prompted us to derive the explicit solution for the tipping point.

DMD # 81307

AUTHORSHIP CONTRIBUTIONS

Participated in research design: Patilea-Vrana and Unadkat

Conducted experiments: Patilea-Vrana

Contributed new reagents or analytical tool: Patilea-Vrana

Performed data analysis: Patilea-Vrana

Wrote or contributed to the writing of the manuscript: Patilea-Vrana and Unadkat

DMD # 81307

REFERENCES

- Backman JT, Kyrklund C, Neuvonen M, and Neuvonen PJ (2002) Gemfibrozil greatly increases plasma concentrations of cerivastatin. *Clin Pharmacol Ther* **72**:685-691.
- Camenisch G and Umehara K (2012) Predicting human hepatic clearance from in vitro drug metabolism and transport data: a scientific and pharmaceutical perspective for assessing drug-drug interactions. *Biopharm Drug Dispos* **33**:179-194.
- Chu X, Korzekwa K, Elsby R, Fenner K, Galetin A, Lai Y, Matsson P, Moss A, Nagar S, Rosania GR, Bai JP, Polli JW, Sugiyama Y, and Brouwer KL (2013) Intracellular drug concentrations and transporters: measurement, modeling, and implications for the liver. *Clin Pharmacol Ther* **94**:126-141.
- Duan P, Zhao P, and Zhang L (2017) Physiologically Based Pharmacokinetic (PBPK) Modeling of Pitavastatin and Atorvastatin to Predict Drug-Drug Interactions (DDIs). *Eur J Drug Metab Pharmacokinet* **42**:689-705.
- Endres CJ, Endres MG, and Unadkat JD (2009) Interplay of drug metabolism and transport: a real phenomenon or an artifact of the site of measurement? *Mol Pharm* **6**:1756-1765.
- Feng B, Varma MV, Costales C, Zhang H, and Tremaine L (2014) In vitro and in vivo approaches to characterize transporter-mediated disposition in drug discovery. *Expert Opin Drug Discov* **9**:873-890.
- Fichtenbaum CJ, Gerber JG, Rosenkranz SL, Segal Y, Aberg JA, Blaschke T, Alston B, Fang F, Kosel B, Aweeka F, and Group NACT (2002) Pharmacokinetic interactions between protease inhibitors and statins in HIV seronegative volunteers: ACTG Study A5047. *AIDS* **16**:569-577.
- Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, Hillgren KM, Hoffmaster KA, Ishikawa T, Keppler D, Kim RB, Lee CA, Niemi M, Polli JW, Sugiyama Y, Swaan PW, Ware JA, Wright SH, Yee SW, Zamek-

DMD # 81307

- Gliszczynski MJ, Zhang L, and Consortium IT (2010) Membrane transporters in drug development. *Nat Rev Drug Discov* **9**:215-236.
- Hillgren KM, Keppler D, Zur AA, Giacomini KM, Stieger B, Cass CE, Zhang L, and Consortium IT (2013) Emerging transporters of clinical importance: an update from the International Transporter Consortium. *Clin Pharmacol Ther* **94**:52-63.
- Ishida K, Ullah M, Toth B, Juhasz V, and Unadkat JD (2017) Successful prediction of in vivo hepatobiliary clearances and hepatic concentrations of rosuvastatin using sandwich-cultured rat hepatocytes, transporter-expressing cell lines, and quantitative proteomics. *Drug Metab Dispos*.
- Jones HM, Barton HA, Lai Y, Bi YA, Kimoto E, Kempshall S, Tate SC, El-Kattan A, Houston JB, Galetin A, and Fenner KS (2012) Mechanistic pharmacokinetic modeling for the prediction of transporter-mediated disposition in humans from sandwich culture human hepatocyte data. *Drug Metab Dispos* **40**:1007-1017.
- Kajosaari LI, Niemi M, Neuvonen M, Laitila J, Neuvonen PJ, and Backman JT (2005) Cyclosporine markedly raises the plasma concentrations of repaglinide. *Clin Pharmacol Ther* **78**:388-399.
- Kim SJ, Yoshikado T, Ieiri I, Maeda K, Kimura M, Irie S, Kusuhara H, and Sugiyama Y (2016) Clarification of the Mechanism of Clopidogrel-Mediated Drug-Drug Interaction in a Clinical Cassette Small-dose Study and Its Prediction Based on In Vitro Information. *Drug Metab Dispos* **44**:1622-1632.
- Kumar V, Nguyen TB, Tóth B, Juhasz V, and Unadkat JD (2017) Optimization and Application of a Biotinylation Method for Quantification of Plasma Membrane Expression of Transporters in Cells. *AAPS J*.
- Kunze A, Poller B, Huwyler J, and Camenisch G (2015) Application of the extended clearance concept classification system (ECCCS) to predict the victim drug-drug interaction potential of statins. *Drug Metabol Personal Ther* **30**:175-188.

DMD # 81307

Kusuhara H and Sugiyama Y (2009) In vitro-in vivo extrapolation of transporter-mediated clearance in the liver and kidney. *Drug Metab Pharmacokinet* **24**:37-52.

Lesko LJ and Lagishetty CV (2016) Are We Getting the Best Return on Investment From Clinical Drug-Drug Interaction Studies? *J Clin Pharmacol* **56**:555-558.

Li R, Barton HA, and Varma MV (2014) Prediction of pharmacokinetics and drug-drug interactions when hepatic transporters are involved. *Clin Pharmacokinet* **53**:659-678.

Maeda K (2015) Organic anion transporting polypeptide (OATP)1B1 and OATP1B3 as important regulators of the pharmacokinetics of substrate drugs. *Biol Pharm Bull* **38**:155-168.

Maeda K, Ikeda Y, Fujita T, Yoshida K, Azuma Y, Haruyama Y, Yamane N, Kumagai Y, and Sugiyama Y (2011) Identification of the rate-determining process in the hepatic clearance of atorvastatin in a clinical cassette microdosing study. *Clin Pharmacol Ther* **90**:575-581.

Miyauchi S, Sugiyama Y, Sawada Y, Morita K, Iga T, and Hanano M (1987) Kinetics of hepatic transport of 4-methylumbelliferone in rats. Analysis by multiple indicator dilution method. *J Pharmacokinet Biopharm* **15**:25-38.

Morse BL, MacGuire JG, Marino AM, Zhao Y, Fox M, Zhang Y, Shen H, Griffith Humphreys W, Marathe P, and Lai Y (2017) Physiologically Based Pharmacokinetic Modeling of Transporter-Mediated Hepatic Clearance and Liver Partitioning of OATP and OCT Substrates in Cynomolgus Monkeys. *AAPS J* **19**:1878-1889.

Mück W, Mai I, Fritsche L, Ochmann K, Rohde G, Unger S, Johne A, Bauer S, Budde K, Roots I, Neumayer HH, and Kuhlmann J (1999) Increase in cerivastatin systemic exposure after single and multiple dosing in cyclosporine-treated kidney transplant recipients. *Clin Pharmacol Ther* **65**:251-261.

Niemi M, Backman JT, Neuvonen M, and Neuvonen PJ (2003) Effects of gemfibrozil, itraconazole, and their combination on the pharmacokinetics and pharmacodynamics of

DMD # 81307

repaglinide: potentially hazardous interaction between gemfibrozil and repaglinide.

Diabetologia **46**:347-351.

Patel M, Taskar KS, and Zamek-Gliszczyński MJ (2016) Importance of Hepatic Transporters in Clinical Disposition of Drugs and Their Metabolites. *J Clin Pharmacol* **56 Suppl 7**:S23-39.

Patilea-Vrana G and Unadkat JD (2016) Transport vs. Metabolism: What Determines the Pharmacokinetics and Pharmacodynamics of Drugs? Insights From the Extended Clearance Model. *Clin Pharmacol Ther* **100**:413-418.

Paul SM, Mytelka DS, Dunwiddie CT, Persinger CC, Munos BH, Lindborg SR, and Schacht AL (2010) How to improve R&D productivity: the pharmaceutical industry's grand challenge. *Nat Rev Drug Discov* **9**:203-214.

Pfeifer ND, Yang K, and Brouwer KL (2013) Hepatic basolateral efflux contributes significantly to rosuvastatin disposition I: characterization of basolateral versus biliary clearance using a novel protocol in sandwich-cultured hepatocytes. *J Pharmacol Exp Ther* **347**:727-736.

Pham PA, la Porte CJ, Lee LS, van Heeswijk R, Sabo JP, Elgadi MM, Piliero PJ, Barditch-Crovo P, Fuchs E, Flexner C, and Cameron DW (2009) Differential effects of tipranavir plus ritonavir on atorvastatin or rosuvastatin pharmacokinetics in healthy volunteers. *Antimicrob Agents Chemother* **53**:4385-4392.

Prasad B and Unadkat JD (2014) Optimized approaches for quantification of drug transporters in tissues and cells by MRM proteomics. *AAPS J* **16**:634-648.

Riede J, Poller B, Huwyler J, and Camenisch G (2017) Assessing the Risk of Drug-Induced Cholestasis Using Unbound Intrahepatic Concentrations. *Drug Metab Dispos* **45**:523-531.

DMD # 81307

- Riede J, Poller B, Umehara K, Huwyler J, and Camenisch G (2016) New IVIVE method for the prediction of total human clearance and relative elimination pathway contributions from in vitro hepatocyte and microsome data. *Eur J Pharm Sci* **86**:96-102.
- Rose RH, Neuhoff S, Abduljalil K, Chetty M, Rostami-Hodjegan A, and Jamei M (2014) Application of a Physiologically Based Pharmacokinetic Model to Predict OATP1B1-Related Variability in Pharmacodynamics of Rosuvastatin. *CPT Pharmacometrics Syst Pharmacol* **3**:e124.
- Shitara Y, Horie T, and Sugiyama Y (2006) Transporters as a determinant of drug clearance and tissue distribution. *Eur J Pharm Sci* **27**:425-446.
- Sirianni GL and Pang KS (1997) Organ clearance concepts: new perspectives on old principles. *J Pharmacokinet Biopharm* **25**:449-470.
- Tsamandouras N, Dickinson G, Guo Y, Hall S, Rostami-Hodjegan A, Galetin A, and Aarons L (2015) Development and Application of a Mechanistic Pharmacokinetic Model for Simvastatin and its Active Metabolite Simvastatin Acid Using an Integrated Population PBPK Approach. *Pharm Res* **32**:1864-1883.
- Varma MV, Bi YA, Kimoto E, and Lin J (2014) Quantitative prediction of transporter- and enzyme-mediated clinical drug-drug interactions of organic anion-transporting polypeptide 1B1 substrates using a mechanistic net-effect model. *J Pharmacol Exp Ther* **351**:214-223.
- Varma MV, Lai Y, Kimoto E, Goosen TC, El-Kattan AF, and Kumar V (2013) Mechanistic modeling to predict the transporter- and enzyme-mediated drug-drug interactions of repaglinide. *Pharm Res* **30**:1188-1199.
- Varma MV, Steyn SJ, Allerton C, and El-Kattan AF (2015) Predicting Clearance Mechanism in Drug Discovery: Extended Clearance Classification System (ECCS). *Pharm Res* **32**:3785–3802.

DMD # 81307

Vildhede A, Karlgren M, Svedberg EK, Wisniewski JR, Lai Y, Norén A, and Artursson P (2014)

Hepatic uptake of atorvastatin: influence of variability in transporter expression on uptake clearance and drug-drug interactions. *Drug Metab Dispos* **42**:1210-1218.

Yoshikado T, Maeda K, Kusuhara H, Furihata KI, and Sugiyama Y (2017) Quantitative Analyses

of the Influence of Parameters Governing Rate-Determining Process of Hepatic Elimination of Drugs on the Magnitudes of Drug-Drug Interactions via Hepatic OATPs and CYP3A Using Physiologically Based Pharmacokinetic Models. *J Pharm Sci* **106**:2739-2750.

Yoshikado T, Yoshida K, Kotani N, Nakada T, Asaumi R, Toshimoto K, Maeda K, Kusuhara H,

and Sugiyama Y (2016) Quantitative Analyses of Hepatic OATP-Mediated Interactions Between Statins and Inhibitors Using PBPK Modeling With a Parameter Optimization Method. *Clin Pharmacol Ther* **100**:513-523.

Yu J, Zhou Z, Owens KH, Ritchie TK, and Ragueneau-Majlessi I (2017) What Can Be Learned

from Recent New Drug Applications? A Systematic Review of Drug Interaction Data for Drugs Approved by the US FDA in 2015. *Drug Metab Dispos* **45**:86-108.

DMD # 81307

FOOTNOTES

This research was supported in part by National Institutes of Health National Institute of Drug Abuse [P01DA032507].

DMD # 81307

FIGURE LEGENDS

Figure 1. Identifying when RDS_{uptake} switches to RDS_{all} , i.e. the tipping point. A) Extensive inhibition of $CL_{\text{met+bile}}$ can lead to a significant increase in the systemic AUC for three theoretical victim drugs that have RDS_{uptake} (i.e. $CL_{\text{met+bile}} \gg CL_{\text{ef}}^{\text{s}}$) in the absence of DDI. When inhibition of $CL_{\text{met+bile}}$ eventually violates the condition required for RDS_{uptake} , the RDS_{uptake} switches to RDS_{all} . An AUCR ≥ 1.25 was observed when $CL_{\text{met+bile}}$ was inhibited $\geq 84\%$, $\geq 98\%$, and $\geq 99.8\%$ for $CL_{\text{met+bile}} = 1, 10, 100$ L/hr, respectively. However, for all three victim drugs, the $CL_{\text{met+bile}}$ value after such inhibition was similar (0.2 L/min) as was the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio (= 2). Simulations were performed as follows: $CL_{\text{in}}^{\text{s}} = 1xQ_{\text{h}}$, $CL_{\text{met+bile}} = 1, 10, 100$ L/hr, $CL_{\text{ef}}^{\text{s}} = 0.1$ L/min. **B)** The systemic AUC (in the absence of any DDI) of a theoretical drug remains unchanged when the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio remains fixed (blue bars) but not when the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio is varied (yellow bars) even though the absolute value of $CL_{\text{met+bile}}$ and $CL_{\text{ef}}^{\text{s}}$ is varied in both scenarios. This trend was observed irrespective of the value of $CL_{\text{in}}^{\text{s}}$ (Supplementary Fig. 1). Furthermore, this trend is true for when $CL_{\text{met+bile}} > CL_{\text{ef}}^{\text{s}}$ or $CL_{\text{met+bile}} < CL_{\text{ef}}^{\text{s}}$ (also refer to Supplementary Fig. 1). Thus, the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio, irrespective of the magnitude of the absolute values of these clearances, is important for establishing the RDS and henceforth when the RDS switches from uptake to all hepatobiliary clearances. Simulations were performed as follows: $CL_{\text{in}}^{\text{s}} = 0.25xQ_{\text{h}}$ and the other input clearance values for scenarios A-E are shown in the table provided. **C)** Since the RDS depends on the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio, we define the tipping point as the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio at which RDS_{uptake} switches to RDS_{all} . Similar to panel A, the RDS_{uptake} switch to RDS_{all} is represented by an AUCR of 1.25 and the decrease in $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio is akin to inhibition of $CL_{\text{met+bile}}$ when $CL_{\text{ef}}^{\text{s}}$ is kept constant. As shown by the gray arrows, the tipping point for a low, mid, and high ER drug is 3.2, 2, and 0.8, respectively. For example, if the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio for the low ER drug is above the tipping point (i.e. $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}} > 3.2$), then the drug will have RDS_{uptake} and therefore DDI's due to

DMD # 81307

CL_{in}^s but not $CL_{met+bile}$ should be expected. However, inhibition of $CL_{met+bile}$ that makes the $CL_{met+bile}/CL_{ef}^s$ ratio lower than the tipping point (i.e. $CL_{met+bile}/CL_{ef}^s < 3.2$) will lead to a significant increase in the systemic AUC. Crossing the tipping point is indicative of the RDS_{uptake} switch to RDS_{all} . Simulations were performed as follows: systemic AUC were simulated for $CL_{in}^s = 0.25x, 1x, 4xQ_h$ (representing low, mid, and high ER, respectively) and $CL_{met+bile}/CL_{ef}^s$ ratios from 1-10, then normalized to a control simulation where the $CL_{met+bile}/CL_{ef}^s$ ratio was set to 1000 (i.e. RDS_{uptake}).

Figure 2. The RDS framework helps identify DDI liabilities. The $CL_{met+bile}/CL_{ef}^s$ ratio and CL_{in}^s magnitude of a drug determines the RDS of the drug and when RDS_{uptake} switches to RDS_{all} . Combinations of hepatobiliary clearances found in the shaded area have RDS_{all} while those in the non-shaded area have RDS_{uptake} . Any alterations in hepatobiliary clearances that causes a drug to switch from the non-shaded to the shaded area will cross the tipping point (dashed line – Eq. 2) and therefore switch the RDS from uptake to all hepatobiliary clearances. The consequence of this switch is that DDI's due to inhibition of $CL_{met+bile}$ will now manifest in the systemic AUC of a victim drug that originally had RDS_{uptake} . Consistent with Fig. 1C, the tipping point decreases as the magnitude of CL_{in}^s (and therefore the drug's ER) increase. This suggests that the greater the ER of the drug, the more likely it will have RDS_{uptake} and it will be more resistant to switch to RDS_{all} . Furthermore, when $CL_{met+bile}/CL_{ef}^s > 4$, the RDS will always be uptake clearance irrespective of the value of CL_{in}^s/Q_h . However, when $CL_{met+bile}/CL_{ef}^s < 4$, the RDS can be either uptake or all hepatobiliary pathways depending on the magnitude of CL_{in}^s . It should be noted that if a drug is administered orally, the tipping point will always be 4 because the blood flow limitations are no longer relevant. Simulations were performed as follows: the tipping point was simulated for CL_{in}^s values ($0.01xQ_h - 4xQ_h$) using Eq. 2.

DMD # 81307

Figure 3. Identifying when drugs with RDS_{uptake} will start to experience a DDI due to inhibition of $CL_{\text{met+bile}}$. **A)** The $PI_{\text{met+bile}}$, defined as the % inhibition of $CL_{\text{met+bile}}$ required for the RDS_{uptake} switch to RDS_{all} , depends on the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio (prior to inhibition) and the magnitude of $CL^{\text{S}}_{\text{in}}$ (represented as low, mid, and high ER drugs). The $PI_{\text{met+bile}}$ identifies when a positive DDI due to inhibition of $CL_{\text{met+bile}}$ for a drug with RDS_{uptake} would be expected. Lower $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratios as well as low ER drugs are the most susceptible for the RDS_{uptake} switch to RDS_{all} due to $CL_{\text{met+bile}}$ inhibition. **B)** In order for RDS_{uptake} to switch to RDS_{all} for a theoretical victim drug with $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio of 6, $CL_{\text{met+bile}}$ must be inhibited by >46%, >66%, or >87% if the drug is low, mid, and high ER, respectively. Visually, the RDS_{uptake} switch to RDS_{all} happens when the theoretical victim drug crosses the dashed line (the tipping point) from the unshaded area (RDS_{uptake}) to the shaded area (RDS_{all}). Additional examples of $PI_{\text{met+bile}}$ are given in the table provided. Simulations were performed as follows: $PI_{\text{met+bile}}$ was calculated using Eq. 5 for $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratios ranging 1-40 and for $CL^{\text{S}}_{\text{in}} = 0.25x, 1x, 4xQ_h$ (representing low, mid, and high ER, respectively).

Figure 4. Summary of the purpose and conclusions for the simulations used to establish the RDS framework.

Figure 5. Applying the RDS framework to identify DDI liabilities for dual transporter/enzyme substrate drugs. If $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}} > 4$, then the drug will have RDS_{uptake} , irrespective of the magnitude of $CL^{\text{S}}_{\text{in}}$. For drugs with RDS_{uptake} , DDI's due to inhibition of $CL_{\text{met+bile}}$ can become significant depending on the drug's $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio and the expected inhibition of $CL_{\text{met+bile}}$. For example, 50% inhibition of $CL_{\text{met+bile}}$ may result in a significant DDI for a drug with RDS_{uptake} and $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio < 8 but no DDI will be observed if the drug has $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio > 8. The DDI liability due to inhibition of $CL_{\text{met+bile}}$ increases as the $CL_{\text{met+bile}}/CL^{\text{S}}_{\text{ef}}$ ratio decrease and the expected $CL_{\text{met+bile}}$ inhibition increases.

DMD # 81307

Figure 6. The distribution of drugs within the RDS framework using hepatobiliary clearance quantified *in vitro* and extrapolated to *in vivo*. Published *in vitro* hepatobiliary clearance values, when extrapolated to *in vivo* via IVIVE, can identify the RDS based on $f_u \text{CL}^{\text{s}_{\text{in}}}/Q_h$ and $\text{CL}_{\text{met+bile}}/\text{CL}^{\text{s}_{\text{ef}}}$ ratio (Eq. 2). **A)** When no empirical scaling factors, such as to scale up active transport, are applied during the IVIVE process, all drugs except for valsartan and pravastatin have RDS_{all} . Most drugs had $\text{CL}_{\text{met+bile}}/\text{CL}^{\text{s}_{\text{ef}}}$ ratio < 4 , indicating drugs primarily exist within the moderate RDS framework space. Furthermore, most drugs have $f_u \text{CL}^{\text{s}_{\text{in}}}/Q_h < 0.4$, indicating severe underprediction of $\text{CL}^{\text{s}_{\text{in}}}$. **B)** When empirical scaling factors are used or hepatobiliary clearances are estimates from *in vivo* data using PBPK modeling, the RDS of the drugs is altered severely. Now, $\text{RDS}_{\text{uptake}}$ occurs more often for mid and high ER drugs with RDS_{all} primarily for low ER drugs (ER was calculated from *in vivo* hepatic clearance and blood flow). Furthermore, since all drugs have $\text{CL}_{\text{met+bile}}/\text{CL}^{\text{s}_{\text{ef}}}$ ratio < 4 , information about both the magnitude of $f_u \text{CL}^{\text{s}_{\text{in}}}$ and the $\text{CL}_{\text{met+bile}}/\text{CL}^{\text{s}_{\text{ef}}}$ ratio is necessary to correctly predict DDI liabilities. The dashed line represents the tipping point (Eq. 2). The data shown are from Jones et al., 2012 and Varma et al., 2014, and represent a subset of the complete data set presented in Supplementary Table 1.

Figure 7. The impact of underpredictions of hepatobiliary clearance on DDI liability predictions. A representative 3-fold underprediction of either **A)** $\text{CL}^{\text{s}_{\text{in}}}$ or **B)** $\text{CL}_{\text{met+bile}}$ can lead to erroneous labeling of the RDS for low, mid and high ER drugs (shown by the filled circles crossing from the non-shaded to shaded area, i.e. $\text{RDS}_{\text{uptake}}$ switches to RDS_{all}). Mislabeling the RDS impacts the expected DDI risk due to transporters versus enzymes. Furthermore, underpredictions of either $\text{CL}^{\text{s}_{\text{in}}}$ or $\text{CL}_{\text{met+bile}}$ leads to identifying both transporters and enzymes as DDI liabilities when truly only uptake transporters are the true DDI liability. Please refer to Supplementary Fig. 4 for more detailed simulations.

Table 1. Comparison of predicted DDI liabilities from *in vitro* data to *in vivo* clinical studies

Hepatobiliary clearances, following IVIVE, can be used to identify the RDS of a drug, such as if the $CL_{met+bile} / CL_{ef}^{s_{ef}}$ ratio is > or < than the tipping point (Eq. 2), then the drug will have RDS_{uptake} or RDS_{all} , respectively. If a drug has RDS_{uptake} , then the $PI_{met+bile}$ can be quantified (Eq. 5) in order to predict when a significant DDI should be expected due to inhibition of metabolic/biliary efflux clearance. An expanded analysis is shown in Supplementary Table 1.

| Drug | Hepatobiliary clearance (ml/min/kg) | | | | $f_u CL_{in}^{s_{in}}$ /Qh ^f | $CL_{met+bile} / CL_{ef}^{s_{ef}}$ | Tipping point | $PI_{met+bile}$ | RDS <i>in vitro</i> | RDS <i>in vivo</i> | REF |
|--------------|-------------------------------------|--------------------|------------|------------------|---|------------------------------------|---------------|-----------------|---------------------|---|---|
| | $CL_{in}^{s_{in}}$ | $CL_{ef}^{s_{ef}}$ | CL_{met} | CL_{bile} | | | | | | | |
| Atorvastatin | 61 ^a | 24.9 | 58.3 | 4.3 | 0.12 | 2.52 | 3.59 | -- | all | uptake ^g | Varma et al., 2014 Gamenisch and Umehara, 2012 Kunze et al., 2015 Maeda et al., 2011 |
| | 1194 ^b | 24.9 | 58.3 | 4.3 | 2.27 | 2.52 | 1.22 | >51% | uptake | | |
| | 405 ^c | 24.9 | 58.3 | 4.3 | 0.77 | 2.52 | 2.26 | >10% | uptake | | |
| | 198 ^a | 359 | 64.6 | 11.8 | 0.80 | 0.21 | 2.22 | -- | all | | |
| | 198 ^a | 57.7 | 64.6 | 11.8 | 0.80 | 1.32 | 2.22 | -- | all | | |
| Bosentan | 132 ^a | 28.9 | 19.5 | 5.8 | 0.36 | 0.87 | 2.95 | -- | all | uptake ^g | Varma et al., 2014 Jones et al., 2012 Yoshikado et al., 2017 |
| | 142 ^b | 28.9 | 19.5 | 5.8 | 0.38 | 0.87 | 2.90 | -- | all | | |
| | 1117 ^c | 28.9 | 19.5 | 5.8 | 3.02 | 0.87 | 0.99 | -- | all | | |
| | 35 ^a | 12.1 | -- | 39 ^e | 0.02 | 3.24 | 3.93 | -- | all | | |
| | 2035 ^d | 14 | -- | 5.0 ^e | 1.09 | 0.36 | 1.91 | -- | all | | |
| Repaglinide | 166 ^a | 63.6 | 128 | 0.3 | 0.19 | 2.01 | 3.35 | -- | all | uptake ^g all ^h | Varma et al., 2014 Jones et al., 2012 Yoshikado et al., 2017 |
| | 1983 ^b | 63.6 | 128 | 0.3 | 2.32 | 2.01 | 1.21 | >40% | uptake | | |
| | 1151 ^c | 63.6 | 128 | 0.3 | 1.35 | 2.01 | 1.71 | >15% | uptake | | |
| | 299 ^a | 223 | 125 | 0.0 | 0.22 | 0.56 | 3.27 | -- | all | | |
| | 3671 ^d | 352 | 125 | 0.0 | 2.73 | 0.35 | 1.07 | -- | all | | |

^a – *in vitro* quantified + IVIVE

^b – *in vitro* quantified + IVIVE + empirical scaling factor for active uptake transport (individual scaling factor)

^c – *in vitro* quantified + IVIVE + empirical scaling factor for active uptake transport (geometric mean scaling factor)

^d – fitted parameters from *in vivo* using PBPK model

^e – composite $CL_{met+bile}$



^f – f_u and Q_h values as noted in each reference were used for analysis – note that f_u may vary for the same drug across different references

^g – RDS_{uptake} was established *in vivo* for atorvastatin and bosentan, since there was no significant AUC change to victim drugs when co-administered with IV itraconazole (CYP3A inhibitor) which resulted in 33% and 73% CYP3A inhibition, respectively (Maeda et al., 2011; Yoshikado et al., 2017). Midazolam, a CYP3A

probe, was used to assess magnitude of CYP3A inhibition. RDS_{uptake} was established for repaglinide via whole-body PBPK modeling of complex transporter- and enzyme-mediated DDI's (Varma et al., 2013).
^h – RDS_{all} was established *in vivo* for repaglinide since even though there was no change to systemic AUC by IV itraconazole, because CYP2C8 is the major hepatic drug metabolizing enzyme (Yoshikado et al., 2017). In a different study, PO trimethoprim, a selective CYP2C8 inhibitor, increased repaglinide AUC by 1.8-fold (Kim et al., 2016).

Downloaded from dmd.aspetjournals.org at ASPET Journals on April 24, 2019

Figure 1



| | A | B | C | D | E | |
|---------------|-----------------------------|-----|-----|----|----|-----|
| Fixed Ratio | $CL_{met+bile}$ (L/min) | 1 | 4 | 10 | 40 | 100 |
| | CL_{ef}^s (L/min) | 0.1 | 0.4 | 1 | 4 | 10 |
| | $CL_{met+bile} / CL_{ef}^s$ | 10 | 10 | 10 | 10 | 10 |
| Varying Ratio | $CL_{met+bile}$ (L/min) | 0.2 | 0.4 | 1 | 4 | 10 |
| | CL_{ef}^s (L/min) | 1 | 1 | 1 | 1 | 1 |
| | $CL_{met+bile} / CL_{ef}^s$ | 0.2 | 0.4 | 1 | 4 | 10 |

Figure 2






Figure 3



| $CL_{\text{met+bile}}/CL_{\text{ef}}^s$ | $PI_{\text{met+bile}}$ (%) | | |
|---|----------------------------|---------|---------|
| | Low ER | Mid ER | High ER |
| 4 – fold | $>19\%$ | $>50\%$ | $>80\%$ |
| 8 – fold | $>60\%$ | $>75\%$ | $>87\%$ |
| 16 – fold | $>80\%$ | $>90\%$ | $>95\%$ |

Figure 4

How to predict DDI liabilities for dual enzyme/transporter substrates?

What factors determine the switch from RDS_{uptake} to RDS_{all} ?

Fig. 1A

The magnitude of $CL_{\text{met+bile}}$ inhibition that leads to a DDI for a drug with RDS_{uptake} depends on the starting $CL_{\text{met+bile}}$ value.

Fig. 1B

The RDS_{uptake} switch to RDS_{all} depends on the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio and not the absolute CL magnitude.

Fig. 1C

The $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio that results in an $AUCR = 1.25$ signifies when RDS_{uptake} will switch to RDS_{all} . This ratio is defined as the tipping point.

How to use the RDS framework to determine DDI liabilities?

Fig. 2

1. Identify the drug's RDS in the absence of DDI.
 - The $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio and $CL_{\text{in}}^{\text{s}}/Q_{\text{h}}$ in relation to the tipping point defines the RDS of the drug.

Fig. 3

2. Identify when RDS_{uptake} will switch to RDS_{all} in the presence of DDI.
 - Inhibition of $CL_{\text{met+bile}}$ that results in the $CL_{\text{met+bile}}/CL_{\text{ef}}^{\text{s}}$ ratio to be less than the drug's tipping point, is defined as $PI_{\text{met+bile}}$.
 - $PI_{\text{met+bile}}$ identifies when inhibition of $CL_{\text{met+bile}}$ starts to become a DDI liability for victim drugs that have RDS_{uptake} .

Figure 5





Figure 6

A)



B)




Figure 7

