

DMD #15438

**Hepatic metabolism of MK-0457, a potent Aurora kinase  
inhibitor: Interspecies comparison and role of human cytochrome  
P450 and flavin-containing monooxygenase**

J. E. BALLARD, T. PRUEKSARITANONT, C. TANG

Department of Drug Metabolism

Merck Research Laboratories

West Point, PA 19486, USA

DMD #15438

**Running Title:** *In vitro* metabolism of MK-0457 in hepatic preparations.

**Correspondence:** Jeanine E. Ballard, Department of Drug Metabolism, WP75A-203,

Merck Research Laboratories, West Point, PA 19486, USA

Tel (215) 652-1398, Fax (215) 652-2410,

Email: [jeanine\\_ballard@merck.com](mailto:jeanine_ballard@merck.com)

**Number of Text Pages:** 11

**Number of Tables:** 0

**Number of Figures:** 4

**Number of References:** 19

**Word Counts:**

**Abstract:** 163

**Introduction:** 281

**Abbreviations:** CYP, cytochrome P450; FMO, flavin-monooxygenase;  $K_m$ , apparent Michaelis constant;  $K_s$ , substrate inhibition constant;  $K_i$ , inhibition constant;  $V_{max}$ , maximum reaction velocity; LC-MS/MS, liquid chromatography coupled with tandem mass spectrometry.

DMD #15438

## ABSTRACT

MK-0457, an Aurora kinase inhibitor in development for the treatment of cancer, was evaluated for its *in vitro* metabolism in different species. This compound primarily underwent *N*-oxidation and *N*-demethylation in human, monkey, dog, and rat liver preparations. However, *N*-demethylation was less significant in dogs. The formation of minor metabolites varied with species, but all metabolites generated in human hepatocytes were observed in animals. Results of immunoinhibition, selective chemical inhibition, thermal inactivation and metabolism by recombinant CYPs and FMOs strongly suggest that CYP3A4 and FMO3 comparably contributed to MK-0457 *N*-oxidation in human liver microsomes, where the reaction conformed to Michaelis-Menten kinetics. These studies indicate a major role of CYP2C8 in the *N*-demethylation reaction, while CYP3A4 only made a minor contribution. However, significant substrate inhibition was observed with MK-0457 *N*-demethylation, at high substrate concentrations ( $>10\ \mu\text{M}$ ) in human liver microsomes relative to the anticipated therapeutic exposure. A multi-enzyme metabolic pathway such as this may mitigate the potential of drug interactions in clinical treatment with MK-0457.

## Introduction

MK-0457 is a potent and selective Aurora kinase inhibitor in clinical development for the treatment of solid tumors and hematopoietic cancers. Aurora kinases are involved in the regulation of chromosome segregation and cytokinesis during mitosis, and overexpression of these proteins is likely to contribute to tumorigenesis (Katayama et al., 2003; Jiang et al., 2006). Inhibition of Aurora kinases disrupts progression of the cancer cell cycle and blocks proliferation *in vitro*, while noncycling cells remain unaffected. Rodent *in vivo* studies have shown that MK-0457 inhibits tumor growth, resulting in regression (Harrington et al., 2004). MK-0457 is the first kinase inhibitor to exhibit clinical activity in patients with T315I BCR-ABL mutated chronic myeloid leukemia or acute lymphocytic leukemia (Giles et al., 2007).

To gain an understanding of the species differences and similarities in the disposition of MK-0457 for the purpose of predicting human disposition and determining appropriate species for toxicology studies, metabolism studies were conducted in rat, dog, monkey, and human liver preparations. For the purpose of predicting potential drug-drug interactions, the drug-metabolizing enzymes involved in the biotransformation of MK-0457 in human liver microsomes were identified. In the present studies, the role of cytochrome P450 (CYP) and flavin-containing monooxygenase (FMO), two major enzyme systems that catalyze oxidative metabolic reactions, have been investigated using a combination of *in vitro* reaction phenotyping strategies. These methods include the use of (1) monoclonal antibodies for immuno-inhibition of specific CYP families, (2) selective chemical inhibitors, (3) cDNA-expressed CYP and FMO isoforms, and (4) thermal inactivation of FMO enzymes. Our results indicate that metabolism of MK-0457 in human liver microsomes was mediated by multiple drug-metabolizing enzymes and the resulting metabolite profile was similar to those in monkeys and rats.

DMD #15438

## Materials and Methods

*Chemicals and Biologics.* MK-0457 was synthesized at Merck Research Laboratories (MRL, Banyu, Japan). Its desmethyl- and *N*-oxide metabolites (VRT-171335 and VRT-750074) were provided by Vertex Pharmaceuticals Incorporated (Cambridge, MA), and [ $^{14}\text{C}$ ]MK-0457 (52 mCi/mmol) was prepared by ViTrax (Placentia, CA). Quercetin, sulfaphenazole, quinidine, ketoconazole, bupropion, NADPH,  $\text{MgCl}_2$ , and phosphate buffer were purchased from Sigma-Aldrich (St. Louis, MO). Preparations of pooled human liver microsomes (from 10 male donors) were purchased from XenoTech LLC (Lenexa, KS). Immuno-inhibitory monoclonal antibodies selective for CYP isoforms (anti-CYP1A2, anti-CYP2C, anti-CYP2D6, and anti-CYP3A) and microsomes prepared from baculovirus-infected insect cells containing cDNA-expressed CYP enzymes (CYP1A2, 2A6, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, 3A4, and 3A5) were provided in-house (MRL, West Point, PA). Microsomes prepared from baculovirus-infected insect cells containing cDNA-expressed FMOs 1, 3, and 5 were purchased from BD Gentest (Woburn, MA). Viable hepatocytes were isolated from fresh beagle dog, cynomolgus monkey and sprague-dawley rat tissue using a modification of the classical two-step collagenase perfusion technique (Li et al., 1992). Fresh human hepatocytes were purchased from In Vitro Technologies (Baltimore, MD), and two batches of cryopreserved cynomolgus monkey hepatocytes were purchased from CellzDirect (Pittsboro, NC).

*Incubations with hepatocytes.* Incubations of hepatocytes (1 million cells/mL) with [ $^{14}\text{C}$ ]MK-0457 (1 or 10  $\mu\text{M}$ ) were performed with continuous shaking (80 times/min) and a gentle flush of gases ( $\text{O}_2/\text{CO}_2$  95%/5%) at 37°C. Reactions were terminated with one half volume of acetonitrile and centrifuged to remove cell pellets prior to radiochromatography.

DMD #15438

*Incubations with human liver microsomes, recombinant human CYPs and FMOs .*

Incubations of human liver microsomes (1 mg protein/mL) with [<sup>14</sup>C]MK-0457 (1 or 10 μM) were carried out in phosphate buffer (0.1 M, pH 7.4) containing 1 mM MgCl<sub>2</sub>. The reactions were initiated by the addition of 1 mM NADPH and terminated with acetonitrile following 30-min or 60-min incubation at 37°C. Samples were centrifuged to remove precipitated protein and the resulting supernatant was subject to radiochromatography. For other *in vitro* studies with MK-0457 (1 or 10 μM, and 0.5 to 50 μM for metabolite formation kinetics), the incubation mixtures (0.25 ml) contained the same components except for respective enzyme preparations (0.1 mg/mL human liver microsomal protein, equivalent to 50.5 pmol/mL cytochrome P450; 20 pmol/ml recombinant human CYP; 0.2 mg/mL FMO). For thermal inactivation studies, human liver microsomes were preincubated at 50°C for 2 min in the absence of NADPH to inactivate FMO enzymes, and control reactions were preincubated at 37°C for 2 min. Following termination of the reaction, an aliquot of 0.5 μM bupropion was added as an internal standard. Samples were centrifuged, and the resulting supernatant was analyzed via LC-MS/MS. Assays were performed in triplicate under linear conditions with respect to protein concentration and incubation time.

*Immuno-inhibition.* Human liver microsomes (0.025 mg) were preincubated (at room temperature for 5 min) with anti-CYP antibody or non-specific IgG preparation (5 μL). The ratio of microsomal protein-to-antibody volume was chosen to give a maximal inhibitory effect. Afterwards, the samples were diluted with the buffer containing MgCl<sub>2</sub> and MK-0457 and reactions were initiated by the addition of 1 mM NADPH. Assays was performed in triplicate.

*Chemical inhibition.* Known chemical inhibitors selective for CYP2C8 (quercetin), CYP2C9 (sulfaphenazole), CYP2D6 (quinidine), and CYP3A (ketoconazole) were used to perform

DMD #15438

inhibition experiments. The concentration range of quercetin (1.2, 4, and 12  $\mu\text{M}$ ), sulfaphenazole (0.15, 0.5, and 1.5  $\mu\text{M}$ ), quinidine (0.03, 0.1, and 0.3  $\mu\text{M}$ ), and ketoconazole (0.015, 0.05, and 0.15  $\mu\text{M}$ ) was chosen based on the  $K_i$  values of previous reports (Bourrie et al., 1996; Rahman et al., 1994). Stock solutions of chemical inhibitors were prepared in 50% acetonitrile/water (v/v). The final concentration of acetonitrile in the incubation mixture was 1.5% (v/v). Control incubations, without chemical inhibitor, contained the same volume of acetonitrile, and did not exhibit enzyme inhibition compared to control incubations containing < 1% acetonitrile. Chemical inhibition experiments were performed in triplicate at MK-0457 concentrations of 1 and 10  $\mu\text{M}$  and were determined using human liver microsomes.

*Qualitative MS-Radiochromatography.* [ $^{14}\text{C}$ ]MK-0457 and its metabolites were separated on a Synergi Hydro-RP column (4.6 x 150 mm, 4  $\mu\text{m}$ , Phenomenex, Torrance, CA) using a Rheos 4000 binary pump (LEAP Technologies, Carrboro, NC) with a flow rate of 1 mL/min. The aqueous mobile phase (solvent A) consisted of 10% acetonitrile in 0.02% acetic acid (pH adjusted to 9.0 using triethylamine), while the organic phase (solvent B) consisted of acetonitrile. The initial mobile phase was composed of 5% of solvent B, which was maintained for 3 min and increased to 15% over a period of 0.5 min, and then further increased to 25% in 11.5 min, to 40% in 10 min, and finally to 70% over 0.5 min. This value was held for 2 min before returning to 5% of solvent B for column equilibration for 2 min prior to the next injection. The post-column flow rate was split 1:9, with 100  $\mu\text{L}/\text{min}$  directed to the LCQ ion trap and 900  $\mu\text{L}/\text{min}$  to the radiochemical director. Post-column radiochemical detection was performed using a Packard C515 flow scintillation analyzer (PerkinElmer Life Sciences, Meriden, CT) with a 500  $\mu\text{L}$  liquid flow cell and Packard scintillation cocktail (Ultima-Flo<sup>TM</sup>M) running at 3 mL/min. Mass spectrometry data were collected on the Finnigan LCQ ion trap system (Finnigan, San Jose, CA)

DMD #15438

equipped with an electrospray ionization (ESI) source operating in positive mode under optimized conditions (voltage, 4.5 kV; sheath gas, 70; auxillary gas, 10; capillary temperature, 220°C).

*Quantitative LC-MS/MS.* Quantitative analysis of two major metabolites of MK-0457 (desmethyl metabolite VRT-171335 and *N*-oxide VRT-750074) was accomplished by LC-MS/MS analysis. Separation was achieved on a Synergi Polar-RP column (2.0 x 50 mm, 4  $\mu$ m, Phenomenex, Torrance, CA) using PE 200 binary pumps. Solvent A consisted of 0.02% aqueous acetic acid and 10% acetonitrile, adjusted to pH 9.0 with NH<sub>4</sub>OH, and solvent B of 90% acetonitrile in water. The mobile phase was delivered at a flow rate of 0.5 mL/min with a linear increase of solvent B from 5% to 95% over 0.6 min and a hold at 95% for 1.4 min, followed by equilibration for an additional 1.5 min. LC-MS/MS was performed on a Sciex Model API 3000 triple quadrupole mass spectrometer (Concord, Ontario, Canada) interfaced to the column eluent via a Sciex turbospray probe operated at 450°C. Operating conditions common to both analytes were IS voltage (5000), NEB gas (8), CUR gas (12), and CAD gas (4). Other parameters were optimized by infusion of a mixture of all analytes, using selected reaction monitoring (SRM) experiments in positive ionization mode to monitor transitions of 451>383 for VRT-171335 and 481>394 for VRT-750074. The calibration curve (0.001 – 5  $\mu$ M) was constructed using the ratio of the analyte peak area to the internal standard peak area. The limit of quantification for this assay was 0.005  $\mu$ M for both metabolites.

*Data analysis.* The apparent enzyme kinetic parameters were determined by fitting the reaction velocities versus substrate concentrations to equation (1) or equation (2) (KaleidaGraph 3.52, Synergy Software, Reading PA), which describe Michaelis-Menten kinetics alone or Michaelis-Menten kinetics coupled with uncompetitive substrate inhibition, respectively.



DMD #15438

$$v = \frac{V_{\max} \cdot S}{K_m + S} \quad (1)$$

$$v = \frac{V_{\max} \cdot S}{K_m + S(1 + S / K_s)} \quad (2)$$

Where  $v$ ,  $S$ ,  $V_{\max}$ ,  $K_m$  and  $K_s$  are the velocity of reaction, substrate concentration, the maximum reaction velocity, apparent Michaelis constant and substrate inhibition constant, respectively.

## Results and Conclusions

Figure 1 outlines the metabolites of MK-0457 generated in human, cynomolgus monkey, dog, and rat hepatocytes. All species gave rise to two major metabolites (Figure 2), namely, *N*-oxide (VRT-750074) and desmethyl products (VRT-171335), although the desmethyl metabolite was much less significant in dogs. Consistent with the findings in human hepatocytes, MK-0457 was metabolized to its *N*-oxide and desmethyl metabolites in a preparation of pooled male human liver microsomes (Figure 3). Other minor metabolites were also observed in hepatocytes, including M1 (human, cynomolgus and rat), M2 (all species tested), and a glucuronide conjugate M3 (human, cynomolgus and rat). Structural confirmation of major metabolites (VRT-750074 and VRT-171335) was achieved using synthesized authentic standards, and identification of M3 as a glucuronide conjugate was based on mass change (data not shown). All of the metabolites detected in human hepatocytes were observed in rat and monkey hepatocytes as well. Also, VRT-750074 and VRT-171335 were the major metabolites formed *in vivo* in rats and monkeys following administration of MK-0457 (data not shown). However, in dog hepatocytes, *N*-desmethyl formation was low and M1 as well as the glucuronide conjugate M3 were not observed. The *in vitro* species differences observed in the metabolite profile of MK-0457 suggest that monkey and rat may potentially be better models than dog for the assessment of toxicological risks associated with MK-0457 metabolites.

Since *N*-demethylation and *N*-oxidation were the primary metabolic pathways of MK-0457 in human liver preparations, metabolizing enzymes involved in these reactions were characterized. Monoclonal antibodies, thermal inactivation, chemical inhibitors, and recombinant enzymes were employed to estimate the relative contribution of CYP and FMO

## DMD #15438

isoforms involved in MK-0457 metabolism in human liver microsomes. The findings are described below.

Immuno-inhibition was first approached to provide a quick assessment of relative importance of metabolizing enzymes. It was found that anti-CYP2C antibodies markedly inhibited the formation of *N*-desmethyl metabolite VRT-171335 ( $85 \pm 9$  and  $70 \pm 8\%$  inhibition at 1 and 10  $\mu\text{M}$ , respectively), and inclusion of anti-CYP3A antibody led to further inhibition ( $92 \pm 7\%$  inhibition at both MK-0457 concentrations). The result suggests that the reaction was primarily mediated by members of the CYP2C subfamily with a minor contribution from CYP3A enzymes. Further studies with recombinant CYPs showed that CYP2C8 was capable of catalyzing the reaction at the highest rate ( $1.69 \pm 0.18$  and  $13.63 \pm 0.93$  pmol/min/pmol CYP at 1 and 10  $\mu\text{M}$  MK-0457), followed by CYP3A4 ( $0.41 \pm 0.04$  and  $1.97 \pm 0.18$  pmol/min/pmol CYP at 1 and 10  $\mu\text{M}$  MK-0457). Moreover, of all the inhibitors tested, only quercetin significantly inhibited VRT-171335 formation ( $\sim 80\%$  inhibition at 12  $\mu\text{M}$ ). Although recombinant CYP2D6 was able to catalyze this reaction efficiently ( $0.22 \pm 0.02$  and  $1.83 \pm 0.14$  pmol/min/pmol CYP) compared with other isoforms ( $< 0.025$  and  $0.25$  pmol/min/pmol for CYP1A2, 2A6, 2B6, 2C9, 2C19, 2E1, and 3A5 at 1 and 10  $\mu\text{M}$  MK-0457), the contribution of this isoform *in vivo* is anticipated to be low based on the low abundance of CYP2D6 ( $\sim 2\%$  of total CYP content) compared to CYP2C8 ( $\sim 8\%$ ) and 3A4 ( $\sim 20\text{--}30\%$ ) (Lasker et al., 1998; Shimada, 1994; Rodrigues, 1999; Shimada et al., 1999). In agreement, anti-CYP2D6 antibodies and quinidine failed to inhibit the reaction. Collectively, *N*-demethylation of MK-0457 in human liver microsomes is primarily catalyzed by CYP2C8, with a minor contribution from CYP3A4.

On the other hand, the formation of *N*-oxide VRT-750074 was moderately inhibited by antibodies selective for CYP3A ( $25 \pm 6$  and  $48 \pm 7\%$  inhibition at 1 and 10  $\mu\text{M}$  MK-0457,

## DMD #15438

respectively) and by a combination of antibodies against CYPs 2C and 3A ( $45 \pm 12$  and  $60 \pm 8\%$  inhibition at 1 and 10  $\mu\text{M}$  MK-0457, respectively). The results suggest that the reaction was only partially catalyzed by the CYP3A and, to a minor extent, CYP2C subfamilies. Consistently, recombinant CYP3A4 and CYP2C8 displayed the highest and the 2<sup>nd</sup> highest metabolic rates ( $0.20 \pm 0.02$  and  $1.17 \pm 0.18$  pmol/min/pmol CYP for CYP3A4,  $0.036 \pm 0.003$  and  $0.33 \pm 0.05$  pmol/min/pmol CYP for CYP2C8 at 1 and 10  $\mu\text{M}$  MK-0457) among the isoforms tested ( $< 0.06$  pmol/min/pmol CYP for the rest of isoforms at 10  $\mu\text{M}$  MK-0457). In addition, moderate inhibition ( $\sim 50\%$ ) was achieved by ketoconazole (0.15  $\mu\text{M}$ ) and quercetin (12  $\mu\text{M}$ ). The partial inhibition of MK-0457 *N*-oxidation observed in the immuno-inhibition study elicited further investigation into the enzyme(s) responsible outside the cytochrome P450 family. A role for flavin-containing monooxygenases was anticipated based on their known ability to catalyze *N*-oxidation reactions (Cashman, 1995). This hypothesis was first investigated using the heat lability of FMOs compared to CYPs in human liver microsomes (Grothusen et al., 1996). It was found that pre-incubation of human liver microsomes at 50°C for 2 min (to inactivate FMOs) inhibited VRT-750074 formation by  $44 \pm 5\%$  (1  $\mu\text{M}$  MK-0457) and  $66 \pm 4\%$  (10  $\mu\text{M}$  MK-0457), while VRT-171335 formation was not significantly altered by the same treatment. Next, the specific isoform involved was elucidated using recombinant FMO enzymes. Among the three FMO isoforms tested, FMO1 and FMO3 displayed good catalytic activity for the formation of the *N*-oxide VRT-750074 ( $1832 \pm 7$  and  $1919 \pm 83$  pmol/min/mg protein for FMO1 and FMO3 at 10  $\mu\text{M}$  MK-0457), but FMO5 showed a  $\sim 10$ -fold lower activity ( $110 \pm 15$  pmol/min/mg protein at 10  $\mu\text{M}$  MK-0457). While FMO1 and FMO3 were both capable of efficiently catalyzing the *N*-oxidation reaction, the absence of FMO1 in adult human livers excludes its role in MK-0457 *N*-oxidation. Supported by another fact that the hepatic abundance of FMO3 is

DMD #15438

similar to CYP3A4 in humans (Cashman, 2002; Haining et al., 1997), it is conceivable that FMO3 appreciably contributes to MK-0457 *N*-oxidation in human liver microsomes.

Another interesting difference between these two major metabolic pathways in human liver microsomes is that *N*-oxidation conformed to typical Michaelis-Menten formation kinetics ( $K_m$  and  $V_{max}$  values of 6.6  $\mu\text{M}$  and 499 pmol/min/mg), while *N*-demethylation kinetics appeared to comprise of substrate inhibition as indicated by the decreased formation rate as substrate concentration exceeded  $\sim 10 \mu\text{M}$  (Figure 4). This observation was reproduced with recombinant human CYP2C8 (Figure 4). The reduced rate of *N*-demethylation cannot be the result of substrate precipitation at higher concentrations since it did not occur with recombinant CYP3A4 under the same conditions. The possibility of generation of an inhibitory metabolite was ruled out as well, because the product VRT-171335 is not a potent CYP2C8 inhibitor ( $\text{IC}_{50} > 20 \mu\text{M}$ ). However, different from other cases of substrate inhibition (Lin et al., 2001; Shou et al., 2000) where substrate inhibition constants ( $K_s$ ) are greater than the  $K_m$ , fitting the current data to equation 2 containing a substrate inhibition component gave rise to  $K_s$  values of 5.6 and 5.0  $\mu\text{M}$  for human liver microsomes and recombinant CYP2C8, respectively. These values are significantly lower than the respective  $K_m$  (24 and 64  $\mu\text{M}$ , respectively) for MK-0457 *N*-demethylation. Attempts to fit the data with a more complex model containing  $\alpha$  and  $\beta$  terms (Shou et al., 2000) led to a similar result. Apparently, better understanding of this unusual kinetic feature of CYP2C8-mediated reactions would be an interesting subject for future investigation. However, this phenomenon is not expected to have any clinical significance for MK-0457 as the inhibition occurs at substrate concentrations well above the anticipated therapeutic exposure ( $\leq 1 \mu\text{M}$ ).

## DMD #15438

In summary, MK-0457 metabolism in human hepatic preparations is characterized by two major pathways, namely, *N*-demethylation and *N*-oxidation, which are well mirrored in rat and cynomolgus monkey *in vitro* systems. The former pathway in humans is primarily catalyzed by CYP2C8 with a minor contribution of CYP3A4, while CYP3A4 and FMO3 appear to be equally important to the latter reaction. This multiple enzyme mediated metabolism may attenuate the potential of drug interactions in clinical treatment with MK-0457, regardless of the known inter-individual variability in enzyme abundance and activity (Shimada, 1994; Shimada et al., 1999; Bosch et al., 2006; Cashman, 2002) of these three enzymes as well as CYP3A4 vulnerability to induction and inhibition (Guengerich, 1999; Lin and Lu, 2001).

DMD #15438

### **Acknowledgements**

The authors thank Vertex Pharmaceuticals Incorporated for providing the metabolites VRT-750074 and VRT-171335, and ViTrax for preparing the radiolabeled MK-0457 used in these studies.

DMD #15438

## References

- Bosch TM, Meijerman I, Beijnen JH, Schellens JHM (2006) Genetic polymorphisms of drug-metabolising enzymes and drug transporters in the chemotherapeutic treatment of cancer. *Clin Pharmacokinet* **45**:253-285.
- Bourrie M, Meunier V, Berger Y, Fabre G (1996) Cytochrome P450 isoform inhibitors as a tool for the investigation of metabolic reactions catalyzed by human liver microsomes. *J Pharmacol Exp Ther* **277**:321-332.
- Cashman JR (1995) Structural and catalytic properties of the mammalian flavin-containing monooxygenase. *Chem Res Toxicol* **8**:165-181.
- Cashman JR (2002) Human flavin-containing monooxygenase (form 3): Polymorphisms and variations in chemical metabolism. *Pharmacogenomics* **3**:325-339.
- Giles FJ, Cortes J, Jones D, Bergstrom D, Kantarjian H, Freedman J (2007) MK-0457, a novel kinase inhibitor, is active in patients with chronic myeloid leukemia or acute lymphocytic leukemia with the T315I BCR-ABL mutation. *Blood* **109**:500-502.
- Grothusen A, Hardt J, Bräutigam L, Lang D, Böcker R (1996) A convenient method to discriminate between cytochrome P450 enzymes and flavin-containing monooxygenases in human liver microsomes. *Arch Toxicol* **71**:64-71.
- Haining RL, Hunter AP, Sadeque AJM, Philpot RM, Rettie AE (1997) Baculovirus-mediated expression and purification of human FMO3: Catalytic, immunochemical, and structural characterization. *Drug Metab Dispos* **25**:790-797.
- Harrington EA, Bebbington D, Moore J, Rasmussen RK, Ajoye-Adeogun AO, Nakayama T, Graham JA, Demur C, Hercend T, Diu-Hercend A, Su M, Golec JMC, Miller KM (2004)



DMD #15438

VX-680, a potent and selective small-molecule inhibitor of the Aurora kinases, suppresses tumor growth *in vivo*. *Nat Med* **10**:262-267.

Jiang N, Wang X, Yang Y, Dai W (2006) Advances in mitotic inhibitors for cancer treatment. *Mini Rev Med Chem* **6**:885-895.

Katayama H, Brinkley WR, Sen S (2003) The Aurora kinases: Role in cell transformation and tumorigenesis. *Cancer Metastasis Rev* **22**:451-464.

Lasker JM, Wester MR, Aramsombatdee E, Raucy JL (1998) Characterization of CYP2C19 and CYP2C9 from human liver: Respective roles in microsomal tolbutamide, s-mephenytoin, and omeprazole hydroxylations. *Arch Biochem Biophys* **353**:16-28.

Li AP, Rogue MA, Beck DJ, Kaminski DL (1992) Isolation and culturing of hepatocytes from human liver. *J Tissue Cult Method* 139-146.

Lin JH, Lu, AYH (2001) Interindividual variability in inhibition and induction of cytochrome P450 enzymes. *Annu Rev Pharmacol Toxicol* **41**:535-567.

Lin Y, Lu P, Tang C, Mei Q, Sandig G, Rodrigues AD, Rushmore TH, Shou M (2001) Substrate inhibition kinetics for cytochrome P450-catalyzed reactions. *Drug Metab Dispos* **29**:368-374.

Rahman A, Korzekwa KR, Grogan J, Gonzalez FJ, Harris JW (1994) Selective biotransformation of taxol to 6- $\alpha$ -hydroxytaxol by human cytochrome P450 2C8. *Cancer Res* **54**:5543-5546.

Rodrigues AD (1999) Integrated cytochrome P450 reaction phenotyping: Attempting to bridge the gap between cDNA-expressed cytochrome P450 and native human liver microsomes. *Biochem Pharmacol* **57**:465-480.

Shimada T (1994) Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: Studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* **270**:414-423.

DMD #15438

Shimada T, Tsumura F, Yamazaki H (1999) Prediction of human liver microsomal oxidations of 7-ethoxycoumarin and chlorzoxazone with kinetic parameters of recombinant cytochrome P-450 enzymes. *Drug Metab Dispos* **27**:1274-1280.

Shou M, Lin Y, Lu P, Tang C, Mei Q, Cui D, Tang W, Ngui JS, Lin CC, Singh R, Wong BK, Yergey JA, Lin JH, Pearson PG, Baillie TA, Rodrigues AD, Rushmore TH (2000) Enzyme kinetics of cytochrome P450-mediated reactions. *Curr Drug Metab* **2**:17-36.

DMD #15438

### Legends for Figures

Figure 1. Representative radiochromatograms of MK-0457 and metabolites extracted from dog (DHp; A), cynomolgus monkey (cMHP; B), human (HHp; C), and rat (RHp; D) hepatocyte incubates. Incubations were carried out with 10  $\mu\text{M}$  [ $^{14}\text{C}$ ]MK-0457 at 37°C for 30 minutes (DHp, cMHP, HHp) or 120 minutes (RHp) using 1 million cells/mL.

Figure 2. Structures of MK-0457 and major metabolites formed in suspensions of rat (RHp), dog (DHp), cynomolgus monkey (cMHP) and human hepatocytes (HHp) as well as human liver microsomes (HLM) supplemented with NADPH.

Figure 3. Representative radiochromatograms of MK-0457 and metabolites extracted from human liver microsome incubates. Incubations were carried out at 37°C with 1  $\mu\text{M}$  [ $^{14}\text{C}$ ]MK-0457 for 30 minutes (A) and 10  $\mu\text{M}$  [ $^{14}\text{C}$ ]MK-0457 for 60 minutes (B) using 1 mg/mL protein concentration.

Figure 4. Formation kinetics of *N*-desmethyl product VRT-171335 over MK-0457 concentration range of 0.5 - 50  $\mu\text{M}$  in human liver microsomes (solid circle) and recombinant CYP2C8 (open square). The kinetic parameters obtained by fitting the data to Equation 2 give rise to  $V_{\text{max}}$ ,  $K_m$  and  $K_s$  of 4950 pmol/min/mg protein, 24  $\mu\text{M}$  and 5.6  $\mu\text{M}$  for human liver microsomes. The corresponding values for recombinant CYP2C8 are 129 pmol/min/pmol CYP, 64  $\mu\text{M}$  and 5.0  $\mu\text{M}$ .





Figure 1




**CYP3A4  
FMO3**



**VRT-7500074**  
(DHp, HHp, cMHP, RHp, HLM)

**CYP2C8  
CYP3A4**



**VRT-171335**  
(DHp, HHp, cMHP, RHp, HLM)

**Figure 2**

CPM





Figure 3




Figure 4