

DMD # 70300

Short Communication

Transcriptional Regulation of SULT1C2 by Vitamin D Receptor in LS180 Human Colorectal Adenocarcinoma Cells

Kathleen G. Barrett, Hailin Fang, Thomas A. Kocarek, and Melissa Runge-Morris
Institute of Environmental Health Sciences, Wayne State University, Detroit, MI 48202
(K.G.B., H.F., T.A.K., M.R.-M.)

DMD # 70300

Running Title: Vitamin D regulation of SULT1C2

Address correspondence to: Dr. Melissa Runge-Morris, Institute of Environmental Health Sciences, 6135 Woodward Avenue, Integrative Biosciences Building, Room 2116, Wayne State University, Detroit, MI 48202, USA. Tel: (313) 577-5598; FAX: (313) 972-8025; E-mail: m.runge-morris@wayne.edu

Number of text pages: 18

Number of tables: 0 (1 Supplemental)

Number of figures: 4

Number of references: 20

Number of words in Abstract: 227

Number of words in Introduction: 283

Number of words in Results and Discussion: 673

Abbreviations: DR3, direct repeat of AGGTCA with 3 intervening nucleotides; GW3965, 3-[3-[N-(2-chloro-3-trifluoromethylbenzyl)-(2,2-diphenylethyl)amino]propyloxy]phenylacetic acid; GW4064, 3-(2,6-dichlorophenyl)-4-(3'-carboxy-2-chlorostilben-4-yl)oxymethyl-5-isopropylisoxazole; HEK, human embryonic kidney cells; Luc, firefly luciferase; PCR, polymerase chain reaction; PXR, pregnane X receptor; PXRE, PXR response element; RXR, retinoid X receptor; SULT, cytosolic sulfotransferase; VitD₃, 1 α ,25-dihydroxyvitamin D₃; VDR, vitamin D receptor; VDRE, vitamin D response element; WT, wild-type.

Abstract

The factors that regulate expression of genes in the 1C family of human cytosolic sulfotransferases (SULT1C) are not well understood. In a recent study evaluating the effects of a panel of transcription factor activators on SULT1C family member expression in LS180 human colorectal adenocarcinoma cells, we found that SULT1C2 expression was significantly increased by 1 α ,25-dihydroxyvitamin D₃ (VitD₃) treatment. The objective of the current study was to identify the mechanism responsible for VitD₃-mediated activation of SULT1C2 transcription. VitD₃ treatment of LS180 cells activated transcription of a transfected luciferase reporter plasmid that contained ~5Kb of the SULT1C2 gene, which included 402 nt of the non-coding exon 1, all of intron 1, and 21 nt of exon 2. Although computational analysis of the VitD₃-responsive region of the SULT1C2 gene identified a pregnane X receptor (PXR)-binding site within exon 1, the transfected 5Kb SULT1C2 reporter was not activated by treatment with rifampicin, a prototypical PXR agonist. However, deletion or mutation of the predicted PXR-binding site abolished VitD₃-mediated SULT1C2 transcriptional activation, identifying the site as a functional vitamin D response element (VDRE). We further demonstrated that vitamin D receptor (VDR) can interact directly with the SULT1C2 VDRE sequence using an ELISA-based transcription factor binding assay. In conclusion VitD₃-inducible SULT1C2 transcription is mediated through a VDRE in exon 1. These results suggest a role for SULT1C2 in VitD₃-regulated physiological processes in human intestine.

Introduction

The cytosolic sulfotransferases (SULTs) are a family of conjugating enzymes that catalyze the biotransformation of a wide variety of exogenous and endogenous substrates. The human SULT1C subfamily consists of three members, SULT1C2, SULT1C3, and SULT1C4. The characterization of SULT1C substrate specificity, expression, and regulation is incomplete, but available evidence suggests that these enzymes might play metabolic roles during development (Runge-Morris and Kocarek, 2013). Human SULT1C2 is expressed in multiple tissues including fetal liver, stomach, and the vitamin D target tissues thyroid, intestine and kidney (Runge-Morris and Kocarek, 2013). SULT1C2 does not metabolize prototypical SULT substrates but has been shown to sulfonate thyroid hormones and some phenols and to bioactivate the procarcinogen N-hydroxy-2-acetylaminofluorene (Sakakibara et al., 1998; Li et al., 2000; Allali-Hassani et al., 2007).

We recently reported that treatment of LS180 human colorectal adenocarcinoma cells with several nuclear receptor activators, including GW3965 (liver X receptor agonist), GW4064 (farnesoid X receptor agonist), rifampicin [pregnane X receptor (PXR) agonist], or 1 α ,25-dihydroxyvitamin D₃ (VitD₃) [vitamin D receptor (VDR) agonist], increased SULT1C2 expression (Rondini et al., 2014). The VitD₃-mediated induction was especially pronounced and occurred at the mRNA, protein, and promoter activation levels. It therefore seems likely that VitD₃-mediated regulation of SULT1C2 could contribute to VitD₃-regulated processes in intestine and possibly other VitD₃ target tissues, such as kidney.

DMD # 70300

VitD₃ regulates gene transcription by activating the ligand-dependent nuclear receptor, vitamin D receptor (VDR), which binds as a heterodimer with retinoid X receptor (RXR) to vitamin D response elements (VDREs) in target genes (Haussler et al., 1997). VDREs are classically direct repeats of AGGTCA separated by three nucleotides (i.e., DR3 motifs). The objective of this study was to identify the *cis*-acting element responsible for VitD₃-mediated regulation of SULT1C2 transcription.

Materials and Methods

Cell Culture. LS180 colorectal adenocarcinoma cells were obtained from the American Type Culture Collection (Manassas, VA) and were cultured as previously described (Rondini et al., 2014). HEK 293 cells (human embryonic kidney cells), used for protein expression, were obtained from Dr. Ye-Shih Ho (Wayne State University) and cultured in Dulbecco's Modified Eagle Medium supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin and 10% fetal bovine serum (all purchased from Life Technologies, Grand Island, NY). Cells were maintained under a humidified atmosphere of 95% air, 5% CO₂ at 37°C.

Preparation of Firefly Luciferase (Luc) Reporter Plasmids. Preparation of the SULT1C2 (-4998:-1)-Luc reporter plasmid has been described previously where it was referred to as SULT1C2#2 (Rondini et al., 2014). The SULT1C2 (-4827:-1)-Luc deletion construct was prepared using SULT1C2 (-4998:-1)-Luc as template, HotStar HiFidelity Polymerase (Qiagen, Valencia, CA), and the primer pairs listed in Supplemental Table 1. The PCR fragment was digested with KpnI and NheI and ligated into the corresponding sites of the pGL4.24 [luc2P/minP] reporter plasmid (Promega, Madison,

DMD # 70300

WI). Site-directed mutagenesis of a computationally predicted (MatInspector; Genomatix, Inc., Cincinnati, OH) (Quandt et al., 1995; Cartharius et al., 2005) PXR-binding site was performed using the QuikChange II Kit (Agilent Technologies, Santa Clara, CA) according to manufacturer's instructions, with SULT1C2 (-4998:-1)-Luc as template and the primers listed in Supplemental Table 1.

Transient Transfection Analysis. Approximately 200,000 LS180 cells were seeded into the wells of 12-well plates. Seventy-two hours later, cultures were transfected with 1.6 μ g of a SULT1C2 Luc reporter plasmid and 1 ng of the *Renilla* luciferase expression plasmid, pRL-CMV (Promega), as previously described (Rondini et al., 2014). After 18h, transfection medium was replaced and cells were treated with 0.1% ethanol (Sigma-Aldrich, St. Louis, MO) or 0.1 μ M VitD₃ (Sigma-Aldrich) for 48h. Cells were harvested for measurement of firefly and *Renilla* luciferase activities using the Dual Luciferase Assay (Promega) and a GloMax Luminometer (Promega).

VDR•VDRE *In Vitro* Binding Assay. A VDR expression plasmid was prepared by amplifying the VDR coding sequence (RefSeq NM_000376.2), using LS180 cDNA as template, Herculase II Fusion DNA Polymerase (Agilent Technologies), and the primers listed in Supplemental Table 1. The amplified fragment was digested with HindIII and XhoI and ligated into the pcDNA3.1 expression plasmid (Life Technologies). Approximately 1,000,000 HEK 293 cells were seeded into 100 mm plates. Once 70-80% confluent, the cells were transfected with 60 μ l Lipofectamine 2000 (Life Technologies), 1.5 μ g of VDR-pcDNA3.1, and 1.5 μ g RXR α -pSG5 (provided by Dr. Steven Kliewer, University of Texas Southwestern, Dallas, TX). Forty-eight hours later,

DMD # 70300

nuclear proteins were extracted using the NucBuster Protein Extraction kit (Novagen, EMD Millipore, Billerica, MA), and protein concentrations were quantified using the bicinchoninic acid protein assay (Thermo Fisher Scientific, Rockford, IL).

The Universal EZ-TFA Chemiluminescent Transcription Factor Assay (EMD Millipore) was performed according to the manufacturer's protocol. Briefly, 0.75 µg HEK 293 nuclear extract containing VDR and RXRα, 2 pmol biotinylated capture probe containing the VDRE consensus sequence from the rat osteocalcin gene [also known as bone gamma-carboxyglutamate, official symbol Bglap; Markose et al., 1990; Demay et al., 1992], and transcription factor assay buffer were added to the wells of a streptavidin-coated microplate. Competition for binding of VDR to the capture probe was assessed by adding one of the following unlabeled double-stranded oligonucleotides to the binding reaction: (1) consensus VDRE; (2) mutated consensus VDRE (Gutierrez et al., 2004); (3) SULT1C2 VDRE; (4) mutated SULT1C2 VDRE. The competitors were added at 1.5- to 50-fold molar excess of the biotinylated capture probe. Background binding was determined by assessing VDR binding to a biotinylated mutated consensus VDRE probe. Capture probe, competitor probe, and negative control probe sequences are listed in Supplemental Table 1. Capture probe-bound VDR was detected using the primary antibody, VDR (D-6) X (sc-13133X; Santa Cruz Biotechnology, CA), diluted 1:5000, followed by horseradish peroxidase-conjugated secondary antibody (1:500 dilution), and chemiluminescence was measured using kit reagents and a GloMax Luminometer. Data are expressed relative to VDR•RXR binding to the consensus VDRE capture probe in the absence of competitor.

DMD # 70300

Statistical Analysis. Transfection and transcription factor binding assay data were analyzed using one-way analysis of variance (ANOVA) followed by the Newman-Keuls or Dunnett's multiple comparison test. $p < 0.05$ was considered statistically significant. Binding curves with 95% confidence intervals were generated using the sigmoidal dose-response algorithm of Prism 6 for Windows (GraphPad Software Inc., La Jolla, CA).

Results and Discussion

We previously reported that VitD₃ treatment of LS180 cells increased activity from a transfected reporter plasmid containing ~5 Kb of the SULT1C2 gene (-4998:-1 relative to the translation start site in exon 2, shown schematically in Fig. 1) (Rondini et al., 2014). Recently, in a genome-wide ChIP-seq analysis of VitD₃-treated LS180 cells, Meyer et al. (2012) detected a VDR•RXR binding peak at nt 108,288,453 to 108,289,105 of chromosome 2 [Gene Expression Omnibus GSE31939; RefSeq NC_000002.12], with the peak center located within the SULT1C2 non-coding exon 1. This information suggested that a VDRE site is located near the 5'-end of our SULT1C2 (-4998:-1) fragment. We therefore deleted 171 nt from the 5'-end of SULT1C2 (-4998:-1), creating the SULT1C2 (-4827:-1)-Luc reporter. Figure 2 shows that this deletion abolished VitD₃-mediated SULT1C2 activation, confirming the presence of a VitD₃-responsive site in this region.

MatInspector software was used to identify putative transcription factor-binding sites within the deleted 171 nt sequence. Instead of detecting a prototypical VDR•RXR DR3 motif in exon 1, a motif identified as a putative PXR•RXR binding site was detected

DMD # 70300

at nt -4887 to -4863 (predicted PXRE, Figure 1). However, we previously reported that treatment of LS180 cells with the prototypical PXR agonist, rifampicin, did not increase expression from the SULT1C2 (-4998:-1)-Luc reporter (Rondini et al., 2014), suggesting that the computationally-predicted sequence is not a functional PXR response element (PXRE), but rather potentially a VDRE site. Mutation of the core sequence of the predicted PXR-binding site (from GGT to AAC) within the SULT1C2 (-4998:-1)-Luc plasmid caused a 94% reduction in VitD₃-mediated SULT1C2 reporter activation compared to the wild-type construct (Figure 3), further supporting the conclusion that this site is a functional VDRE.

An ELISA-based transcription factor-binding assay was used to determine whether VDR•RXR can bind directly to the VDRE site in exon 1 of the human SULT1C2 gene. The capture probe containing the VDRE consensus sequence from the rat osteocalcin gene promoter was incubated with unlabeled competitor probes added in 50-fold molar excess and nuclear protein extract from HEK 293 cells expressing VDR and RXR α . The SULT1C2 VDRE competitor significantly inhibited VDR•RXR binding to the capture probe by 74%, which was similar to the inhibition (79%) that was observed when consensus VDRE competitor was added (Figure 4). Mutated consensus and SULT1C2 VDRE competitors did not significantly inhibit binding to the capture probe (25% and 31% inhibition, respectively). Incubation with increasing amounts (1.5- to 50-fold molar excess) of SULT1C2 VDRE competitor inhibited VDR•RXR binding to the capture probe with an IC₅₀ that was comparable to that of the consensus VDRE competitor (Figure 4 Inset).

DMD # 70300

VDR and PXR are both members of the NR1I nuclear receptor group, and therefore have high sequence similarity (Kliwer et al., 1998). Additionally, they have been shown to recognize similar ligands and regulate similar target genes. For example, VDR and PXR are both activated by the secondary bile acid lithocholic acid (Staudinger et al., 2001; Makishima et al., 2002) and both receptors regulate expression of CYP3A4 (Bertilsson et al., 1998; Schmiedlin-Ren et al., 2001; Thummel et al., 2001). In the case of CYP3A4, PXR and VDR activate transcription through the same *cis*-acting elements, a proximal everted repeat with six intervening nucleotides motif and a distal DR3 motif (Goodwin et al., 1999; Thummel et al., 2001; Thompson et al., 2002). By comparison, while both PXR and VDR activate SULT1C2 expression in LS180 cells (Rondini et al., 2014), in this case the two receptors do not appear to utilize the same response element. While VitD₃ treatment activates SULT1C2 transcription through the element at nt -4887 to -4863, rifampicin treatment did not increase transcription from the (-4998:-1) reporter construct containing that element (Rondini et al., 2014). Therefore, VDR/PXR regulation of CYP3A4 differs somewhat from SULT1C2 regulation by these two receptors. Presumably the restriction by the SULT1C2 motif to VDR is physiologically important for ensuring appropriate SULT1C2 expression in response to hormonal versus xenobiotic signals. Further research is needed to determine the role that this mode of regulation plays in intestinal physiology.

DMD # 70300

Authorship Contributions

Participated in research design: Barrett, Kocarek, and Runge-Morris

Conducted experiments: Barrett and Fang

Performed data analysis: Barrett and Kocarek

Wrote or contributed to the writing of the manuscript: Barrett, Kocarek, and Runge-Morris

DMD # 70300

References

Allali-Hassani A, Pan PW, Dombrovski L, Najmanovich R, Tempel W, Dong A, Loppnau P, Martin F, Thornton J, Edwards AM, Bochkarev A, Plotnikov AN, Vedadi M and Arrowsmith CH. (2007) Structural and chemical profiling of the human cytosolic sulfotransferases. *PLoS.Biol.* 5(5), e97.

Bertilsson G, Heidrich J, Svensson K, Asman M, Jendeberg L, Sydow-Backman M, Ohlsson R, Postlind H, Blomquist P and Berkenstam A. (1998) Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc.Natl.Acad.Sci.U.S.A.* 95(21), 12208-12213.

Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, Frisch M, Bayerlein M, and Werner T (2005) MatInspector and beyond: promoter analysis based on transcription factor binding sites. *Bioinformatics.* 21 (13):2933-2942.

Demay MB, Kiernan MS, DeLuca HF and Kronenberg, HM. (1992) Characterization of 1,25-dihydroxyvitamin D3 receptor interactions with target sequences in the rat osteocalcin gene. *Mol.Endocrinol.* 6(4), 557-562.

Goodwin B, Hodgson E and Liddle C. (1999) The orphan human pregnane X receptor mediates the transcriptional activation of CYP3A4 by rifampicin through a distal enhancer module. *Mol.Pharmacol.* 56(6), 1329-1339.

DMD # 70300

Gutierrez S, Liu J, Javed A, Montecino M, Stein GS, Lian JB and Stein JL. (2004) The vitamin D response element in the distal osteocalcin promoter contributes to chromatin organization of the proximal regulatory domain. *J.Biol.Chem.* 279(42), 43581-43588.

Haussler MR, Haussler CA, Jurutka PW, Thompson PD, Hsieh JC, Remus LS, Selznick SH and Whitfield GK. (1997) The vitamin D hormone and its nuclear receptor: molecular actions and disease states. *J Endocrinol.* 154:S57-S73.

Kliwer SA, Moore JT, Wade L, Staudinger JL, Watson MA, Jones SA, McKee DD, Oliver BB, Willson TM, Zetterstrom RH, Perlmann T and Lehmann JM. (1998) An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell.* 92(1), 73-82.

Li X, Clemens DL and Anderson RJ. (2000) Sulfation of iodothyronines by human sulfotransferase 1C1 (SULT1C1)*. *Biochem.Pharmacol.* 60(11), 1713-1716.

Makishima M, Lu TT, Xie W, Whitfield GK, Domoto H, Evans RM, Haussler MR and Mangelsdorf DJ. (2002) Vitamin D receptor as an intestinal bile acid sensor. *Science.* 296(5571), 1313-1316.

DMD # 70300

Markose ER, Stein JL, Stein GS and Lian JB. (1990) Vitamin D-mediated modifications in protein-DNA interactions at two promoter elements of the osteocalcin gene.

Proc.Natl.Acad.Sci.USA. 87(5), 1701-1705.

Meyer MB, Goetsch PD, and Pike JW. (2012) VDR/RXR and TCF4/ β -Catenin Cistromes in Colonic Cells of Colorectal Tumor Origin: Impact on c-FOS and c-MYC Gene Expression. *Mol Endocrinol.* 26(1):37-51.

Quandt K, Frech K, Karas H, Wingender E, and Werner T (1995) MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res.* 23 (23):4878-4884.

Rondini EA, Fang H, Runge-Morris M, and Kocarek TA. (2014) Regulation of cytosolic sulfotransferases 1C2 and 1C3 by nuclear signaling pathways in LS180 colorectal adenocarcinoma cells. *Drug Metab Dispos.* 42:361-368.

Runge-Morris M and Kocarek TA. (2013) Expression of the sulfotransferase 1C family: Implications for xenobiotic toxicity. *Drug Metab Rev.* 45(4):450-459.

DMD # 70300

Sakakibara Y, Yanagisawa K, Katafuchi J, Ringer DP, Takami Y, Nakayama T, Suiko M and Liu MC. (1998) Molecular cloning, expression, and characterization of novel human SULT1C sulfotransferases that catalyze the sulfonation of N-hydroxy-2-acetylaminofluorene. *J.Biol.Chem.* 273(51), 33929-33935.

Schmiedlin-Ren P, Thummel KE, Fisher JM, Paine MF and Watkins PB. (2001) Induction of CYP3A4 by 1 alpha,25-dihydroxyvitamin D3 is human cell line-specific and is unlikely to involve pregnane X receptor. *Drug Metab Dispos.* 29(11), 1446-1453.

Staudinger JL, Goodwin B, Jones SA, Hawkins-Brown D, MacKenzie KI, LaTour A, Liu Y, Klaassen CD, Brown KK, Reinhard J, Willson TM, Koller BH and Kliewer SA. (2001) The nuclear receptor PXR is a lithocholic acid sensor that protects against liver toxicity. *Proc.Natl.Acad.Sci.U.S.A.* 98(6), 3369-3374.

Thompson, P. D., Jurutka, P. W., Whitfield, G. K., Myskowski, S. M., Eichhorst, K. R., Dominguez, C. E., Haussler, C. A., and Haussler, M. R. (2002) Liganded VDR induces CYP3A4 in small intestinal and colon cancer cells via DR3 and ER6 vitamin D responsive elements. *Biochem.Biophys.Res.Comm.* 20;299(5), 730-738.

Thummel KE, Brimer C, Yasuda K, Thottassery J, Senn T, Lin Y, Ishizuka H, Kharasch E, Schuetz J and Schuetz E. (2001) Transcriptional control of intestinal cytochrome P-4503A by 1alpha, 25-dihydroxy vitamin D3. *Mol.Pharmacol.* 60(6), 1399-1406.

DMD # 70300

Footnotes

This research was supported by the National Institutes of Health National Institute of Environmental Health Sciences [Grants R01 ES022606 (to M.R.-M.) and Center Grant P30 ES020957].

These results were presented in preliminary form at the Society of Toxicology 2015 Annual Meeting.

DMD # 70300

Figure Legends

Fig. 1. Schematic representation of the SULT1C2 (-4998:-1) fragment. A ~5Kb fragment of the SULT1C2 gene containing nt -4998 to -1 relative to the translation start site was amplified and ligated into a luciferase reporter plasmid. This fragment includes 402 nt of the non-coding exon 1, intron 1, and 21 nt of exon 2. Computational analysis identified a putative PXR-binding site (predicted PXRE, core sequence underlined) at nt -4887 to -4863. SULT1C2 (-4827:-1) shows the PXR-binding site deletion fragment.

Fig. 2. VitD₃ treatment activates reporter expression from SULT1C2 construct (-4998:-1) but not from deletion construct (-4828:-1) in LS180 cells. LS180 cells were transiently transfected with SULT1C2 (-4998:-1)-Luc, SULT1C2 (-4827:-1)-Luc, or empty reporter plasmid (pGL4.24) and treated with 0.1% ethanol or 0.1μM VitD₃ for 48h. Cells were harvested and luciferase activities were measured. Each bar represents the mean ± S.D. of normalized (firefly/Renilla) luciferase activity relative to the activity measured in ethanol-treated, SULT1C2 (-4998:-1)-Luc transfected cells (n=9 wells per group, derived from combining data from three independent experiments with triplicate transfection). ***Significantly different from ethanol-treated cells transfected with the same reporter plasmid, $P < 0.001$.

Fig. 3. Mutation of the predicted PXR-binding site in exon 1 attenuates VitD₃-mediated SULT1C2 (-4998:-1)-Luc reporter activation. LS180 cells were transiently transfected with SULT1C2 (-4998:-1)-Luc containing either wild-type (WT) or mutated (Mut) predicted PXR-binding site and treated with 0.1% ethanol or 0.1μM VitD₃. Forty-

DMD # 70300

eight hours later, the cells were harvested and luciferase activities measured. Each bar represents the mean \pm S.D. normalized (firefly/*Renilla*) luciferase activity relative to the activity measured in ethanol-treated WT SULT1C2 (-4998:-1)-Luc transfected cells (n=6 wells per group, derived from combining data from two independent experiments with triplicate transfection). ***Significantly different from ethanol-treated control, $P < 0.001$.

Fig. 4. VDR•RXR binds to the predicted PXR-binding site (VDRE) in exon 1 of SULT1C2. An ELISA-based *In vitro* transcription factor-binding assay was performed as described in Materials and Methods. Binding reactions contained nuclear protein extract and a biotinylated consensus or mutated VDRE capture probe, alone or in the presence of a 50-fold molar excess of either wild-type (WT) or mutated (Mut) consensus VDRE or WT or Mut SULT1C2 VDRE competitor probe. Each bar represents the mean \pm S.E.M. binding of VDR•RXR to a VDRE capture probe relative to the binding to the consensus VDRE capture probe that was detected in the absence of competitor (n=4, derived from means of four independent binding experiments, each performed with duplicate wells). ***Significantly different from consensus VDRE capture probe in absence of competitor, $P < 0.001$. Inset: Binding affinity was assessed by adding different amounts (1.5- to 50-fold molar excess) of competitor probes to the consensus VDRE capture probe and nuclear protein extract. IC₅₀ values with 95% confidence intervals (CI) are shown. Each data point is the mean from two independent experiments, each performed in duplicate wells.

Fig. 1




Fig. 2




Fig. 3





Fig. 4



Supplemental Data

Transcriptional Regulation of SULT1C2 by Vitamin D Receptor in LS180 Human Colorectal Adenocarcinoma Cells

Kathleen G. Barrett, Hailin Fang, Thomas A. Kocarek, and Melissa Runge-Morris. Institute of Environmental Health Sciences, Wayne State University, Detroit, MI 48202 (K.G.B., H.F., T.A.K., M.R.-M.)

Drug Metabolism and Disposition

Supplemental Table 1. PCR Primers and oligonucleotides used to prepare constructs and probes in this study.

	Sequence	Comments
Primers to prepare SULT1C2 (-4998:-1)		
FOR	5' GGG <u>GGTACC</u> CAGGCAGCTGAGGGCCAGGA 3'	KpnI site underscored.
REV	5' GGG <u>GCTAGC</u> AGTGTCTCAGGGTTGGGGTC 3'	NheI site underscored.
Primers to prepare SULT1C2 (-4827:-1) deletion construct		
FOR	5' GGG <u>GGTACC</u> ACAGGGTTTGTCTGTTCATCCG 3'	KpnI site underscored.
REV	5' GGG <u>GCTAGC</u> AGTGTCTCAGGGTTGGGGTC 3'	NheI site underscored; same REV primer for SULT1C2 (-4998:-1) (Rondini et al., 2014).
Primers to prepare mutant VDRE SULT1C2		
FOR	5' TTGGTATTTGCACAGCAGTCAGTTGCACATGAGTGATCATGGTACAG 3'	Mutated bases underscored.
REV	5' CTGTACCATGATCACTCATGTGCAACTGACTGCTGTGCAAATACCAA 3'	Mutated bases underscored.
Primers to prepare VDR expression plasmid		
FOR	5' GGG <u>AAGCTT</u> CCCCCTGCTCCTTCAGGGATG 3'	HindIII site underscored.
REV	5' GGG <u>CTCGAG</u> TCAGGAGATCTCATTGCCAAACACT 3'	XhoI site underscored.
Oligonucleotides for VDR-VDRE <i>In Vitro</i> Binding Assay		
Capture Probe: 5'-end biotin-modified consensus VDRE probe		
Sense	5' TGGATGAGCGGAGCTGCCCTGCACTGGGTGAATGAGGACATTACTGACCG 3'	VDRE consensus sequence (DR3) underscored. 5'-end biotin modification of sense strand.
Antisense	5' CGGTCAGTAATGTCCCTCATTCACCCAGTGCAGGGCAGCTCCGCTCATCCA 3'	
Negative Probe: 5'-end biotin-modified mutated consensus VDRE probe		
Sense	5' TGGATGAGCGGAGCTGCCCTGCACTGtaTGAATGActACATTACTGACCG 3'	Consensus VDRE sequence (DR3) underscored; mutated bases lower case. 5'-end biotin modification of sense strand.
Antisense	5' CGGTCAGTAATGTaGTCATTCAtaCAGTGCAGGGCAGCTCCGCTCATCCA 3'	
Competitor 1: Consensus VDRE probe		
Sense	5' GCCCTGCACTGGGTGAATGAGGACATTACTGACCG 3'	Consensus VDRE sequence (DR3) underscored.
Antisense	5' CGGTCAGTAATGTCCCTCATTCACCCAGTGCAGGGC 3'	
Competitor 2: Mutated consensus VDRE probe		
Sense	5' GCCCTGCACTGtaTGAATGActACATTACTGACCG 3'	Mutated consensus VDRE sequence (DR3) underscored; mutated bases lower case.
Antisense	5' CGGTCAGTAATGTaGTCATTCAtaCAGTGCAGGGC 3'	
Competitor 3: SULT1C2 VDRE probe		
Sense	5' ACAGCAGTCAGGGTCACATGAGTGATCATGGTACA 3'	Computationally predicted SULT1C2 PXRE core sequence underscored.
Antisense	5' TGTACCATGATCACTCATGTGACCCTGACTGCTGT 3'	
Competitor 4: SULT1C2 mutated VDRE probe		
Sense	5' ACAGCAGTCAGaacCACATGAGTGATCATGGTACA 3'	Computationally predicted PXRE core sequence underscored; mutated bases lower case.
Antisense	5' TGTACCATGATCACTCATGTGgttCTGACTGCTGT 3'	