

**DMD/2019/086595**

**Title:**

**Carfilzomib is not an appropriate payload of antibody-drug conjugates as rapid  
inactivation by lysosomal enzymes**

**Authors:**

Yong Ma, Josefa Dela Cruz-Chuh, S Cyrus Khojasteh, Peter S Dragovich, Thomas H Pillow,  
Donglu Zhang

Affiliation: Drug Metabolism and Disposition (YM, SCK, DZ), Biochemical and Cellular Pharmacology  
JDC), Discovery Chemistry (PD, TP), Genentech, 1 DNA Way, South San Francisco, CA 98080

Running Title: Carfilzomib as an inactive ADC payload

Corresponding author: Donglu Zhang or Yong Ma

Drug Metabolism and Pharmacokinetics, Genentech, 1 DNA Way, South

San Francisco, CA 94080. [zhang.donglu@gene.com](mailto:zhang.donglu@gene.com) or [yongma87@gmail.com](mailto:yongma87@gmail.com)

Number of Table:

Number of Figures: 7

Number of references: 20

Number of words in sections:

Abstract: 225

Introduction: 374

Results and discussion: 1445

Abbreviations: CFZ, carfilzomib; ADC, antibody drug conjugate; MMAE, monomethyl auristatin E; AE, adverse event; MC, maleimidocaproyl; ValCit, valine-citrulline; PABQ, *p*-aminobenzyl quaternary ammonium; DAR, drug-to-antibody ratio; LC-HRMS, liquid chromatography-high resolution mass spectrometry.

**Abstract:**

Carfilzomib (CFZ) is a proteasome inhibitor used for oncology indications including treating multiple myeloma. CFZ is a potent cytotoxic agent with an  $IC_{50}$  in the nanomolar range in various cancer cell lines and was considered as a potential payload for antibody drug conjugates (ADCs). However, the conjugated CFZ to anti-CD22 or anti-HER2 antibody totally abolishes the *in vitro* potency. This was a surprise since with other payloads such as monomethyl auristatin E (MMAE) where a potent anti-proliferation efficacy was retained as MMAE alone or as a payload in an ADC. Further investigations were conducted using CFZ alone, CFZ with a linker and CFZ-ADC with tissue matrices including lysosomal enzymes. With CFZ linked to the ADC, cathepsin B (a lysosomal enzyme) was efficient in liberating CFZ from the ADC by cleavage of valine-citrulline linker. At the same time, the liberated CFZ in the lysosome was inactivated due to further metabolism by lysosomal enzymes. The products from epoxide and amide hydrolysis were identified from these incubations. These results suggested that the CFZ-ADC upon uptake and internalization specifically delivers CFZ payload to the lysosomes, where the CFZ was inactivated. On the other hand, CFZ by itself is not as vulnerable and could reach its target. Therefore, the lysosomal stability is an important criterion in the selection of a payload for making the next generation of potent ADC therapeutics.

## **Introduction:**

Carfilzomib (CFZ) is an anti-cancer drug that can selectively inhibit the proteasome function in human cancer cells. Derived from a peptide epoxyketone nature product called expomicin, CFZ possesses a labile epoxy ketone pharmacophore (Kim and Crews, 2013). The epoxy ketone warhead in CFZ was demonstrated to have a high selectivity to form a covalent adduct with the 20S proteasome and thus disrupt the cellular protein homeostasis to induce the cell death. Despite of the essential role of proteasomes in all eukaryotic cells, a reasonable therapeutic window was still demonstrated for the proteasome inhibitors to selectively kill cancer cells (Adams, 2004a; Adams, 2004b). In 2012, CFZ was approved by FDA to treat multiple myeloma in patients who have received at least two prior therapies (the first generation proteasome inhibitor bortezomib and an immunomodulatory agent like thalidomide) and demonstrated disease progress in 60 days after the last therapy was finished (Herndon et al., 2013).

In multiple clinical trials, the treatment by CFZ alone or combined with dexamethasone has been demonstrated as tolerable and effective to induce responses in patients. However, the risk of severe adverse events (AEs) was also very high in patients receiving CFZ and even discontinued the treatment in some patients (Lendvai et al., 2014; Dimopoulos et al., 2016; Schlafer et al., 2017). The common AEs induced by CFZ include anaemia, hypertension, thrombocytopenia, and pneumonia. Like other small molecular cytotoxic agents, CFZ may largely decrease the health-related quality of life for patients in chemotherapy. Therefore, CFZ was tested as a payload of antibody drug conjugates (ADCs) to investigate the potential of improving its efficacy and safety. ADC technology is a demonstrated strategy to effectively enlarge the therapeutic windows of cytotoxic agents and minimize their side effects (Flygare et al., 2013; Polakis, 2016; Zhang et al., 2018). Also, because of the broad cytotoxicity to all cancer cells

containing proteasomes, conjugation to antibodies can help explore the usage of CFZ to treat cancers other than multiple myeloma. Recently, Staben et al. reported a new linker design strategy for payloads containing tertiary and heteroaryl amines, which was applied in the conjugation of CFZ to antibodies (Staben et al., 2016). The current article describes the challenges faced to investigate the feasibility to use CFZ as an ADC payload.

## **Experimental**

### **Materials and Methods:**

Ammonium acetate, sodium citrate, Tris-HCl, formic acid and cathepsin B were purchased from Sigma-Aldrich (St. Louis, MO). Pooled human liver S9 fractions were purchased from BD Biosciences (Bedford, MA). Human cellular lysosomal fractions prepared from human liver (mixed genders) were purchased from Xenotech (Lenexa, KS). Cancer cell lines including MDA-MB-231, MES-SA, MES-SA/Dx5, KPL-4, HCT116, A2058, BJAB, SK-BR-3 and WSU-DLCL2 were obtained from ATCC (Manassas, Virginia). These selected cell lines that include Human breast cancer cell lines MDA-MB-231 and KPL-4, human sarcoma cell lines MES-SA and MES-SA/Dx5, human colon cancer cell line HCT116, human melanoma cell line A2058, and human lymphoma cell line BJAB comprise a panel of human cancer cell lines representing various human cancers. The cell culture medium is DMEM/F-12 supplemented with 10% FBS. The medium and drug solution was not changed and replenished daily because the incubation time is short. The final DMSO concentration is 0.1 %. Vehicle controls were included in the study and cell viability in vehicle controls serves as the baseline (0% changes) to determine the cell viability at different drug concentrations. Maleimidocaproyl-valine-citrulline-*p*-aminobenzoyl quaternary ammonium (MC-ValCit-PABQ) linker-drug of CFZ (Sigma-Aldrich, St. Louis, MO) was synthesized as described previously (Staben et al., 2016). Human CD22 and HER2 antibodies with two engineered cysteine residues were generated as described previously (Polson et al., 2010; Bhakta et al., 2013). The anti-HER2 and anti-CD22 conjugates for CFZ were prepared and characterized as described previously (Zhang et al., 2016). The drug-to-antibody ratio (DAR) and aggregation percentage for the anti-HER2-CFZ conjugate (CNJ3264) were characterized as 1.9 and 2.2%, respectively. The DAR and aggregation percentage for the

anti-CD22-CFZ conjugate (CNJ3409) were characterized as 1.9 and 4%, respectively. The DAR and aggregation percentage for the anti-HER2-MMAE conjugate (CNJ1472) were characterized as 1.9 and < 5%, respectively. The DAR and aggregation percentage for the anti-CD22-MMAE conjugate (CNJ1994) were characterized as 1.9 and < 5%, respectively.

**Cell viability assay.** Cells were seeded in 96-well plates and allowed to adhere overnight at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>. The cell culture medium was then removed and replaced by 100 µl fresh medium containing various concentrations of small molecule drugs (stock solutions in DMSO) or antibody drug conjugates (stock solutions in 20 mM histidine-acetate, pH 5.5, 240 mM sucrose, and 0.02% Tween-20). After 4 days of drug incubation, the cell viability was determined using Promega CellTiter-Glo luminescent reagent, which measures ATP level (an indirect measure of cell number). The luminescent intensity was measured on PerkinElmer Envision reader. The relative cell viability was calculated by normalizing to non-drug treatment control and was graphed using GraphPad Prism software. IC<sub>50</sub> value was determined as the concentration to obtain 50% of the maximum cell killing.

***In vitro* immolation of linker-drug by cathepsin B.** The linker-drug was incubated at 15 µM in 100 mM citrate buffer (pH 5.5) containing 25 µg/mL cathepsin B and 5 mM cysteine at 37 °C. Aliquots were taken at 0, 1, 4, and 24 hours during the incubation. The samples were mixed with equal volume of acetonitrile, vortexed, and then centrifuged at 13,500 × g for 15 minutes. The supernatant was then analyzed by liquid chromatography-high resolution mass spectrometry (LC-HRMS).

**Stability of CFZ in cancer cell homogenates/human liver S9 fractions.** Cancer cells ( $5 \times 10^6$ ) were suspended and homogenized by beads in 0.5 mL ice-cold 100 mM Tris-HCl buffer (pH 7.4). The protein concentrations of cancer cell homogenates were determined by Bradford assay using

bovine serum albumin as the standard. CFZ (5  $\mu$ M) was incubated with 1 mg/mL cell homogenates and incubated at 37°C and aliquots were taken, processed, and analyzed in the same way described above. The stability of CFZ was also tested in the same buffer containing 1 mg/mL human liver S9 fractions.

**Metabolite identification of CFZ by LC-HRMS.** Identification of CFZ metabolites was conducted by LC-HRMS on a 1290 Infinity LC coupled with Thermo Orbitrap Elite mass spectrometer. The LC conditions were: column, Waters Atlantis dC18, 100 Å, 3  $\mu$ m, 75  $\mu$ m  $\times$  100 mm; mobile phase A, 0.1% formic acid in 10 mM ammonium acetate in H<sub>2</sub>O; mobile phase B, 0.1% formic acid and 10 mM ammonium acetate in 90% acetonitrile; gradient, 0–1.0 minutes 5% B, 1.0–16.0 minutes 5–34% B, 16.0–29.0 minutes 34–56 % B, 29.0–30.0 minutes 56–95% B, 30.0–31.0 minutes 95% B, 31.0–31.5 minutes 95–5% B; flow rate, 0.4 mL/min; column temperature, 25 °C. The cysteine adduct of MC-ValCit-PABQ-CFZ was identified by the molecular ion [M+H]<sup>2+</sup> at m/z found: 698.3723 and calculated: 698.3727, C<sub>72</sub>H<sub>103</sub>N<sub>12</sub>O<sub>15</sub>S<sup>2+</sup> and by major fragments at m/z: 720.4323, 659.2857, 633.3065, 571.2540, 554.2278 and 549.3073 (retention time: 24.0 minutes). CFZ was identified by the molecular ion [M+H]<sup>+</sup> at m/z found: 720.4312 and calculated: 720.4331, C<sub>40</sub>H<sub>58</sub>N<sub>5</sub>O<sub>7</sub><sup>+</sup> and by major fragments at m/z 702.4203, 566.3317, 549.3047, 521.3104, 432.2843, 402.2372, 289.1537, and 261.1589 (retention time: 27.9 minutes). M0 was identified by the molecular ion [M+H]<sup>+</sup> at m/z found: 589.2653 and calculated: 589.2650 C<sub>24</sub>H<sub>41</sub>N<sub>6</sub>O<sub>9</sub>S<sup>+</sup> and by major fragments at m/z 572.2375, 546.2584, 528.2480, 414.1688 and 386.1738 (retention time: 6.4 minutes). M1 was identified by the molecular ion [M+H]<sup>+</sup> at m/z found: 420.2492 and calculated: 420.2493 C<sub>22</sub>H<sub>34</sub>N<sub>3</sub>O<sub>5</sub><sup>+</sup> and by major fragments at m/z 289.1537 and 261.1589 (retention time: 13.7 minutes). M2 was identified by the molecular ion [M+H]<sup>+</sup> at m/z found: 307.1653 and calculated: 307.1652 C<sub>16</sub>H<sub>23</sub>N<sub>2</sub>O<sub>4</sub><sup>+</sup> and



by major fragments at  $m/z$  261.1592 and 100.0754 (retention time: 7.9 minutes). M3 was identified by the molecular ion  $[M+H]^+$  at  $m/z$  found: 738.4417 and calculated: 738.4436  $C_{40}H_{60}N_5O_8^+$  and by major fragments at  $m/z$  549.3052, 521.3109, 450.2951, 402.2375, 337.2112, 289.1540 and 261.1591 (retention time: 22.4 minutes).

## **Results**

As a free drug, CFZ is an excellent cytotoxic agent which can effectively kill cancer cells at nanomolar concentrations. Consistent with previous studies, free CFZ has been tested to be very cytotoxic to multiple cancer cell lines in our study (Figure 1) (Kuhn et al., 2007; Baker et al., 2014). However, when CFZ was conjugated to antibodies, the conjugates were ineffective to kill cancer cells. An ADC comprising anti-HER2 antibody and CFZ is unable to inhibit the proliferation of the HER2-positive human breast cancer cell lines SKBR-3 and KPL-4 at a concentration which is equivalent to as high as 1  $\mu$ M CFZ (Figure 2, A and B). Also, CFZ conjugated to anti-CD22 antibody shows no efficacy in the CD22-positive cancer cell lines BJAB and WSU-DLCL2 (Figure 2, C and D). As the positive control, MMAE, a peptide-like cytotoxic agent employed as the payload in the marketed ADC brentuximab vedotin (Adcetris®), shows a great cell-killing activity in those cell lines after conjugated to the same anti-HER2 or -CD22 antibodies (Figure 2).

To determine whether CFZ can be successfully released from ADC in lysosomes, an *in vitro* assay was conducted using purified cathepsin B and the linker-drug as a surrogate to determine whether the proteolytic cleavage of linker can occur (Figure 3A). LC-HRMS analysis indicated the rapid disappearance of cysteine adduct of MC-ValCit-PABQ-CFZ in the incubation (Figure 3A). In the first hour, more than 95% of linker-drug disappeared. The cysteine adduct of the MC-vc linker residue (M0) was detected as a product from the cleavage and the peak areas of M0 at 1, 4, and 24 hours were comparable to each other. This is consistent with the previous study in which the MC-ValCit-PABQ linker has been demonstrated to be efficient in releasing payloads conjugated via a quaternary ammonium (Staben et al., 2016). Unfortunately, no free CFZ was detected in the incubation, nor the possible PABQ-drug intermediate was detected (Figure 3, A).

Given the fact that CFZ has a peptide backbone, the proteolytic cleavage by the lysosomal cathepsin(s) might result in the liability of unconjugated CFZ in the lysosomes.

*In vitro* incubations of CFZ with lysosomal fractions provide some insights. The incubation of CFZ with lysosomal fractions prepared from human livers at pH 5.5 showed that almost 100% CFZ disappeared rapidly in one hour (Figure 3, B). Two metabolites were detected in the incubation and they were more hydrophilic based on retention times and with lower molecular weights than the parent CFZ. Based on the exact mass and MS fragmentation patterns, the two metabolites were identified as hydrolytic metabolites (Figure 4, B and C). The same metabolites M1 and M2 were also observed in the incubation of MC-ValCit-PABQ-CFZ with cathepsin B (Figure 3, A), indicating that cathepsin B hydrolyzed not only the amide bond in the valine-citrulline dipeptide and the PAB spacer, but also the amide bonds within CFZ. Besides cathepsin B, many other members in the cathepsin family have been identified in the lysosomes of human cancer cells (Nomura and Katunuma, 2005; Caculitan et al., 2017). It is unknown whether the cathepsins other than cathepsin B can also catalyze the hydrolysis of CFZ.

Further investigation was conducted to determine whether the non-lysosomal hydrolases inside the cells can also contribute to the hydrolysis of CFZ. At 37°C and pH 7.4, moderate hydrolysis of CFZ was observed in the homogenates of all the three cancer lines (KPL4, SKBR3, and BJAB) tested. After 1 hour of incubation in the homogenates of KPL4, SKBR3, and BJAB cells, approximately 40% of the CFZ (starting concentration 5 µM) disappeared, respectively (Figure 5). The hydrolytic metabolites with cancer cell fractions were similar to those observed in the lysosomal fraction hydrolysis of CFZ. For comparison, we also conducted the incubation of CFZ in human liver S9 fractions at pH 7.4 because it provides an abundant resource of hydrolysis enzymes (Fiddler et al., 1979). Interestingly, the peptide backbone of CFZ was found to be very

stable in the human liver S9 fractions incubation at physiological pH, but the epoxy group was hydrolyzed to give a dihydrodihydroxy metabolite M3 (Figure 3, C and Figure 4, D). This reaction is likely catalyzed by an epoxide hydrolase under a neutral pH condition (Arand et al., 2005). We proposed that hydrolysis of CFZ in the cytosol of cancer cells can impair its efficacy. However, the low  $IC_{50}$  values of free CFZ in various cancer cell lines indicated that the impact of cytosolic hydrolysis on its efficacy is very limited, perhaps because the hydrolysis rates are relatively slow and enough free CFZ can bind to the proteasomes before extensive hydrolysis occurs.

## **Discussion**

In the current study, the feasibility of utilizing CFZ, a drug used to treat multiple myeloma, was investigated as an ADC payload. An ADC underwent antigen binding, tissue uptake, cell internalization, and biochemical transformation (proteolytic degradation of antibody, linker cleavage, and immolation) to release payload in lysosomes. The biotransformation pathways associated with CFZ and its linker drug in lysosomes are summarized in Figure 6. CFZ was hydrolyzed to multiple metabolites in lysosomes. CFZ is highly cytotoxic to cancer cell lines by diffusing into the cells and covalently binding to the 20S subunit of proteasomes to illicit its action (Figure 7, A). However, the mechanism of action of ADC is much more complicated and involves a series of events including antibody-antigen binding, endosome incorporation, lysosomal catalysis, linker cleavage (and immolation), payload relocation and finally the cell-killing event (Schrama et al., 2006). ADC may fail to show efficacy if any one of these steps fails. In the current study, *in vitro* assays using cathepsin B and lysosomal fractions have demonstrated that CFZ is susceptible to hydrolysis in the lysosomes. As a part of the incorporated ADC, CFZ may be rapidly hydrolyzed by cathepsin(s) during lysosomal catalysis

before or after any CFZ molecule is released from the ADC. It should be noted that due to lysosomal preparations, the protein concentration in the *in vitro* incubation was diluted to 0.3 mg/mL, while the actual protein concentration inside the lysosomes of living cells can be as high as 200 mg/mL (Dean and Barrett, 1976). It can be expected that there is no or very limited escape of CFZ off an ADC from the lysosomes to reach the target and thus no proteasome adduction and dysfunction will occur (Figure 7, B). In contrast, MMAE exemplifies that good lysosomal stability can be critical to make an excellent payload for ADC. MMAE embodies 4 amide bonds in its structure, but it is 100% resistant to lysosomal hydrolysis under the similar *in vitro* incubation conditions (Doronina et al., 2003). After ADC incorporation and linker cleavage and immolation, the release of free MMAE from lysosomes to cytosol is a necessary step before MMAE can block the polymerisation of tubulin and exert antineoplastic effects (Francisco et al., 2003). Retrospectively, a conclusion can be drawn that the stability of MMAE in lysosomes is a prerequisite for the efficacy and successful development of brentuximab vedotin. The experiences and lessons learned from MMAE and CFZ have indicated that the lysosomal stability should be considered as one of the criteria to evaluate the potential of a compound to serve as the cytotoxic warhead in ADC. At the early stage in the discovery of new payloads, *in vitro* incubation with cathepsin(s) or lysosomal fractions is a very helpful assay in evaluating the lysosomal stability of the lead compounds, especially those with functional groups vulnerable to hydrolases (e.g., amides and esters). In future, a more quantitative relationship between the rate of hydrolysis *in vitro* and the anti-cancer efficacy may be established for new payloads with designing experiments to learn the kinetics of hydrolysis reactions.

ADC technology makes it possible to deliver the cytotoxins to specific antigen-bound cancer cells and possibly minimize the toxicity-induced side effects in normal tissues and organs,

enabling the application of these cytotoxins in cancer chemotherapy (Flygare et al., 2013; Polakis, 2016). In addition, ADC could enable the delivery of payloads with poor cell membrane permeability into cancer cells and consequently enhance their efficacy. Several categories of substances are usually being investigated as ADC payloads, which are mainly anti-mitotic and DNA-damaging agents (McCombs and Owen, 2015). Meanwhile, researchers are also looking for novel ADC payloads which can act via new targets or pathways, or trying to conjugate existed small molecule drugs to antibodies for improving their efficacy or safety.

In summary, CFZ is a potent cytotoxic agent in various cancer cell lines, however, CFZ is an inactive payload in ADCs. Our studies demonstrated that CFZ is highly vulnerable to hydrolysis in lysosomal fractions. CFZ conjugated to antibodies loses its cytotoxic potency because of the extensive hydrolysis of the drug occurring in lysosomes, which is necessary for the ADC catabolism. The results indicate that when selecting an appropriate payload in the development of ADC, the lysosomal stability is a property of payloads which should be taken into considerations.

Authors were employees of Genentech, Inc.

Current address: Yong Ma, Calithera Biosciences, 343 Oyster Point Boulevard, Suite 200, South San Francisco, CA, 94080

Author contributions:

Research design: Ma, Cruz-Chuh, Khojasteh, Dragovich, Pillow, Zhang

Experimental: Ma, Cruz-Chuh

Contributed to new reagents: Pillow

Performed data analysis: Ma, Cruz-Chuh, Khojasteh, Zhang

Contributed to new reagents: Ma, Khojasteh, Dragovich, Zhang

## References:

- Adams J (2004a) The development of proteasome inhibitors as anticancer drugs. *Cancer Cell* **5**:417-421.
- Adams J (2004b) The proteasome: a suitable antineoplastic target. *Nat Rev Cancer* **4**:349-360.
- Arand M, Cronin A, Adamska M, and Oesch F (2005) Epoxide Hydrolases: Structure, Function, Mechanism, and Assay. *Methods in Enzymology* **400**:569-588.
- Baker AF, Hanke NT, Sands BJ, Carbajal L, Anderl JL, and Garland LL (2014) Carfilzomib demonstrates broad anti-tumor activity in pre-clinical non-small cell and small cell lung cancer models. *Journal of Experimental & Clinical Cancer Research* **33**:111.
- Bhakta S, Raab H, and Junutula JR (2013) Engineering THIOMABs for Site-Specific Conjugation of Thiol-Reactive Linkers, in: *Antibody-Drug Conjugates* (Ducry L ed), pp 189-203, Humana Press, Totowa, NJ.
- Caculitan NG, dela Cruz Chuh J, Ma Y, Zhang D, Kozak KR, Liu Y, Pillow TH, Sadowsky J, Cheung TK, Phung Q, Haley B, Lee B-C, Akita R, Sliwkowski MX, and Polson AG (2017) Cathepsin B is dispensable for cellular processing of cathepsin B-cleavable antibody-drug conjugates. *Cancer Research:canres.2391.2017*.
- Dean RT and Barrett AJ (1976) Lysosomes. *Essays in biochemistry* **12**:1-40.
- Dimopoulos MA, Moreau P, Palumbo A, Joshua D, Pour L, Hajek R, Facon T, Ludwig H, Oriol A, Goldschmidt H, Rosinol L, Straub J, Suvorov A, Araujo C, Rimashevskaya E, Pika T, Gaidano G, Weisel K, Goranova-Marinova V, Schwarzer A, Minuk L, Masszi T, Karamanesht I, Offidani M, Hungria V, Spencer A, Orlowski RZ, Gillenwater HH, Mohamed N, Feng S, Chng WJ, and Investigators E (2016) Carfilzomib and dexamethasone versus bortezomib and dexamethasone for patients with relapsed or refractory multiple myeloma (ENDEAVOR): a randomised, phase 3, open-label, multicentre study. *The Lancet Oncology* **17**:27-38.
- Doronina SO, Toki BE, Torgov MY, Mendelsohn BA, Cerveny CG, Chace DF, DeBlanc RL, Gearing RP, Bovee TD, Siegall CB, Francisco JA, Wahl AF, Meyer DL, and Senter PD (2003) Development of potent monoclonal antibody auristatin conjugates for cancer therapy. *Nature Biotechnology* **21**:778.
- Fiddler MB, Ben-Yoseph Y, and Nadler HL (1979) Binding of human liver hydrolases by immobilized lectins. *Biochemical Journal* **177**:175-180.
- Flygare JA, Pillow TH, and Aristoff P (2013) Antibody-Drug Conjugates for the Treatment of Cancer. *Chemical Biology & Drug Design* **81**:113-121.
- Francisco JA, Cerveny CG, Meyer DL, Mixan BJ, Klussman K, Chace DF, Rejniak SX, Gordon KA, DeBlanc R, Toki BE, Law C-L, Doronina SO, Siegall CB, Senter PD, and Wahl AF (2003) cAC10-vcMMAE, an anti-CD30-monomethyl auristatin E conjugate with potent and selective antitumor activity. *Blood* **102**:1458-1465.
- Herndon TM, Deisseroth A, Kaminskas E, Kane RC, Koti KM, Rothmann MD, Habtemariam B, Bullock J, Bray JD, Hawes J, Palmby TR, Jee J, Adams W, Mahayni H, Brown J, Dorantes A, Sridhara R, Farrell AT, and Pazdur R (2013) U.S. Food and Drug Administration Approval: Carfilzomib for the Treatment of Multiple Myeloma. *Clinical Cancer Research* **19**:4559.
- Kim KB and Crews CM (2013) From epoxomicin to carfilzomib: chemistry, biology, and medical outcomes. *Natural Product Reports* **30**:600-604.
- Kuhn DJ, Chen Q, Voorhees PM, Strader JS, Shenk KD, Sun CM, Demo SD, Bennett MK, van Leeuwen FWB, Chanan-Khan AA, and Orlowski RZ (2007) Potent activity of carfilzomib, a novel, irreversible inhibitor of the ubiquitin-proteasome pathway, against preclinical models of multiple myeloma. *Blood* **110**:3281-3290.



- Lendvai N, Hilden P, Devlin S, Landau H, Hassoun H, Lesokhin AM, Tsakos I, Redling K, Koehne G, and Chung DJ (2014) A phase 2 single-center study of carfilzomib 56 mg/m<sup>2</sup> with or without low-dose dexamethasone in relapsed multiple myeloma. *Blood* **124**:899-906.
- McCombs JR and Owen SC (2015) Antibody Drug Conjugates: Design and Selection of Linker, Payload and Conjugation Chemistry. *The AAPS Journal* **17**:339-351.
- Nomura T and Katunuma N (2005) Involvement of cathepsins in the invasion, metastasis and proliferation of cancer cells. *The Journal of Medical Investigation* **52**:1-9.
- Polakis P (2016) Antibody Drug Conjugates for Cancer Therapy. *Pharmacological Reviews* **68**:3-19.
- Polson AG, Williams M, Gray AM, Fuji RN, Poon KA, McBride J, Raab H, Januario T, Go M, Lau J, Yu SF, Du C, Fuh F, Tan C, Wu Y, Liang WC, Prabhu S, Stephan JP, Hongo JA, Dere RC, Deng R, Cullen M, de Tute R, Bennett F, Rawstron A, Jack A, and Ebens A (2010) Anti-CD22-MCC-DM1: an antibody-drug conjugate with a stable linker for the treatment of non-Hodgkin's lymphoma. *Leukemia* **24**:1566-1573.
- Schlafer D, Shah KS, Panjic EH, and Lonial S (2017) Safety of proteasome inhibitors for treatment of multiple myeloma. *Expert Opinion on Drug Safety* **16**:167-183.
- Schrama D, Reisfeld RA, and Becker JC (2006) Antibody targeted drugs as cancer therapeutics. *Nat Rev Drug Discov* **5**:147-159.
- Staben LR, Koenig SG, Lehar SM, Vandlen R, Zhang D, Chuh J, Yu S-F, Ng C, Guo J, Liu Y, Fourie-O'Donohue A, Go M, Linghu X, Segraves NL, Wang T, Chen J, Wei B, Phillips GDL, Xu K, Kozak KR, Mariathasan S, Flygare JA, and Pillow TH (2016) Targeted drug delivery through the traceless release of tertiary and heteroaryl amines from antibody-drug conjugates. *Nat Chem* **8**:1112-1119.
- Zhang D, Yu S-F, Khojasteh SC, Ma Y, Pillow TH, Sadowsky JD, Su D, Kozak KR, Xu K, Polson AG, Dragovich PS, and Hop CECA (2018) Intratumoral Payload Concentration Correlates with the Activity of Antibody-Drug Conjugates. *Molecular Cancer Therapeutics* **17**:677-685.
- Zhang D, Yu S-F, Ma Y, Xu K, Dragovich PS, Pillow TH, Liu L, Del Rosario G, He J, Pei Z, Sadowsky JD, Erickson HK, Hop CE, and Khojasteh SC (2016) The chemical structure and concentration of intratumor catabolites determine the efficacy of antibody drug conjugates. *Drug Metabolism and Disposition*:dmd.116.070631.

List of Figures:

Figure 1. Cytotoxicity of CFZ in various cancer cell lines

IC<sub>50</sub>s: MDA-MB-231, 9.3 nM; MES-SA, 4.1 nM; MES-SA/Dx5, 11.8 nM; KPL-4, 19.7 nM; HCT116, 4.8 nM; A2058, 4.6 nM; BJAB, 1.9 nM

Figure 2. Cytotoxicity of anti-HER2 and CD22 antibody conjugates of CFZ and MMAE in various cancer cell lines (A) SK-BR-3, (B) KPL-4, (C) BJAB, (D) WSU-DLCL2

Figure 3. Selected ion chromatograms of lysosomal metabolic products of CFZ or its linker drug.

(A) MC-ValCit-PABQ-CFZ in incubation with cathepsin B at pH 5.5; (B) CFZ in incubation with lysosomal fractions prepared from human liver at pH 5.5; (C) CFZ in incubation with human liver S9 at pH 7.4.

Figure 4. Product ion scans of CFZ (A) and metabolites M1 (B), M2 (C), and M3 (D) formed in lysosomal and S9 incubations

Figure 5. Disappearance of CFZ in incubations with various cellular fractions of cancer cells or buffers

Figure 6. Lysosomal degradation pathways of CFZ linker-drug

Figure 7. Proposed cellular distribution and trafficking of (A) CFZ and (B) antibody-CFZ conjugate

Figure 1




Figure 2




Figure 3





Figure 4



Downloaded from dmd.aspetjournals.org at ASPET Journals on April 20, 2024

Figure 5




Figure 6






Figure 7

DMD/2019/086595

