

DMD # 89557

Acetaminophen-induced Liver Injury Alters Expression and Activities of Cytochrome P450 Enzymes in an Age-dependent Manner in Mouse Liver

Yifan Bao, Pei Wang, Xueyan Shao, Junjie Zhu, Jingcheng Xiao, Jian Shi, Lirong Zhang, Hao-Jie Zhu, Xiaochao Ma, José E. Manautou, and Xiao-bo Zhong

Department of Pharmaceutical Sciences, School of Pharmacy, University of Connecticut, Storrs, Connecticut (Y.B., P.W., X.S., J.E.M., X.B.Z.); Department of Pharmacology, School of Basic Medical Sciences, Zhengzhou University, Zhengzhou, Henan, China (P.W., L.Z.); Department of Pharmaceutical Sciences, School of Pharmacy, University of Pittsburgh, Pittsburgh, Pennsylvania (J.Z., X.M.); Department of Pharmaceutical Sciences, College of Pharmacy, University of Michigan, Ann Arbor, Michigan (J.X); and Department of Clinical Pharmacy, College of Pharmacy, University of Michigan, Ann Arbor, Michigan (J.S., H.J.Z.)

DMD # 89557

Running Title: Alterations of P450s by APAP-induced liver injury

Corresponding author: Dr. Xiao-bo Zhong, Department of Pharmaceutical Sciences, School of Pharmacy, University of Connecticut, 69 N Eagleville Road, Storrs, Connecticut 06269, USA. Telephone: 860-486-3697; Fax: 860-486-5792; E-mail: xiaobo.zhong@uconn.edu

Number of text pages: 37

Number of figures: 7

Number of tables: 2

Number of references: 47

Number of words in Abstract: 246

Number of words in Introduction: 538

Number of words in Discussion: 1405

DMD # 89557

ABBREVIATIONS:

ALLI, acetaminophen-induced liver injury; ALT, alanine aminotransferase; ANOVA, one-way analysis of variance; APAP, acetaminophen; AST, aspartate aminotransferase; BSA, bovine serum albumin; CHZ, chlorzoxazone; DILI, drug-induced liver injury; EFV, efavirenz; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; H&E, hematoxylin and eosin; HCC, hepatocellular carcinoma; LC-MS, liquid chromatography-tandem mass spectrometry; lncRNA, long non-coding RNA; MDZ, midazolam; miRNA, microRNA; NAFLD, non-alcoholic fatty liver disease; NAPQI, *N*-acetyl-para-benzo-quinone imine; P450s, cytochrome P450s; PBS, phosphate-buffered saline; RSF, resorufin; RSG, rosiglitazone; RT-PCR, real-time polymerase chain reaction; SD, standard deviation; UNL, upper limit of normal level; UPLC-QTOFMS, ultra-performance liquid chromatography-quadrupole time of flight mass spectrometry.

DMD # 89557

ABSTRACT

Drug-induced liver injury (DILI) is a global medical problem. The risk of DILI is often related to expression and activities of drug-metabolizing enzymes, especially cytochrome P450s (P450s). However, changes on expression and activities of P450s after DILI have not been determined. The aim of this study is to fill this knowledge gap. Acetaminophen (APAP) was used as a model drug to induce DILI in C57BL/6J mice at different ages of day 10 (infant), 22 (child), and 60 (adult). DILI was assessed by levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) in plasma with a confirmation by H&E staining on liver tissue sections. The expression of selected P450s at mRNA and protein levels was measured by RT-PCR and LC-MS/MS, respectively. The activities of these P450s were determined by the formation of metabolites from probe drugs for each P450 using UPLC-QTOFMS. DILI was induced at mild to severe levels in a dose-dependent manner in 200, 300, and 400 mg/kg APAP treated groups at child and adult ages, but not at the infant age. Significantly decreased expression at mRNA and protein levels as well as enzymatic activities of CYP2E1, 3A11, 1A2, and 2C29 was found at child and adult ages. Adult male mice were more susceptible to ALIL than female mice with more decreased expression of P450s. These results implicate that altered levels of P450s in severely injured livers caused by drugs may affect the therapeutic efficacy of drugs, which are metabolized by P450s, more particularly for males.

DMD # 89557

Significance Statement

The current study in an animal model demonstrates that ALI results in decreased expression and enzyme activities of several examined drug-metabolizing P450s. The extent of such decreases is correlated to the degree of liver injury severity. The generated data may be translated to human health for patients who have DILI with decreased capability to metabolize drugs by certain P450s.

DMD # 89557

Introduction

Drug therapy is an important part of the current practical medicine to treat human diseases, but has a potential risk to develop drug-induced liver injury (DILI). The incidence data in various populations in Europe (Sgro et al., 2002), North America (Bell and Chalasani, 2009; Leise et al., 2014; Vega et al., 2017), and Asia (Shen et al., 2019) indicate that there are approximately 10-24 new cases of DILI per 100,000 persons each year. In the over 7.7 billion global individuals in 2019 documented by the World Population Prospects of the United Nations, the estimated total cases of DILI may reach to 0.7-1.4 million worldwide. More than 900 drugs have been implicated in causing DILI (Larrey, 2002; Maddrey, 2005; Hoofnagle et al., 2013). The current drug therapy approaches do not specially consider whether patients with DILI have different efficacy and risks for adverse reactions in comparison to patients who do not have DILI.

Efficacy and adverse reactions of drug therapy are largely relied on drug metabolism in liver. The role of drug metabolism in causing DILI has been defined (Corsini and Bortolini, 2013). Especially, drug metabolism by phase-I enzymes of cytochrome P450s (P450s) can lead to formation of toxic metabolites in liver (Pasanen et al., 1995; Brewer and Chen, 2017; Hassan et al., 2018). One of such examples is P450-mediated metabolism of acetaminophen (APAP), a popular analgesic drug. APAP is the leading cause of acute liver failure in the United States (Fontana, 2008; Yoon et al., 2016). APAP-induced liver injury (ALI) is mainly due to the formation of a toxic metabolite of *N*-acetyl-para-benzo-quinone imine (NAPQI) in liver during APAP metabolism by CYP2E1, 1A2, and 3A4 (Dahlin et al., 1984; Yoon et al., 2016). Increased enzyme activities of CYP2E1, 1A2, and 3A4 in liver can lead to increased susceptibility of ALI.

A knowledge gap exists in the current practice of pharmacotherapy on whether injured liver caused by DILI alters drug metabolism by P450s, resulting in changes in efficacy and adverse

DMD # 89557

reactions of drug therapy. We hypothesize that DILI results in altered hepatic expression and activities of several P450s. To test the hypothesis, the current study used APAP as a model drug and mouse as the animal model to systematically determine the impact of AILI on expression and activities of P450s in liver at different ages. APAP is the leading cause of DILI worldwide and its toxicity is dose-dependent (Bunchorntavakul and Reddy, 2013). Comparing *in vivo* and *in vitro* models for AILI, the mouse model has the most similarity in toxicity initiation with human primary hepatocytes and toxic doses have been well defined (Jaeschke et al., 2014; Xie et al., 2014). Mouse CYP3A11 contributes to convert APAP to NAPQI (Guo et al., 2004). In consideration of the changes of expression and activities of P450s in mouse liver during postnatal maturation (Hart et al., 2009; Peng et al., 2012), we hypothesized that the consequence of APAP treatment on P450 expression is in an age-dependent manner. The current study determined expression and activities of various key drug-metabolizing P450s (CYP1A2, 2B10, 2C29, 2E1, and 3A11) in mouse liver at different ages of postnatal day 10 (infant), 22 (child), or 60 (adult) with different levels of AILI caused by exposure to different doses of APAP.

Materials and Methods

Chemicals and Reagents. Phosphate-buffered saline (PBS), APAP, resorufin (RSF), and midazolam (MDZ) were purchased from Sigma-Aldrich (St. Louis, MO). Buffered formalin (10%) was purchased from Fisher Science (Fair Lawn, NJ). Efavirenz (EFV) was purchased from TCI American (Portland, OR). 1-OH-MDZ and 8-OH-EFV were purchased from Toronto Research Chemicals (Ontario, Canada). Rosiglitazone (RSG), demethyl-RSG, chlorzoxazone (CHZ), 7-

DMD # 89557

ethoxyresorufin (7-ethoxy-RSF), and 6-OH-CHZ were obtained from Cayman Chemical (Ann Arbor, MI).

Animals and Drug Treatment. C57BL/6J mice were obtained from Jackson Laboratory (Bar Harbor, ME). Mice were bred under standard conditions and housed in compliance with the animal care guidelines outlined by the American Association for Animal Laboratory Sciences in the Animal Resources Facility at the University of Connecticut. All animal procedures were approved by the University of Connecticut's Institutional Animal Care and Use Committee. At the age of 6 weeks, mice were set up into breeding pairs to produce sufficient mice at postnatal day 10 (infant), 22 (child), or 60 (adult). Given the fact that male mice are more susceptible to APAP than female mice (Dai et al., 2006), male mice were used to investigate the role of age- and dose-dependency in ALI in the current study. In brief, male mice at different ages (postnatal day 10, 22, or 60) were intraperitoneally administered with a single dose of APAP (200, 300, or 400 mg/kg) or PBS (vehicle control). To study the different effects of ALI by gender, both male and female mice at day 60 after birth were included. Before 16 hrs and after 8 hrs of APAP dosing, mice at day 22 and 60 were fasted with free access to water to avoid food-induced changes in liver genes (Gavito et al., 2016). All the mice were sacrificed at 24 hrs after APAP treatment to collect blood and liver samples. A part of the liver was fixed with 10% formalin buffer for histopathological analysis. Remaining other liver tissues were snap frozen in liquid nitrogen and then stored at -80°C for further gene expression analysis.

Plasma Alanine Aminotransferase (ALT) and Aspartate Aminotransferase (AST) Assay. Blood samples were collected in anticoagulant-coated tubes (Fisher Scientific, Fair Lawn, NJ) and centrifuged at 3,000 g for 10 min to obtain plasma. Levels of plasma ALT and AST were measured as biomarkers of hepatocellular injury. Scientific™ AST/GOT and

DMD # 89557

ALT/GPT Reagents from Thermo Fisher (Waltham, MA) were used to determine AST and ALT levels, respectively, with a method described previously (Rudraiah et al., 2014). A BioTek Power Wave X Spectrophotometer (BioTek, Winooski, VT) was used to analyze the samples.

Hematoxylin & Eosin (H&E) Staining. Liver samples were fixed with 10% buffered formalin and kept at 4°C for H&E staining. The embedding and slicing process of H&E staining was performed by the Connecticut Veterinary Medical Diagnostic Laboratory (<http://cvmdl.uconn.edu/>). Images were taken by a phase-contrast microscope with a 20 × objective lens. The scale bars were added by Image J software.

Quantification of P450 mRNAs by Real-Time PCR (RT-PCR). Total RNAs were isolated from frozen liver tissues using a TRIzol™ reagent (Life Technologies, Guilford, CT) according to the manufacturer's protocol. RNA concentrations were measured by a NanoDrop spectrophotometer from NanoDrop Technologies (Wilmington, DE) at 260 nm and mRNA integrity was evaluated using an Agilent 2,200 Tape Station from Agilent Technologies (Santa Clara, CA). To obtain cDNAs, 1 µg of total mRNAs were reversely transcribed to cDNAs using an iScript™ cDNA Synthesis kit from Bio-Rad Laboratories (Hercules, CA). RT-PCR reactions were performed using a CFX96™ Real-Time System from Bio-Rad Laboratories (Hercules, CA) with the TaqMan gene expression assays for glyceraldehyde-3-phosphate dehydrogenase (GAPDH), CYP1A2, 2B10, 2C29, 2E1, and 3A11 from Thermo Fisher (Waltham, MA). Fold changes of P450 expression in the APAP-treated groups compared to the PBS controls at each age were calculated using the $2^{-\Delta\Delta C_t}$ method with an internal normalization to GAPDH. The values were calculated for $\text{Log}_2(\text{fold changes})$.

Quantification of P450 Proteins by Mass Spectrometry. Frozen liver tissues were chopped and homogenized in a PBS buffer (pH 7.4). Protein samples were prepared and quantified by a previously described method using liquid chromatography-tandem mass

DMD # 89557

spectrometry (LC-MS/MS)-based approach (Piekos et al., 2018). Briefly, liver microsomes were obtained by centrifugation. The microsomes were mixed with an internal standard bovine serum albumin (BSA) and digested by lysyl endopeptidase and trypsin. The digested peptides were extracted using Waters Oasis HLB columns (Waters Corporation, Milford, MA) and analyzed on a Triple TOF 5,600+ mass spectrometer (AB Sciex, Framingham, MA) coupled with an Eksigent 2D Plus LC System (Eksigent Technologies, Dublin, CA). The proteomics data obtained from the sequential window acquisition of all theoretical mass spectra (SWATH) method were analyzed using the Spectronaut software (version 13.0, Biognosys AG, Schlieren, Switzerland) with the reference spectral library generated from the previous information-dependent acquisition searches (Piekos et al., 2018). The MS2 peak areas of top 3 signature peptides of each protein were used for quantification with the default settings (BGS Factory Setting). Data were normalized using the "Local Normalization" strategy embedded in the Spectronaut™ software.

Quantification of P450 Enzyme Activities by Mass Spectrometry. S9 fractions of liver tissues were isolated by centrifugation with a previously described method (Piekos et al., 2018). To determine the enzyme activities of CYP1A2, 2B10, 2C29, 2E1, and 3A11, the reaction rates of 7-ethoxy-RSF to RSF, EFV to 8-OH-EFV, RSG to demethyl-RSG, CHZ to 6-OH-CHZ, and MDZ to 1-OH-MDZ were measured, respectively. P450 enzyme activities in the S9 fractions were determined by ultra-performance liquid chromatography-quadrupole time of flight mass spectrometry (UPLC-QTOFMS) as previously described (Tien et al., 2015).

Quantification of *in vitro* Metabolism of APAP. Liver microsomes were isolated from liver tissues by centrifugation with a previously described method (Piekos et al., 2018). Microsome proteins were determined using a Qubit Protein Assay Kit (Thermo Fisher Scientific, Rockford, IL). The metabolism of APAP was initiated by adding 1 ml of 15 mM N-

DMD # 89557

acetyl cysteine and 1 ml of 3 μ M NADPH to 0.5 ml solution containing 3 mg microsomal proteins and incubated at 37°C for 10 min. One ml of 60 mM APAP solution was added subsequently and incubated at 37°C for 30 min. The reaction was terminated by adding 3 ml of ice-cold methanol. After centrifuged at 2,800 *g* for 30 min and filtered through 0.22 μ m Millipore filter, 20 μ l aliquot was injected into a Shimadzu CTO-20A/2-AC HPLC-UV system for quantification of APAP. Chromatographic separation of metabolites was performed on an Agilent 5 HC-C18(2) column (4.6 \times 250 mm, 5 μ m; Agilent Technologies). Mobile phase A consisted of 1% glacial acetic acid in water and mobile phase B was methanol. The flow rate of the mobile phase was 0.80 ml/min and the column temperature was maintained at 25°C. The absorbance wavelength of photodiode array detector was 254 nm. The rate of APAP metabolism was calculated by the decreased amount of APAP divided by the amount of microsomal proteins within the reaction time.

Quantification of Total Liver Proteins. A 5-mg frozen liver sample was homogenized with 300 μ l RIPA buffer (Thermo Fisher Scientific, Rockford, IL). After centrifuged at 14,000 *g* for 30 min at 4°C, total protein concentration in the supernatant was determined by a Qubit Protein Assay Kit (Thermo Fisher Scientific, Rockford, IL).

Statistical Analysis. The data were presented as mean \pm SD (standard deviation). Statistical analysis was performed by one-way analysis of variance (ANOVA) with the Dunnett's post-hoc test using the GraphPad Prism 7 software program from GraphPad Software, Ln. (La Jolla, CA). A value of $p < 0.05$ was considered to be statistically significant.

Results

DMD # 89557

The normal ALT and AST levels in plasma of adult C57BL/6J mice collected by cardiac puncture have been defined (Fernández et al., 2010; Otto et al., 2016). In male adult mice, the upper limit of normal level (UNL) of ALT is 94 U/L, while the UNL of AST is 122 U/L. The grade of liver injury can be estimated by using the UNL of ALT and AST of mice and the grading standard of human (Hoofnagle et al., 2013). Grade 0 has a level of ALT or AST within 1.25-fold higher than the UNL. A level of ALT or AST is considered as Grade 1 (mild) between 1.25- and 2.5-fold, Grade 2 (moderate) between 2.5- and 5.0-fold, Grade 3 (severe) between 5.0- and 10-fold, and Grade 4 (extreme severe and life-threatening) over 10-fold, respectively. The grading system based on ALT and AST levels for mice is summarized in Table 1. A grading system of liver injury based on Ishak scores on the morphology changes is shown in Table 2 (Goodman, 2007).

In this study, male mice at different ages (day 10, 22, or 60 after birth) were administrated with a single dose of APAP (200, 300, or 400 mg/kg) to investigate the role of ontogeny of P450s in the severity of ALI as well as the impact of ALI on the expression of P450s. To study the gender differences, mice at day 60 were used. Briefly, both male and female mice were treated with different doses of APAP (200, 300, or 400 mg/kg). The levels of ALT and AST and the hepatic expression of P450s were determined.

Age Affected the Severity of ALI in Male Mice. As seen in Fig. 1A, APAP treatment only led to a higher ALT level in mice at day 10 after birth at the highest dose (400 mg/kg). Whereas, even with the lowest dose of APAP (200 mg/kg), plasma ALT levels in male mice dramatically increased at day 22 and 60. Moreover, higher ALT levels were observed in the groups administered with higher doses of APAP in the mice at day 22 and 60. For instance, the mice at day 60 treated with APAP at 200, 300, or 400 mg/kg resulted in ALT levels of 8.57- (grade 3 injury), 10.69- (grade 4 injury), or 14.83-fold (grade 4 injury) of the UNL, respectively. When

DMD # 89557

treated with a same dose of APAP, the ALT levels were found to be the highest in the mice at day 60 and the lowest in the mice at day 10. For instance, the ALT levels in the mice treated with 400 mg/kg APAP at day 10, 22, or 60 were 66, 472, or 1394 U/L, respectively.

Similar findings were observed with AST values (Fig. 1B). Male mice treated with 200 or 300 mg/kg APAP showed increased AST levels at day 22 and 60, but not at day 10. APAP treatment at 400 mg/kg led to AST levels of 2.39- (grade 1 injury), 5.83- (grade 3), or 10.02-fold (grade 4) of the UNL at day 10, 22, or 60, respectively. At the same age, AILI was more severe in the group with a higher dose of APAP exposure. For the mice at day 60, APAP treatment with a dose of 200, 300, or 400 mg/kg resulted in AILI at grade 3, 3, or 4, respectively.

H&E staining is another standard measurement for DILI. The severity of liver injury was also confirmed by H&E staining in this study. The characteristics of AILI is predominantly hepatocellular necrosis, featured as the abruption of the cell membranes and nuclei of hepatocytes around the central veins. Liver samples from the PBS control and 400 mg/kg APAP treated groups with the highest ALT or AST levels were selected for staining. Representative images of the control and APAP groups are shown in Fig. 1C. No liver injury was observed in any of the control groups of any age (top row). In the APAP-treated groups (bottom row), no necrosis could be observed in the infant male mice at day 10, but ballooning of hepatocytes was found in the child group at day 22 and large injured areas around the central veins were seen in the adult group at day 60. The severity of the liver injury was scored by the Ishak system based on H&E staining following the described method in Table 2. Consistent with the results in ALT or AST levels, the treatment with 400 mg/kg APAP led to the most severe injury in male mice at day 60 among the three age groups (Fig. 1C). In detail, for the infant mice at day 10, the score in the APAP-treated group was 0, which means no injury happened. For the child mice at day 22, the Ishak score was 2, suggesting moderate AILI. For

DMD # 89557

the adult mice at day 60, APAP-treated liver had an Ishak score of 5, suggesting extremely severe AILI. The *in vitro* metabolism rate of APAP by liver microsomes from the mice at day 10 was significantly lower than that of the mice at day 60, indicating that the differential severity of AILI between the ages was partly due to age-related differential metabolism of APAP, which was mainly mediated by CYP1A2, 2E1, and 3A11 (Supplemental Fig. S1).

Overall, older male mice were more susceptible to AILI than younger ones. The severity of AILI was also dose-dependent.

Effects of AILI on Expression of CYP1A2 in Mouse Liver at Different Ages. Among the three age groups, the basal mRNA expression of CYP1A2 was the highest in the mice at day 60 followed by the mice at day 22 (Fig. 2A). After same doses of APAP treatment, the mRNA expression of CYP1A2 was also higher in the mice at day 22 and 60 than the mice at day 10 (Fig. 2A). APAP treatment significantly decreased the mRNA expression of CYP1A2 in all age groups. Compared to the age-matched controls, the mRNA expression of CYP1A2 was eighty-five percent lower in the APAP treatment mice at day 60. Consistent with the most severe AILI at day 60, the extent of repression of CYP1A2 expression was the largest among the three ages with same doses of APAP treatment.

The relative protein levels of CYP1A2 were measured in randomly selected male mouse livers from control (0 mg/kg) and APAP-treated (400 mg/kg) mice at each age (n = 3-4). Consistent with the CYP1A2 mRNA expression, the basal protein expression of CYP1A2 was also higher in older age mice (Fig. 2B). Moreover, a higher protein expression of CYP1A2 was observed in the APAP-treated older age groups compared to the dose-matched mice at day 10. A significant decrease of CYP1A2 protein expression was found in the APAP-treated mice receiving 400 mg/kg at ages of day 22 and 60 in comparison to age-matched controls (Fig. 2B). The changes of CYP1A2 protein were directly related to the APAP-treatment because the total

DMD # 89557

amount of proteins per gram liver tissue (Supplemental Fig. S2A) and microsomal recovery (Supplemental Fig. S2B) were unchanged and expression of the housekeeping protein GAPDH was unaffected by APAP treatment (Supplemental Fig. S2C).

The enzyme activity of CYP1A2 was determined by a UPLC-QTOFMS method using the formation rates of RSF from 7-ethoxy-RSF in the liver S9 samples of four randomly selected mice from the control and 400 mg/kg APAP-treated groups. The basal activity of CYP1A2 was higher in the livers of mice at day 20 and 60 compared to the mice at day 10 (Fig. 2C). No differences in the enzyme activity of CYP1A2 were found between control and APAP-treated groups in the mice at day 10. However, after the APAP treatment, the mice at day 60 and day 22 had lower CYP1A2 enzyme activity than the age-matched control groups (Fig. 2C).

Overall, APAP treatment decreased the expression of CYP1A2 at mRNA and protein levels as well as enzyme activity in the liver of mice at different ages.

Effects of ALI on Expression and Activity of CYP2B10 in Mouse Liver at Different Ages. As seen in Fig. 3A, the basal mRNA expression of CYP2B10 was lower in the mice at day 22, but higher in the mice at day 60 compared to the mice at day 10. Differential effects of APAP treatment on the expression of CYP2B10 at the three age groups were observed. Mice at day 10 receiving higher doses of APAP treatment (300 and 400 mg/kg) exhibited decreased mRNA expression of CYP2B10 in liver. Whereas, increased mRNA expression of CYP2B10 by APAP was seen in the liver of mice at day 22 and 60.

The relative protein levels of CYP2B10 were measured in randomly selected male mouse livers from control (0 mg/kg) and APAP-treated mice (400 mg/kg) at each age (n = 3-4). Inconsistent with CYP2B10 mRNA expression, the basal protein expression of CYP2B10 was significantly decreased in the mice at day 60 compared to the mice at day 10 (Fig. 3B). Different effects of APAP treatment on the expression of CYP2B10 were found between mRNA and

DMD # 89557

protein levels. The mice at day 22 treated with APAP had lower protein expression of CYP2B10 while mRNA levels were higher (Fig.3). Comparable protein levels of CYP2B10 were observed in the mice at day 10 and 60 between APAP-treated and age-matched controls (Fig. 3B).

The enzyme activity of CYP2B10 was determined by an UPLC-QTOFMS based method using the formation rates of 8-OH-EFV in liver S9 samples from the control and APAP-treated mice at 400 mg/kg (n = 4). The basal activity of CYP2B10 was higher in liver of the mice at day 22 and 60 compared to the mice at day 10 (Fig. 3C). In contrast to the effect of APAP on CYP2B10 mRNA expression, the activity of CYP2B10 was significantly decreased by APAP in the mice at day 60 (Fig. 3C). No statistical differences were seen between the control and APAP-treated mice at day 10 and 22.

Overall, APAP treatment of mice had a suppression or induction effect on the expression of CYP2B10 that is age-dependent. There is a discrepancy on the impact of APAP on CYP2B10 expression at levels of mRNA, protein, and enzyme activity.

Effects of ALI on Expression and Activity of CYP2C29 in Mouse Liver at Different Ages. As seen in Fig. 4, the expression of CYP2C29 was measured in male mice at three ages and different doses of APAP. At the ages of day 22 and 60, the basal mRNA expression of CYP2C29 was 8- and 4-fold, respectively, higher than that in the mice at day 10 (Fig. 4A). Treatment with APAP decreased the mRNA expression of CYP2C29 at each age (Fig. 4A). In the mice at day 10, mRNA level of CYP2C29 was significantly decreased in the APAP-treated mice at all three doses in comparison to the controls. For the mice at day 22, mRNA level of CYP2C29 was repressed in the mice treated with 400 mg/kg APAP compared to aged-matched controls. Although the mRNA expression of CYP2C29 was decreased in a dose-dependent manner in the mice at day 22, that was still 3-fold higher after APAP treatment than the control mice at day 10 (Fig. 4A).

DMD # 89557

The relative protein levels of CYP2C29 were measured in randomly selected male mouse livers from control (0 mg/kg) and APAP-treated mice (400 mg/kg) at each age (n = 3-4). Consistent with the basal CYP2C29 mRNA expression, the protein expression of CYP2C29 was increased dramatically at day 22 and kept steady at day 60 compared to the mice at day 10 (Fig. 4B). APAP treatment also significantly decreased the protein expression of CYP2C29 in the mice at day 22 and 60. Although the expression of CYP2C29 at protein level was decreased by APAP in the mice at day 22 and 60, they were still much higher than the control mice at day 10 (Fig. 4B).

The enzyme activity of CYP2C29 was determined by an UPLC-QTOFMS method using the formation rates of demethyl-RSG in liver S9 samples from the control and APAP-treated mice at 400 mg/kg (n = 4). Consistent with the basal expression of CYP2C29 at mRNA and protein levels, the enzyme activity of CYP2C29 was increased in the mice at day 22 compared to the mice at day 10 (Fig. 4C). After APAP treatment, the mice at day 60 had a lower enzyme activity of CYP2C29 than the age-matched controls (Fig. 4C).

Overall, APAP treatment resulted in decreased expression of CYP2C29 at levels of mRNA, protein, and enzyme activity at each age.

Effects of ALLI on Expression and Activity of CYP2E1 in Mouse Liver at Different Ages.

The expression of CYP2E1 at the levels of mRNA, protein, and enzyme activity was also measured in this study. For the basal mRNA expression of CYP2E1, slightly higher expression was found in the mice at day 22 and a dramatically elevated level was seen in the mice at day 60 with 4-fold higher than the mice at day 10 (Fig. 5A). Generally, APAP treatment decreased the mRNA expression of CYP2E1 at all ages. The mice at day 10 treated with 400 mg/kg APAP had a significantly lower mRNA expression of CYP2E1, while only lower doses of APAP treatment (200 and 300 mg/kg) decreased the mRNA expression of CYP2E1 in the mice at day

DMD # 89557

60 (Fig. 5A). However, APAP treatment at each dose repressed the mRNA expression of CYP2E1 in the mice at day 22 compared to the age-matched controls (Fig. 5A).

The relative protein levels of CYP2E1 were measured in control (0 mg/kg) and APAP-treated (400 mg/kg) mice at each age (n = 3-4). Similar to the basal mRNA expression of CYP2E1, the protein levels were also higher in the mice at day 22 and 60 than day 10 (Fig. 5B). Consistent with the decreased mRNA expression of CYP2E1 by APAP treatment, reduced protein expression of CYP2E1 was also observed in the mice at day 22 (Fig. 5B). Moreover, significantly lower protein expression of CYP2E1 was found in the mice at day 60 treated with APAP compared to the age-matched controls (Fig. 5B).

The enzyme activity of CYP2E1 was determined by an UPLC-QTOFMS method using the formation rates of 6-OH-CHZ in liver S9 samples from the control and APAP-treated mice at 400 mg/kg (n = 4). Consistent with the basal mRNA and protein expression of CYP2E1, the enzyme activity was also higher in the mice at day 22 and 60 compared to the mice at day 10 (Fig. 5C). A trend of decrease of enzyme activity of CYP2E1 was found in the mice at day 22 and 60 treated with 400 mg/kg APAP compared to age-matched controls (Fig. 5C).

Overall, APAP treatment reduced the expression of CYP2E1 at the levels of mRNA, protein, and enzyme activity at each age.

Effects of ALI on Expression and Activity of CYP3A11 in Mouse Liver at Different Ages. The basal expression of CYP3A11 at mRNA level was higher in the older mice at day 22 and 60 with a 2- and 8-fold increase, respectively, than the mice at day 10 (Fig. 6A). This is reflective to a normal ontogenic expression pattern for CYP3A11 during postnatal liver maturation. In general, APAP treatment resulted in decreased mRNA expression of CYP3A11, which was correlated with the severity degree of tissue injury. In the mice at day 10, CYP3A11 mRNA expression was decreased only in the 400 mg/kg APAP-treated mice (Fig. 6A). In the

DMD # 89557

mice at day 22, CYP3A11 mRNA levels were decreased in all APAP-treated mice compared to the controls, but by less than 2-fold. In the mice at day 60, CYP3A11 mRNA decreased more than 2-fold in all APAP-treated mice compared to the controls, which is correlated to the AILI severity.

The relative protein levels of CYP3A11 were measured in control (0 mg/kg) and APAP-treated (400 mg/kg) mice at each age (n = 3-4). Consistent with higher basal mRNA expression of CYP3A11 in the mice at day 22 and 60, the protein expression of CYP3A11 was also higher in the mice at day 22 and 60 compared to day 10 (Fig. 6B). Similar to the effect of APAP on the mRNA expression of CYP3A11, the APAP treatment also resulted in a significantly decreased protein expression of CYP3A11 in the mice at day 22 and 60. Especially, the protein level of CYP3A11 in the mice at day 60 with APAP treatment was decreased to a similar level as the controls at day 10 (Fig. 6B).

The enzyme activity of CYP311 was determined by an UPLC-QTOFMS method using the formation rates of 1-OH-MDZ in liver S9 samples from the control and APAP-treated mice at 400 mg/kg (n = 4). Consistent with the mRNA and protein expression of CYP3A11, the basal enzyme activity of CYP3A11 was increased with age and decreased by APAP treatment (Fig. 6C).

Overall, APAP treatment reduced the expression of CYP3A11 at the levels of mRNA, protein, and enzyme activity at each age in a dose-dependent manner.

Impact on APAP Treatment in Mice by Gender. We then investigated whether gender affected the severity of AILI and the consequence of AILI on the expression of P450s. In this study, both male and female mice at day 60 were used and the mRNA expression of CYP3A11 was determined.

DMD # 89557

As seen in Fig. 7, APAP treatment significantly increased the levels of ALT and AST in both genders, indicating that AILI occurred. After treated with same doses of APAP, alteration magnitudes of ALT and AST changes were greater in the males than females. Consistent with grading of liver injury based on ALT or AST levels, H&E staining results showed more severe injury induced by 400 mg/kg APAP in male mice than in female mice (Fig. 7C).

Comparable basal mRNA levels of CYP3A11 were observed between male and female mice at day 60 (Fig. 7D). Accompanied with AILI, the hepatic mRNA expression of CYP3A11 was decreased in both males and females. Same doses of APAP treatment resulted in lower mRNA expression in males than in females. After same doses of APAP treatment, the mRNA expression of CYP3A11 in the male mice, which had more severe liver injury, was decreased even more than that in female mice.

Overall, APAP treatment led to severe liver injury and lower mRNA expression of CYP3A11 in male more than female mice.

Discussion

The current study confirmed the role of ontogeny of P450s in the severity of AILI at different ages and investigated the age-dependent differences on the consequences of AILI on the hepatic expression and activities of P450s in mice. In this study, the mice at day 10, which had lower basal expression levels of P450s, were observed to be less susceptible to AILI in comparison to the mice at day 22 and 60. CYP1A2, 2B10, 2C29, 2E1, and 3A11 were selected because they are the major P450s in mice correlated to the human P450s responsible for drug metabolism. Decreased expression levels of P450s (except CYP2B10) by AILI were also observed in mice at different ages.

DMD # 89557

Susceptibility to APAP toxicity has been studied in various species (McGill et al., 2012; Jaeschke et al., 2014). Mice are more vulnerable to AILI compared with rats (McGill et al., 2012). Importantly, mice are identified as a more clinically relevant model of AILI than rats due to similar mechanisms of toxicity, including severe mitochondrial dysfunction accompanied by high oxidative stress, in mice and humans (Jaeschke et al., 2014). In addition, cell models are not suitable to study age-related effects. Thus, it is reasonable to use the mouse model in the present study. In this study, mice at day 10, 22, and 60 were selected to mimic the stages of infant, child, and adult in humans, respectively, based on our previous studies related to the ontogenic expression of P450s (Hart et al., 2009; Peng et al., 2012).

Intensity of liver injury is dose-dependent (Jaeschke, 2015). Generally, the doses of 200 to 500 mg/kg APAP are used to treat adult mice to cause AILI (Harrill et al., 2009; Imaeda et al., 2009). In this study, three doses of APAP (200, 300, and 400 mg/kg) were selected based on previous studies. The results of ALT and AST levels in plasma as well as H&E staining confirmed an injury of liver in adult mice as expected. Mice at younger ages were reported to be more resistant to AILI (Adamson et al., 1991). In the present study, the resistance to AILI in the mice at the infant age was further confirmed. For details, the treatment with APAP at 200, 300, and 400 mg/kg did not induce liver injury in the mice at day 10, while same doses of APAP caused mild to severe liver injury in the mice at day 22 and severe to life threatening liver injury in the mice at day 60. Consistent with literature, lower expression levels of selected P450s were observed in the liver of mice at younger ages in the current study. The resistance to AILI in the mice at day 10 is partly due to the lower activities of CYP1A2, 2E1, and 3A11, which are responsible to metabolism of APAP to a toxic metabolite NAPQI. The differences in the capability of metabolism of APAP in the mice between day 10 and 60 have been confirmed by

DMD # 89557

a lower rate of *in vitro* metabolism of APAP by microsomal proteins from the mice at day 10 compared to the mice at day 60 (Supplemental Fig. S1).

Previous studies have reported altered expression of P450s by APAP treatment at subtoxic doses and overdoses (Gill et al., 2017; Jeong et al., 2019; Ohtsuki et al., 2019). However, contradictory results were observed in these studies. It is also unclear the influence of APAP treatment at infant age on the expression of P450s. In HepaRG cells, a high concentration of APAP was found to down-regulate the mRNA expression of CYP1A2, 2E1, and 3A4 (Gill et al., 2017). Whereas, *in vivo* studies in mice treated with 300 mg/kg APAP increase the expression of CYP1A2 (Chowdhary et al., 2017). In the present study, lower expression levels of CYP1A2, 2C29, 2E1, and 3A11 were seen in the mice with AILI at day 22 and 60, which is consistent with the previous *in vitro* study. A subtoxic dose of APAP was found to elevate the expression of hepatic CYP2E1 and CYP3A proteins in adult rats (Kim et al., 2007). However, in the current study, APAP-treated mice at day 10 had decreased expression of all selected P450s though no liver injury was evident. This may be due to different species as well as different ages of rodents utilized. In the previous study, adult rats were used to investigate the impact of subtoxic doses of APAP; whereas, infant mice groups were selected as no liver injury was induced by APAP. Importantly, the present results were consistent to the study in HepaRG cells, which is a good *in vitro* model to mimic the APAP treatment in humans (McGill et al., 2011; Xie et al., 2015).

In addition, differences in susceptibility to APAP and its consequence on P450 expression were investigated in different genders in the current study. APAP treatment at three doses elevated the ALT and AST levels in both male and female mice (Fig. 7 A and B). Liver injury was also confirmed by H&E staining and severe injury was seen in all the APAP-treated mice (Fig. 7C). Compared with male mice, lower levels of ALT and AST as well as relatively less

DMD # 89557

severe injury were found in female mice treated with same doses of APAP, indicating male mice are more vulnerable to ALI than female mice. This finding is consistent with previous studies (Du et al., 2014).

Although the mechanisms of ALI have been well established, the underlying mechanism of altered expression of P450s by APAP treatment has not been elucidated. Events involved in the regulation of P450s may play roles in the altered expression of P450s by APAP. For instance, necrosis causing cell mass loss may be responsible for the decrease of P450 expression. In ALI, necrosis mainly happened around a central vein area of the liver, where P450s are primarily expressed (Kietzmann, 2017; Kitaoka et al., 2018). Further immunostaining experiment may be required to test the zonation changes of P450s with APAP treatment. In addition, endoplasmic reticulum stress (Nagy et al., 2007), which activates gene expression in ALI, represses the expression of CYP3A4 in HepG2 cells (Vachirayonsti et al., 2015). Treatment with IL-6 has a similar effect, indicating the potential relationship between inflammation and P450 expression. Further knock-in and knock-down experiments may need to investigate the signaling pathways involved. Recent research demonstrated that miRNAs may contribute to the altered expression of P450s in ALI. Elevated expression of miRNAs, eg. miR-122-5p and miR-378a-5p, in ALI contributed to altered expression of CYP1A2, 2E1, and 3A4 in HepRG2 cells (Gill et al., 2017). Knock-down of long non-coding RNAs (lncRNAs) HNF1a-AS1 and HNF4a-AS1 alters susceptibility of APAP-induced cytotoxicity in HepaRG cells (Chen et al., 2020). The roles of noncoding RNAs (including miRNAs and lncRNAs) in the alterations of P450 expression by APAP should be investigated in future studies. In addition to APAP, many other drugs can result in DILI by various mechanisms. Impact on P450 expression by DILI should be studied with more drugs than APAP.

DMD # 89557

Currently, there is no discussion in clinical setting on how to handle co-administered drugs in patients who experience ALI. Very likely, current practice is to stop all drugs when ALI happens. However, such decision is not based on sound science, instead by a logical concern by clinicians that patients with liver injury may have an altered capacity to metabolize drugs. Many ALI patients may have chronic conditions, such as diabetes, hypertension, hyperlipidemia, and depression under controls by multiple medications. P450s play an important role in metabolism of drugs to treat many chronic diseases. Management of the diseases requires a long period of treatment with medications, even lifetime. Understanding how expression and activities of the P450s are altered under an ALI condition is clinically critical. The data in this manuscript may stimulate more studies on how to make evidence-based decisions on doses of drugs in management of various conditions in patients who experience ALI.

In conclusion, the present study indicates that adult mice are more vulnerable to ALI compared with infant mice, which may be due to the higher expression levels of P450s at the adult age. It is also demonstrated that ALI results in reduced expression of P450s (except CYP2B10) in mice, which may influence the metabolism and toxicity of drugs that are substrates for these enzymes when co-administrated with APAP. Moreover, male mice are more susceptible to ALI than female mice. Further studies are needed to explore the underlying mechanisms of reduction of P450s by ALI.

DMD # 89557

Authorship Contributions:

Participated in research design: Bao, Wang, Zhang, Zhu, Ma, Manautou, Zhong.

Conducted experiments: Bao, Wang, Shao, Xiao, Shi, Zhu.

Performed data analysis: Bao, Wang, Zhong.

Wrote or contributed to the writing of the manuscript: Bao, Wang, Shao, Zhang, Zhu, Ma,
Manautou, Zhong

DMD # 89557

References

- Adamson GM, Papadimitriou JM, and Harman AW (1991) Postnatal mice have low susceptibility to paracetamol toxicity. *Pediatric research* **29**:496.
- Bell LN and Chalasani N (2009) Epidemiology of idiosyncratic drug-induced liver injury. *Semin Liver Dis* **29**:337-347.
- Brewer CT and Chen T (2017) Hepatotoxicity of Herbal Supplements Mediated by Modulation of Cytochrome P450. *International journal of molecular sciences* **18**.
- Bunchorntavakul C and Reddy KR (2013) Acetaminophen-related hepatotoxicity. *Clin Liver Dis* **17**:587-607, viii.
- Chen L, Wang P, Manautou JEE, and Zhong XB (2020) Knockdown of LncRNAs HNF1alpha-AS1 and HNF4alpha-AS1 Alters Susceptibility of Acetaminophen-induced Cytotoxicity in HepaRG Cells. *Molecular pharmacology*.
- Chowdhary V, Teng KY, Thakral S, Zhang B, Lin CH, Wani N, Bruschiweiler-Li L, Zhang X, James L, Yang D, Junge N, Bruschiweiler R, Lee WM, and Ghoshal K (2017) miRNA-122 Protects Mice and Human Hepatocytes from Acetaminophen Toxicity by Regulating Cytochrome P450 Family 1 Subfamily A Member 2 and Family 2 Subfamily E Member 1 Expression. *The American journal of pathology* **187**:2758-2774.
- Corsini A and Bortolini M (2013) Drug-induced liver injury: the role of drug metabolism and transport. *J Clin Pharmacol* **53**:463-474.
- Dahlin DC, Miwa GT, Lu AY, and Nelson SD (1984) N-acetyl-p-benzoquinone imine: a cytochrome P-450-mediated oxidation product of acetaminophen. *Proc Natl Acad Sci U S A* **81**:1327-1331.
- Dai G, He L, Chou N, and Wan Y-JY (2006) Acetaminophen Metabolism Does Not Contribute to Gender Difference in Its Hepatotoxicity in Mouse. *Toxicological Sciences* **92**:33-41.
- Du K, Williams CD, McGill MR, and Jaeschke H (2014) Lower susceptibility of female mice to acetaminophen hepatotoxicity: Role of mitochondrial glutathione, oxidant stress and c-jun N-terminal kinase. *Toxicology and applied pharmacology* **281**:58-66.

DMD # 89557

Fernández I, Peña A, Del Teso N, Pérez V, and Rodríguez-Cuesta J (2010) Clinical biochemistry parameters in C57BL/6J mice after blood collection from the submandibular vein and retroorbital plexus. *Journal of the American Association for Laboratory Animal Science : JAALAS* **49**:202-206.

Fontana RJ (2008) Acute liver failure including acetaminophen overdose. *Med Clin North Am* **92**:761-794, viii.

Gavito AL, Cabello R, Suarez J, Serrano A, Pavon FJ, Vida M, Romero M, Pardo V, Bautista D, Arrabal S, Decara J, Cuesta AL, Valverde AM, Rodriguez de Fonseca F, and Baixeras E (2016) Single administration of recombinant IL-6 restores the gene expression of lipogenic enzymes in liver of fasting IL-6-deficient mice. *Br J Pharmacol* **173**:1070-1084.

Gill P, Bhattacharyya S, McCullough S, Letzig L, Mishra PJ, Luo C, Dweep H, and James L (2017) MicroRNA regulation of CYP 1A2, CYP3A4 and CYP2E1 expression in acetaminophen toxicity. *Scientific reports* **7**:12331.

Goodman ZD (2007) Grading and staging systems for inflammation and fibrosis in chronic liver diseases. *Journal of hepatology* **47**:598-607.

Guo GL, Moffit JS, Nicol CJ, Ward JM, Aleksunes LA, Slitt AL, Kliewer SA, Manautou JE, and Gonzalez FJ (2004) Enhanced acetaminophen toxicity by activation of the pregnane X receptor. *Toxicological sciences : an official journal of the Society of Toxicology* **82**:374-380.

Harrill AH, Watkins PB, Su S, Ross PK, Harbourt DE, Stylianou IM, Boorman GA, Russo MW, Sackler RS, Harris SC, Smith PC, Tennant R, Bogue M, Paigen K, Harris C, Contractor T, Wiltshire T, Rusyn I, and Threadgill DW (2009) Mouse population-guided resequencing reveals that variants in CD44 contribute to acetaminophen-induced liver injury in humans. *Genome research* **19**:1507-1515.

Hart SN, Cui Y, Klaassen CD, and Zhong XB (2009) Three patterns of cytochrome P450 gene expression during liver maturation in mice. *Drug Metab Dispos* **37**:116-121.

Hassan HM, Yousef BA, Guo H, Xiaoxin L, Zhang L, and Jiang Z (2018) Investigating the CYP2E1 Potential Role in the Mechanisms Behind INH/LPS-Induced Hepatotoxicity. *Frontiers in pharmacology* **9**:198.

DMD # 89557

Hoofnagle JH, Serrano J, Knoblen JE, and Navarro VJ (2013) LiverTox: a website on drug-induced liver injury.

Hepatology **57**:873-874.

Imaeda AB, Watanabe A, Sohail MA, Mahmood S, Mohamadnejad M, Sutterwala FS, Flavell RA, and Mehal WZ

(2009) Acetaminophen-induced hepatotoxicity in mice is dependent on Tlr9 and the Nalp3 inflammasome. *The Journal of clinical investigation* **119**:305-314.

Jaeschke H (2015) Acetaminophen: Dose-Dependent Drug Hepatotoxicity and Acute Liver Failure in Patients.

Digestive diseases **33**:464-471.

Jaeschke H, Xie Y, and McGill MR (2014) Acetaminophen-induced Liver Injury: from Animal Models to Humans.

J Clin Transl Hepatol **2**:153-161.

Jeong TB, Kim J-H, Kim SH, Lee S, Son SW, Lim Y, Cho J-Y, Hwang DY, Kim KS, Kwak J-H, and Jung Y-S (2019)

Comparison of toxic responses to acetaminophen challenge in ICR mice originating from different sources. *Laboratory Animal Research* **35**.

Kietzmann T (2017) Metabolic zonation of the liver: The oxygen gradient revisited. *Redox Biology* **11**:622-630.

Kim SN, Seo JY, Jung DW, Lee MY, Jung YS, and Kim YC (2007) Induction of hepatic CYP2E1 by a subtoxic dose of acetaminophen in rats: increase in dichloromethane metabolism and carboxyhemoglobin elevation.

Drug metabolism and disposition: the biological fate of chemicals **35**:1754-1758.

Kitaoka S, Hatogai J, Ochiai W, and Sugiyama K (2018) Zonation of the drug-metabolizing enzyme cytochrome

P450 3A in infant mice begins in pre-weaning period. *J Toxicol Sci* **43**:223-227.

Larrey D (2002) Epidemiology and individual susceptibility to adverse drug reactions affecting the liver. *Seminars*

in liver disease **22**:145-155.

Leise MD, Poterucha JJ, and Talwalkar JA (2014) Drug-induced liver injury. *Mayo Clin Proc* **89**:95-106.

Maddrey WC (2005) Drug-induced hepatotoxicity: 2005. *J Clin Gastroenterol* **39**:S83-89.

McGill MR, Williams CD, Xie Y, Ramachandran A, and Jaeschke H (2012) Acetaminophen-induced liver injury in

rats and mice: comparison of protein adducts, mitochondrial dysfunction, and oxidative stress in the mechanism of toxicity. *Toxicol Appl Pharmacol* **264**:387-394.

DMD # 89557

McGill MR, Yan HM, Ramachandran A, Murray GJ, Rollins DE, and Jaeschke H (2011) HepaRG cells: a human model to study mechanisms of acetaminophen hepatotoxicity. *Hepatology* **53**:974-982.

Nagy G, Kardon T, Wunderlich L, Szarka A, Kiss A, Schaff Z, Banhegyi G, and Mandl J (2007) Acetaminophen induces ER dependent signaling in mouse liver. *Arch Biochem Biophys* **459**:273-279.

Ohtsuki Y, Sanoh S, Santoh M, Ejiri Y, Ohta S, and Kotake Y (2019) Inhibition of cytochrome P450 3A protein degradation and subsequent increase in enzymatic activity through p38 MAPK activation by acetaminophen and salicylate derivatives. *Biochemical and biophysical research communications* **509**:287-293.

Otto GP, Rathkolb B, Oestreich MA, Lengger CJ, Moerth C, Micklich K, Fuchs H, Gailus-Durner V, Wolf E, and Hrabě de Angelis M (2016) Clinical Chemistry Reference Intervals for C57BL/6J, C57BL/6N, and C3HeB/FeJ Mice (*Mus musculus*). *Journal of the American Association for Laboratory Animal Science : JAALAS* **55**:375-386.

Pasanen M, Pellinen P, Stenback F, Juvonen RO, Raunio H, and Pelkonen O (1995) The role of CYP enzymes in cocaine-induced liver damage. *Archives of toxicology* **69**:287-290.

Peng L, Yoo B, Gunewardena SS, Lu H, Klaassen CD, and Zhong XB (2012) RNA sequencing reveals dynamic changes of mRNA abundance of cytochromes P450 and their alternative transcripts during mouse liver development. *Drug Metab Dispos* **40**:1198-1209.

Piekos SC, Chen L, Wang P, Shi J, Yaqoob S, Zhu HJ, Ma X, and Zhong XB (2018) Consequences of Phenytoin Exposure on Hepatic Cytochrome P450 Expression during Postnatal Liver Maturation in Mice. *Drug Metab Dispos* **46**:1241-1250.

Rudraiah S, Rohrer PR, Gurevich I, Goedken MJ, Rasmussen T, Hines RN, and Manautou JE (2014) Tolerance to acetaminophen hepatotoxicity in the mouse model of autoprotection is associated with induction of flavin-containing monooxygenase-3 (FMO3) in hepatocytes. *Toxicol Sci* **141**:263-277.

Sgro C, Clinard F, Ouazir K, Chanay H, Allard C, Guilleminet C, Lenoir C, Lemoine A, and Hillon P (2002) Incidence of drug-induced hepatic injuries: a French population-based study. *Hepatology* **36**:451-455.

DMD # 89557

Shen T, Liu Y, Shang J, Xie Q, Li J, Yan M, Xu J, Niu J, Liu J, Watkins PB, Aithal GP, Andrade RJ, Dou X, Yao L, Lv F, Wang Q, Li Y, Zhou X, Zhang Y, Zong P, Wan B, Zou Z, Yang D, Nie Y, Li D, Wang Y, Han X, Zhuang H, Mao Y, and Chen C (2019) Incidence and Etiology of Drug-Induced Liver Injury in Mainland China. *Gastroenterology*.

Tien YC, Liu K, Pope C, Wang P, Ma X, and Zhong XB (2015) Dose of Phenobarbital and Age of Treatment at Early Life are Two Key Factors for the Persistent Induction of Cytochrome P450 Enzymes in Adult Mouse Liver. *Drug Metab Dispos* **43**:1938-1945.

Vachirayonsti T, Ho KW, Yang D, and Yan B (2015) Suppression of the pregnane X receptor during endoplasmic reticulum stress is achieved by down-regulating hepatocyte nuclear factor-4alpha and up-regulating liver-enriched inhibitory protein. *Toxicological sciences : an official journal of the Society of Toxicology* **144**:382-392.

Vega M, Verma M, Beswick D, Bey S, Hossack J, Merriman N, Shah A, Navarro V, and Drug Induced Liver Injury N (2017) The Incidence of Drug- and Herbal and Dietary Supplement-Induced Liver Injury: Preliminary Findings from Gastroenterologist-Based Surveillance in the Population of the State of Delaware. *Drug Saf* **40**:783-787.

Xie Y, McGill MR, Cook SF, Sharpe MR, Winefield RD, Wilkins DG, Rollins DE, and Jaeschke H (2015) Time course of acetaminophen-protein adducts and acetaminophen metabolites in circulation of overdose patients and in HepaRG cells. *Xenobiotica; the fate of foreign compounds in biological systems* **45**:921-929.

Xie Y, McGill MR, Dorko K, Kumer SC, Schmitt TM, Forster J, and Jaeschke H (2014) Mechanisms of acetaminophen-induced cell death in primary human hepatocytes. *Toxicol Appl Pharmacol* **279**:266-274.

Yoon E, Babar A, Choudhary M, Kutner M, and Prysopoulos N (2016) Acetaminophen-Induced Hepatotoxicity: a Comprehensive Update. *J Clin Transl Hepatol* **4**:131-142.

DMD # 89557

Footnotes

This study was partly supported by the National Institutes of Health (NIH) National Institute of General Medical Sciences [Grant R01GM-118367] (to X.B.Z); the National Institute of Allergy And Infectious Diseases [Grant R01 AI131983] (to X.M); the NIH National Heart, Lung, and Blood Institute [Grant R01HL126969] (to H.J.Z). This study was also partly supported by the Institute for System Genomics at the University of Connecticut (to X.B.Z). Pei Wang was a visiting scholar supported by the China Scholarship Council [Grant 201707040007].

DMD # 89557

Figure Legends

Fig. 1. Impact of APAP treatment on liver injury in male mice at different ages. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Blood samples and liver tissues were collected at 24 hrs after the treatment. (A) Left panel: plasma ALT activities; right panel: grades of liver injury defined based on ALT levels. (B) Left panel: plasma AST activities; right panel: grades of liver injury defined based on AST levels. (C) Left panel: representative images of liver sections stained with H&E from mice treated with 400 mg/kg APAP and age-matched controls (0 mg/kg); right panel: grades of liver injury defined based on Ishak scores. Data are shown as mean \pm SD ($n = 4-6$) and analyzed by one-way ANOVA with the Dunnett's *post-hoc* test. * $p < 0.05$ and *** $p < 0.001$ versus age-matched controls.

Fig. 2. Effects of APAP treatment on the expression of CYP1A2. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Liver tissues were collected at 24 hrs after the treatment. (A) Hepatic mRNA expression of CYP1A2 ($n = 4-6$). The fold changes of relative mRNA expression compared to the control mice at day 10 were calculated using the $2^{-\Delta\Delta t}$ method after normalization with GAPDH. (B) The protein expression of CYP1A2 in liver microsomes from the mice treated with 0 or 400 mg/kg APAP ($n = 3-4$). The relative protein expression of CYP1A2 was determined by LC-MS/MS with normalization to external added BSA. (C) Enzyme activity of CYP1A2 in liver S9 fractions of mice treated with 0 or 400 mg/kg APAP ($n = 4$). Formation rates of 7-ethoxy-RSF to RSF were determined by UPLC-QTOFMS. Data were presented as mean \pm SD. Statistical significance was determined by one-way ANOVA with the Dunnett's *post-hoc* test. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ versus age-matched controls; ### $p < 0.001$ versus 0 mg/kg APAP treatment at day 10.

DMD # 89557

Fig. 3. Effects of APAP treatment on the expression and activity of CYP2B10. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Liver tissues were collected at 24 hrs after the treatment. (A) Hepatic mRNA expression of CYP2B10 (n = 4-6). The fold changes of relative mRNA expression compared to the control mice at day 10 were calculated using the $2^{-\Delta\Delta t}$ method after normalization with GAPDH. (B) The protein expression of CYP2B10 in liver microsomes from the mice treated with 0 or 400 mg/kg APAP (n = 3-4). The relative protein expression of CYP2B10 was determined by LC-MS/MS with normalization to external added BSA. (C) Enzyme activity of CYP2B10 in liver S9 fractions of mice treated with 0 or 400 mg/kg APAP (n = 4). Formation rates of 8-OH-EFV were determined by UPLC-QTOFMS. Data were presented as mean \pm SD. Statistical significance was determined by one-way ANOVA with the Dunnett's post-hoc test. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ versus age-matched controls; # $p < 0.05$, ## $p < 0.01$, and ### $p < 0.001$ versus 0 mg/kg APAP treatment at day 10.

Fig. 4. Effects of APAP treatment on the expression and activity of CYP2C29. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Liver tissues were collected at 24 hrs after the treatment. (A) Hepatic mRNA expression of CYP2C29 (n = 4-6). The fold changes of relative mRNA expression compared to the control mice at day 10 were calculated using the $2^{-\Delta\Delta t}$ method after normalization with GAPDH. (B) The protein expression of CYP2C29 in liver microsomes from the mice treated with 0 or 400 mg/kg APAP (n = 3-4). The relative protein expression of CYP2C29 was determined by LC-MS/MS with normalization to external added BSA. (C) Enzyme activity of CYP2C29 in liver S9 fractions of mice treated with 0 or 400 mg/kg APAP (n

DMD # 89557

= 4). Formation rates of demethyl-RSF were determined by UPLC-QTOFMS. Data were presented as mean \pm SD. Statistical significance was determined by one-way ANOVA with the Dunnett's post-hoc test. * p <0.05, ** p <0.01, and *** p <0.001 *versus* age-matched controls; # p <0.05 and ### p <0.001 *versus* 0 mg/kg APAP treatment at day 10.

Fig. 5. Effects of APAP treatment on the expression and activity of CYP2E1. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Liver tissues were collected at 24 hrs after the treatment. (A) Hepatic mRNA expression of CYP2E1 (n = 4-6). The fold changes of relative mRNA expression compared to the control mice at day 10 were calculated using the $2^{-\Delta\Delta t}$ method after normalization with GAPDH. (B) The protein expression of CYP2E1 in liver microsomes from the mice treated with 0 or 400 mg/kg APAP (n = 3-4). The relative protein expression of CYP2E1 was determined by LC-MS/MS with normalization to external added BSA. (C) Enzyme activity of CYP2E1 in liver S9 fractions of mice treated with 0 or 400 mg/kg APAP (n = 4). Formation rates of 6-OH-CHZ were determined by UPLC-QTOFMS. Data were presented as mean \pm SD. Statistical significance was determined by one-way ANOVA with the Dunnett's post-hoc test. * p <0.05, ** p <0.01, and *** p <0.001 *versus* age-matched controls; # p <0.05, ## p <0.01, and ### p <0.001 *versus* 0 mg/kg APAP treatment at day 10.

Fig. 6. Effects of APAP treatment on the expression and activity of CYP3A11. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at a series of doses (0, 200, 300, or 400 mg/kg). Liver tissues were collected at 24 hrs after the treatment. (A) Hepatic mRNA expression of CYP3A11 (n = 4-6). The fold changes of relative mRNA expression compared to the control mice at day 10 were calculated using the $2^{-\Delta\Delta t}$ method after

DMD # 89557

normalization with GAPDH. (B) The protein expression of CYP3A11 in liver microsomes from the mice treated with 0 or 400 mg/kg APAP (n = 3-4). The relative protein expression of CYP3A11 was determined by LC-MS/MS with normalization to external added BSA. (C) Enzyme activity of CYP3A11 in liver S9 fractions of mice treated with 0 or 400 mg/kg APAP (n = 4). Formation rates of 1-OH-MDZ were determined by UPLC-QTOFMS. Data were presented as mean \pm SD. * $p < 0.05$ and *** $p < 0.001$ versus age-matched controls; ## $p < 0.01$ and ### $p < 0.001$ versus 0 mg/kg APAP treatment at day 10.

Fig. 7. Gender differential impacts of APAP treatment on mouse liver. Both male and female mice at day 60 after birth were treated with different doses of APAP (0, 200, 300, or 400 mg/kg). Blood and liver tissues were collected at 24 hrs after the treatment. (A) Left panel: plasma ALT activities; right panel: grades of liver injury defined based on ALT levels. (B) Left panel: plasma AST activities; right panel: grades of liver injury defined based on AST levels. (C) Upper panel: representative images of liver sections stained with H&E from mice treated with 400 mg/kg APAP and gender-matched controls (0 mg/kg); lower panel: grades of liver injury defined based on Ishak scores. (D). The fold changes of relative CYP3A11 mRNA expression compared to the gender-matched controls calculated using the $2^{-\Delta\Delta t}$ method after normalization with GAPDH. Data were presented as mean \pm SD. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ versus gender-matched controls.

DMD # 89557

Table 1. Grading for liver injury

	Grade 0	Grade 1	Grade 2	Grade 3	Grade 4
ALT level*					
Fold (× URL)	<1.25	1.25-2.5	2.6-5.0	5.1-10.0	>10.0
Male (U/L)	<117	117-235	236-470	471-940	>940
Female (U/L)	<110	110-220	221-440	441-880	>880
AST level*					
Fold (× URL)	<1.25	1.25-2.5	2.6-5.0	5.1-10.0	>10.0
Male (U/L)	<152	152-305	306-610	611-1220	>1220
Female (U/L)	<220	220-440	441-880	881-1760	>1760

Footnote: *Value of lower limit was included, while value of upper limit was not included.

URL, upper reference limit.


DMD # 89557

Table 2. Ishak evaluation system

Periportal or periseptal injury		Confluent necrosis		Focal (spotty) lytic injury		Portal inflammation	
Absent	0	Absent	0	Absent	0	None	0
Mild	1	Focal confluent necrosis	1	One focus or less per 10x objective	1	Mild, some or all portal areas	1
Mild/moderate	2	Zone 3 necrosis in some areas	2	Two to four foci per 10x objective	2	Moderate, some or all portal areas	2
Moderate	3	Zone 3 necrosis in most areas	3	Five to ten foci per 10x objective	3	Moderate/marked, all portal areas	3
Severe	4	Zone 3 necrosis + occasional portal-central (P-C) bridging	4	More than ten foci per 10x objective	4	Marked, all portal areas	4
Severe		Zone 3 necrosis + multiple P-C bridging	5				
Severe		Panacinar or multiacinar necrosis	6				


Downloaded from dmd.aspetjournals.org at ASPET Journals on April 19, 2024

Fig. 1




Grading of liver injury based on ALT levels

APAP (mg/kg)	0	200	300	400
Day 10	0	0	0	0
Day 22	0	2	3	3
Day 60	0	3	4	4



Grading of liver injury based on AST levels

APAP (mg/kg)	0	200	300	400
Day 10	0	0	0	1
Day 22	0	2	3	3
Day 60	0	3	3	4



Grading of liver injury based on Ishak scores

Age	Day 10	Day 22	Day 60
APAP 0 mg/kg	0	0	0
APAP 400 mg/kg	0	2	5

Fig. 2

CYP1A2




Fig. 3

CYP2B10




Fig. 4

CYP2C29






Fig. 5

CYP2E1






Fig. 6

CYP3A11






Fig. 7




Grading of liver injury based on ALT levels

APAP (mg/kg)	0	200	300	400
Male	0	2	3	4
Female	0	2	2	3



Grading of liver injury based on AST levels


APAP (mg/kg)	0	200	300	400
Male	0	2	3	4
Female	0	2	2	2



Grading of liver injury based on Ishak scores


Gender	Male	Female
APAP 0 mg/kg	0	0
APAP 400 mg/kg	5	4

D




Yifan Bao, Pei Wang, Xueyan Shao, Junjie Zhu, Jingcheng Xiao, Jian Shi, Lirong Zhang, Hao-Jie Zhu, Xiaochao Ma, José E. Manautou, and Xiao-bo Zhong

Acetaminophen-Induced Liver Injury Alters Expression and Activities of Cytochrome P450 Enzymes in an Age-Dependent Manner in Mouse Liver



Supplemental Fig. S1. Metabolism rate of APAP by liver microsomes isolated from mouse livers at day 10 and 60. Metabolism rate of APAP was determined by HPLC-UV (n = 3). Data were presented as mean \pm SD, ** p <0.01.



Supplemental Fig. S2. Effects of APAP-treatment on the quantification of proteins. Male mice at different ages (postnatal day 10, 22, or 60) were treated with APAP at 0 or 400 mg/kg. Liver tissues were collected at 24 hrs after the treatment. **(A)** Total amount of proteins per gram liver with 0 or 400 mg/kg APAP treatment (n = 3). Total amount of proteins was determined by a Qubit Protein Assay Kit. **(B)** Microsomal proteins in the mice with 0 or 400 mg/kg APAP treatment (n = 3). Amount of microsomal proteins per gram liver was determined by a Qubit Protein Assay Kit. **(C)** The hepatic expression of GAPDH protein in the mice with 0 or 400 mg/kg APAP treatment (n = 3-4). The protein expression of GAPDH was determined by LCT-MS with normalization to external added BSA. Data were presented as mean \pm SD.