

In Vitro–In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicted Drug Disposition[□]

Matthew D. Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R. Russell, Gordon Carlson, Geoffrey Warhurst, and Amin Rostami-Hodjegan

Gut Barrier Group, Inflammation and Repair, University of Manchester, Salford Royal NHS Trust, Salford, United Kingdom (M.D.H., G.C., G.W.); Simcyp Limited (a Certara company), Blades Enterprise Centre, Sheffield, United Kingdom (M.D.H., S.N., A.R.-H.); and Centre for Applied Pharmacokinetic Research, Manchester Pharmacy School, Stopford Building, Manchester, United Kingdom (B.A., M.R.R., A.R.-H.)

Received October 9, 2015; accepted February 1, 2016

ABSTRACT

Relative expression factors (REFs) are used to scale in vitro transporter kinetic data via in vitro–in vivo extrapolation linked to physiologically based pharmacokinetic (IVIVE-PBPK) models to clinical observations. Primarily two techniques to quantify transporter protein expression are available, immunoblotting and liquid chromatography–tandem mass spectrometry. Literature-collated REFs ranged from 0.4 to 5.1 and 1.1 to 90 for intestinal P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP), respectively. The impact of using human jejunum–Caco-2 REFs for P-gp (REF_{IP-gp}) and BCRP (REF_{IBCRP}), generated from the same samples and using different proteomic methodologies from independent laboratories, on PBPK outcomes was assessed. A 5-fold decrease in REF_{IP-gp} for a single oral dose of digoxin resulted in a 1.19- and 1.31-fold higher plasma area under the curve and C_{max}, respectively. All generated REF_{IP-gp} values

led to simulated digoxin C_{max} values within observed ranges; however, combining kinetic data generated from a different laboratory with the 5-fold lower REF_{IP-gp} could not recover a digoxin–rifampicin drug–drug interaction, emphasizing the necessity to obtain transporter-specific kinetic estimates and REFs from the same in vitro system. For a theoretical BCRP compound, with absorption taking place primarily in the jejunum, a decrease in the REF_{IBCRP} from 2.22 (University of Manchester) to 1.11 (Bertin Pharma) promoted proximal intestinal absorption while delaying t_{max} 1.44-fold. Laboratory-specific differences in REF may lead to different IVIVE-PBPK outcomes. To understand the mechanisms underlying projected pharmacokinetic liabilities, it is important to assess the potential impact of bias on the generation of REFs on an interindividual basis within a target population.

Introduction

In vitro–in vivo extrapolation linked to physiologically based pharmacokinetic (IVIVE-PBPK) models aim to predict profiles of drug disposition dynamically. This is accomplished by incorporating “drug” data, generated in vitro, and physicochemical knowledge together with “systems” data in a population (Rostami-Hodjegan, 2012). Kinetic data [i.e., maximal flux capacity of the transporter protein (J_{max}) and K_m] describing the active transport processes generated from cell systems can also be included in IVIVE-PBPK models. To scale these data to in vivo, human and in vitro system transporter protein expression or activity data are also required in combination with physiologic, demographic, and genetic information (Rostami-Hodjegan, 2012). To date, intestinal transporter IVIVE scaling factors (Neuhoff et al., 2013a) have been generated based on Western blotting, a relative quantitative

technique to quantify transport expression (Troutman and Thakker, 2003b). Yet, absolute transporter protein abundances quantified by liquid chromatography–tandem mass spectrometry (LC-MS/MS) have recently been explored for hepatic application in IVIVE-PBPK (Vildhede et al., 2014).

In this study, we provide a systematic analysis of the mRNA and protein expression data available in the literature for generating the relative expression factor (REF), an IVIVE scalar that describes the ratio of in vivo to in vitro systems transporter expression for human jejunum and Caco-2 monolayer P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP). We also evaluate the impact of intestinal P-gp and BCRP REFs generated by different laboratories and methodologies on drug absorption in a PBPK model.

Materials and Methods

Literature Review of the Intestinal Expression Data for P-gp and BCRP. Starting with a previously reported meta-analysis that established human intestinal P-gp and BCRP region-specific protein expression (Harwood et al.,

dx.doi.org/10.1124/dmd.115.067777.

[□]This article has supplemental material available at dmd.aspetjournals.org.

ABBREVIATIONS: BCRP, breast cancer resistance protein; BPh, Bertin Pharma; DDI, drug–drug interaction; HV, healthy Caucasian volunteer population; IVIVE, in vitro–in vivo extrapolation; LC-MS/MS, liquid chromatography–tandem mass spectrometry; PBPK, physiologically based pharmacokinetic; P-gp, P-glycoprotein; REF, relative expression factor; REF_{IBCRP}, intestinal relative expression factor for breast cancer resistance protein; REF_{IP-gp}, intestinal relative expression factor for P-glycoprotein; TC, theoretical BCRP compound; UoM, University of Manchester.

2013), a new search for the relevant published data quantifying P-gp and BCRP mRNA or protein expression in human jejunum and filter-grown Caco-2 monolayers using PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) was undertaken. The following keyword combinations were used in PubMed: human; jejunum; Caco-2; P-gp; MDR1; BCRP; ABCB1; ABCG2; mRNA; protein; expression; absolute; abundance; proteomics. Graphical data were extracted, where required, by GetData Graph Digitizer (<http://getdata-graph-digitizer.com>).

Generation of the Relative Expression Factors. A REF for P-gp or BCRP was generated, where P-gp or BCRP mRNA/protein expression for human jejunum and Caco-2 monolayers was available using the same technique within a laboratory, including reference genes. These were compared with REFs generated from two different LC-MS/MS workflows (two independent laboratories; matching samples) for P-gp and BCRP in human jejunum and 21-day-cultivated Caco-2 monolayers. Methodological details and individual values are provided in the companion study (Harwood et al., 2016).

Incorporating Intestinal REFs into IVIVE-PBPK Models. The impact of the laboratory-specific intestinal relative expression factor for P-glycoprotein (REF_{IP-gp}) and intestinal relative expression factor for breast cancer resistance protein (REF_{IBCRP}) in virtual healthy Caucasian volunteers (HVs) was assessed in a PBPK model (version 14.1; Simcyp, a Certara company, Sheffield, UK) containing the regional distribution of intestinal P-gp and BCRP and their population variability (Harwood et al., 2013). P-gp and BCRP transport in the model is driven by the unbound intracellular enterocyte concentration and is multiplied by REF and the regional-specific transporter expression to yield effective permeability (Yang et al., 2007; Neuhoff et al., 2013a).

The Impact of P-gp and BCRP REF in IVIVE-PBPK. The impact of REF_{IP-gp} values generated by the LC-MS/MS [University of Manchester (UoM), Manchester, UK; Harwood et al., 2016] compared with the immunoblotting approach (Troutman and Thakker, 2003b) on digoxin C_{max} was investigated using identical digoxin parameter inputs as the previously reported digoxin IVIVE-PBPK model (Neuhoff et al., 2013a). Caco-2-derived J_{max} and K_m data (Troutman and Thakker, 2003a) were applied to intestinal and hepatic P-gp, assuming P-gp activity in vitro in healthy individuals corresponds to that in vivo, and that J_{max} is related to P-gp protein expression. Simulations were run with a single oral digoxin dose of 0.5 mg in 100 HV individuals to evaluate if kinetic data for digoxin generated in a Caco-2 system from another laboratory (Troutman and Thakker, 2003a) to the REF_{IP-gp} from the UoM Caco-2 system, could capture the observed digoxin-rifampicin drug-drug interaction (DDI) via induction of intestinal P-gp (Greiner et al., 1999), thus verifying the correct contribution of the active transport built into the digoxin PBPK model.

The impact of laboratory-specific differences for REF_{IBCRP} on pharmacokinetic parameter predictions was evaluated using the Simcyp simulator. A

permeable theoretical BCRP compound (TC); see (Supplemental Materials, Technical Note with limited gut metabolism and a specific BCRP activity was administered orally (10 mg in solution) to 100 HV individuals, with the default region-specific BCRP expression within the PBPK model, as published by Harwood et al. (2013).

Results and Discussion

REF_{IP-gp} and REF_{IBCRP} Generation from Different Laboratories. According to our literature analysis, human jejunum mRNA and protein expression was identified in 19 studies for P-gp and nine studies for BCRP [(Supplemental Materials; (Supplemental Table 1)]. Expression data for Caco-2 P-gp and BCRP from the same laboratory using the same protocol to generate an REF_{IP-gp} or REF_{IBCRP} were found for five and four studies, respectively (Table 1). For P-gp, relative mRNA expression analysis (reverse-transcription polymerase chain reaction) enabled the generation of REF_{IP-gp} from two laboratories in three studies (Taipalensuu et al., 2001; Seithel et al., 2006; Hilgendorf et al., 2007), as the data from Seithel et al. (2006) and Hilgendorf et al. (2007) used Caco-2 monolayers cultivated in the same laboratory for 23 and 16 days, respectively. An REF_{IP-gp} from two independent laboratories that used Western blotting (Troutman and Thakker, 2003b; von Richter et al., 2009) is available, but not for LC-MS/MS quantification for either P-gp or BCRP, as the Caco-2 cell abundances reported by Oswald et al. (2013) were from plastic, not filter-grown cells (Dr. Stefan Oswald, personal communication). The REF_{IP-gp} based on mRNA expression ranged 7.1-fold, and for LC-MS/MS quantification, 2.6-fold. The REF_{IBCRP} of 90 (Taipalensuu et al., 2001) may result from low BCRP levels or variability in the housekeeping gene used (Seithel et al., 2006). The REF_{IP-gp} and REF_{IBCRP} generated from P-gp and BCRP quantification by LC-MS/MS from two different laboratories, UoM and Bertin Pharma (BPh), for the same samples are provided in Table 1 (Harwood et al., 2016). The REF_{IP-gp} (Troutman and Thakker, 2003b) from independent samples quantified by Western blotting was 5-fold higher than the REF_{IP-gp} generated by the UoM (UoM- REF_{IP-gp}), whereas the UoM- REF_{IBCRP} was approximately 2-fold higher than BPh (LC-MS/MS) and Altana AG (Western blot; Wesel, Germany) (von Richter et al., 2009).

Assessing the Sensitivity of REF_{IP-gp} in IVIVE-PBPK. Both the UoM- REF_{IP-gp} of 0.4 (Harwood et al., 2016) and the REF_{IP-gp} of 2

TABLE 1

Generation of REF_{IP-gp} and REF_{IBCRP} from human jejunum and Caco-2 from relative mRNA expression and protein abundance (relative and absolute) data available in the literature for P-gp (MDR1) and BCRP

Transporter and Method	Reference Standard Gene/Protein/Peptide	Jejunum Abundance Mean (S.D., Sample n)	Caco-2 Abundance Mean (S.D., Sample n)	REF	Source
P-gp (MDR1)					
RT-PCR	Villin	7.90 (\pm 1.4, n = 13)	11.0 (n = 1)	0.7	Taipalensuu et al., 2001 ^a
RT-PCR	Cyclophilin-A	0.63 (\pm 0.23, n = 4)	0.13 (n = 1)	5.0	Seithel et al., 2006 ^b
RT-PCR	Cyclophilin-A	1.1 (\pm 0.67, n=5)	0.21 (\pm 0.04, n=3 - 6)	5.1	Hilgendorf et al., 2007 ^b
Western blot	Not run in blot	2.12 (n = 1)	1.04 (\pm 0.16, n = 3)	2.0	Troutman and Thakker, 2003b ^c
Western blot	Not run in blot	1.00 (n = 5, pooled)	1.29 (n = 2)	0.8	Von Richter et al., 2009 ^c
LC-MS/MS	AGAVAEVLAAIR	1.89 (\pm 1.07, n = 3)	4.67 (\pm 0.47, n = 3)	0.4	Harwood et al., 2016 ^d
LC-MS/MS	FYDPLAGK	0.77 (\pm 0.35, n = 3)	2.08 (\pm 0.19, n = 3)	0.4	Harwood et al., 2016 ^e
BCRP					
RT-PCR	Villin	2.7 (\pm 1.4, n = 13)	0.03 (n = 1)	90	Taipalensuu et al., 2001 ^a
RT-PCR	Cyclophilin-A	0.38 (\pm 0.08, n = 4)	0.07 (n = 1)	5.5	Seithel et al., 2006 ^b
RT-PCR	Cyclophilin-A	2.36 (\pm 0.29, n = 5)	0.36 (n = 3 - 6)	6.6	Hilgendorf et al., 2007 ^b
Western blot	Not run in blot	1.00 (n = 5, pooled)	0.84 (n = 2)	1.2	Von Richter et al., 2009 ^c
LC-MS/MS	VIQELGLDK	(2.56 \pm 0.82, n = 3)	(1.16 \pm 0.04, n = 3)	2.2	Harwood et al., 2016 ^d
LC-MS/MS	SLLDVLAAR	(2.06 \pm 1.11, n = 3)	(1.86 \pm 0.14, n = 3)	1.1	Harwood et al., 2016 ^e

RT-PCR, reverse-transcription polymerase chain reaction.

^aUnits for expression of mRNA are number of transcripts per microgram of total RNA.

^bUnits for expression given as relative units of target to reference gene ($2^{-\Delta C_t}$).

^cValues for expression given as signal intensity as measured by densitometry image analysis.

^dUnits for abundance given as fmol protein/ μ g total membrane protein; abundance determined at UoM. The same samples were quantified.

^eUnits for abundance given as fmol protein/ μ g total membrane protein; abundance determined at BPh. The same samples were quantified.

(Troutman and Thakker, 2003b) led to digoxin C_{max} values within observed ranges after a single oral dose of 0.5 mg of digoxin (Fig. 1A), implying both REF_{IP-gp} reflect realistic contributions of P-gp in estimating observed C_{max} values when using the J_{max} and K_m for P-gp reported by Troutman and Thakker (2003a). Using the UoM- REF_{IP-gp} of 0.4 compared with the REF_{IP-gp} of 2 led to a modest 1.31- and 1.19-fold lower mean C_{max} and area under the curve, respectively, in 100 HV individuals (Fig. 1B). A previous study showed that the observed digoxin-rifampicin DDI, which was attributed to a 3.5-fold increase in intestinal P-gp expression (Greiner et al., 1999), could be recovered using an IVIVE-PBPK strategy, in which the REF_{IP-gp} of 2.0 was increased 3.5-fold to 7 after (Fig. 1C) induction (Neuhoff et al., 2013b). A 3.5-fold increase in the UoM- REF_{IP-gp} of 0.4 gave an REF_{IP-gp} of 1.4, leading to a simulated underprediction in the observed DDI (Fig. 1D). This indicates that the lower UoM- REF_{IP-gp} (that is not derived from the same Caco-2 system in which the apparent kinetic data were generated) is not sufficient to recover the contribution of P-gp induction by rifampicin to digoxin plasma concentration in HVs. The inability to recover the observed DDI when using the UoM- REF_{IP-gp} may result from lower P-gp expression and hence lower activity in the UoM Caco-2 systems. This can be due to laboratory differences in methods of expression quantification, Caco-2 cell cultivation, and the variability in jejunum expression. To recover the activity shortfall when using the UoM- REF_{IP-gp} , a 4.3-fold increase in J_{max} (1874 pmol/min/cm²) was required to recover the observed DDI (Fig. 1E) after using the

Nelder-Mead minimization method and weighted least-squares algorithm in the simulators parameter estimation module (Jamei et al., 2014).

Assessing the Sensitivity of REF_{IBCRP} in IVIVE-PBPK. The sensitivity of the region-specific fraction of dose absorbed and enterocyte concentrations to REF_{IBCRP} generated by BPh (1.11) and the UoM (2.22) (Table 1; Harwood et al., 2016) was assessed for the BCRP test compound TC. Figure 2 shows the free segmental enterocyte concentration for TC, used as the driving force for apical efflux transporters. As expected, the lower BPh- REF_{IBCRP} leads to higher TC enterocyte concentrations in proximal regions than the higher UoM- REF_{IBCRP} , whereas an increasing importance of intestinal BCRP UoM- (REF_{IBCRP}) results in higher TC absorption and higher enterocyte concentration in the distal intestine due to the efflux activity promoting TC retention in the gut lumen and transit to the colon, a region with 7.7-fold lower BCRP levels (Harwood et al., 2013). The higher REF_{IBCRP} has a limited impact on lowering C_{max} and area under the curve (1.22- and 1.03-fold, respectively), but increases t_{max} 1.44-fold to 2.8 hours and is in line with clinical observations, where an inhibition of intestinal BCRP leads to a decrease in t_{max} (Schneck et al., 2004). Alongside differences in BCRP expression, interindividual variability in system parameters, such as the small intestinal transit time (range 0.5–10 hours; Yu et al., 1996), also contributes to region-specific fraction of dose absorbed [Supplemental Fig. 1; (Supplemental Materials)]. Acidic BCRP substrates, such as rosuvastatin, are expected to possess higher enterocyte concentrations, as limited metabolism, low




Fig. 1. (A) Sensitivity of C_{max} to REF_{IP-gp} for digoxin (single oral dose, 0.5 mg) in 100 HV individuals. The ranges of observed C_{max} values are given between the dashed lines, and the REF_{IP-gp} of 2.0 from Troutman and Thakker (2003b) and the UoM- REF_{IP-gp} (Harwood et al., 2016) (dashed arrows) are shown. (B) The mean digoxin plasma concentration when using the REF_{IP-gp} of 2.0 (Troutman and Thakker, 2003b) and the UoM REF_{IP-gp} of 0.4 in 100 HVs. The observed and predicted plasma concentrations for digoxin (single oral dose, 1 mg) in 80 HV individuals after the dosing of rifampicin (600 mg, 11 doses, once daily) using an REF_{IP-gp} of 2 (Troutman and Thakker, 2003b) (C), REF_{IP-gp} of 0.4 (Harwood et al., 2016) (D), and REF_{IP-gp} of 0.4 (E) after optimizing the J_{max} of P-gp by parameter estimation. The thin gray lines represent mean values for 10 individual virtual trials of eight individuals, males aged 21–37 years; the thick black lines are the overall means of the virtual population ($n = 80$); and the dashed gray lines are the 95th and 5th percentiles of the confidence interval. Open circles mark the observed digoxin concentrations when coadministered with multiple doses of rifampicin (Greiner et al., 1999).




Fig. 2. Simulated enterocyte concentration profiles for all intestinal segments of the model for the BCRP compound TC (10 mg oral, single dose, in solution) in 100 HVs using the BPh REF_{IBCRP} (1.11) or UoM- REF_{IBCRP} (2.22) with a mean small intestinal transit time of 3.34 hours, a passive apparent permeability of 115.2×10^{-6} cm/s, and an intrinsic clearance for BCRP of $17 \mu\text{l}/\text{min}/\text{cm}^2$.

passive permeation, and apical uptake transporters operate (Li et al., 2012; Jamei et al., 2014). Therefore, increased BCRP expression alters t_{max} and regional absorption, while not limiting overall absorption and bioavailability. This is dissimilar to the cooperation of P-gp and CYP3A4 activities that facilitate a drug's repeated exposure to intestinal CYP3A4, increasing overall gut metabolism and reducing bioavailability (Wacher et al., 1998). To our knowledge, the current study is the first highlighting this difference of the colocalized transporters P-gp and BCRP.

Combining IVIVE scalars and activity data generated from different laboratories for ATP-dependent transporters may not lead to successful IVIVE, whereas we postulate that laboratory-specific differences in REF may impact the mechanistic understanding of projected pharmacokinetic liabilities (efficacy/toxicity). This is due to in vitro activity, reproducibility of in vitro assays, culture conditions, and proteomic workflows. As discussed previously, direct translation of protein expression to activity may not always occur; therefore, accounting for deviations in this linear relationship via activity-abundance scalars will be required (Harwood et al., 2013). Ideally, scaling factors should be defined on a laboratory-specific basis against a common reference and combined with activity data from the same system. However, it is improbable within an industrial setting that groups will possess a bank of human intestinal tissues by which to obtain the in vivo abundance for

in-house intestinal REF generation. It is therefore advocated that commercially available pooled human intestinal microsomes (constituting ≥ 20 intestines) are used to generate an REF using the same proteomic methods as those used for quantifying in vitro system abundances used for determining activity. Alternatively, a link between human liver microsomes, intestinal microsomes, and Caco-2 cells can be approached.

Authorship Contributions

Participated in research design: Harwood, Neuhoff, Warhurst, Rostami-Hodjegan.

Conducted experiments: Harwood, Achour.

Contributed new reagents or analytic tools: Russell, Carlson.

Performed data analysis: Harwood, Achour, Neuhoff.

Wrote or contributed to the writing of the manuscript: Harwood, Achour, Neuhoff, Warhurst, Rostami-Hodjegan.

References

- Greiner B, Eichelbaum M, Fritz P, Kreichgauer HP, von Richter O, Zundler J, and Kroemer HK (1999) The role of intestinal P-glycoprotein in the interaction of digoxin and rifampin. *J Clin Invest* **104**:147–153.
- Harwood MD, Achour B, Neuhoff S, Russell MR, Carlson G, Warhurst G and Rostami-Hodjegan A (2016). In vitro-in vivo extrapolation scaling factors for intestinal P-glycoprotein and breast cancer resistance protein: Part I: A cross-laboratory comparison of transporter protein

- abundances and relative expression factors in human intestine and Caco-2 cells. *Drug Metab Dispos* **44**:297–307.
- Harwood MD, Neuhoff S, Carlson GL, Warhurst G, and Rostami-Hodjegan A (2013) Absolute abundance and function of intestinal drug transporters: a prerequisite for fully mechanistic in vitro-in vivo extrapolation of oral drug absorption. *Biopharm Drug Dispos* **34**:2–28.
- Hilgendorf C, Ahlin G, Seithel A, Artursson P, Ungell AL, and Karlsson J (2007) Expression of thirty-six drug transporter genes in human intestine, liver, kidney, and organotypic cell lines. *Drug Metab Dispos* **35**:1333–1340.
- Jamei M, Bajot F, Neuhoff S, Barter Z, Yang J, Rostami-Hodjegan A, and Rowland-Yeo K (2014) A mechanistic framework for in vitro-in vivo extrapolation of liver membrane transporters: prediction of drug-drug interaction between rosuvastatin and cyclosporine. *Clin Pharmacokinet* **53**:73–87.
- Li J, Wang Y, Zhang W, Huang Y, Hein K, and Hidalgo IJ (2012) The role of a basolateral transporter in rosuvastatin transport and its interplay with apical breast cancer resistance protein in polarized cell monolayer systems. *Drug Metab Dispos* **40**:2102–2108.
- Neuhoff S, Yeo KR, Barter Z, Jamei M, Turner DB, and Rostami-Hodjegan A (2013a) Application of permeability-limited physiologically-based pharmacokinetic models: part I-digoxin pharmacokinetics incorporating P-glycoprotein-mediated efflux. *J Pharm Sci* **102**:3145–3160.
- Neuhoff S, Yeo KR, Barter Z, Jamei M, Turner DB, and Rostami-Hodjegan A (2013b) Application of permeability-limited physiologically-based pharmacokinetic models: part II - prediction of P-glycoprotein mediated drug-drug interactions with digoxin. *J Pharm Sci* **102**:3161–3173.
- Oswald S, Gröer C, Drozdziak M, and Siegmund W (2013) Mass spectrometry-based targeted proteomics as a tool to elucidate the expression and function of intestinal drug transporters. *AAPS J* **15**:1128–1140.
- Rostami-Hodjegan A (2012) Physiologically based pharmacokinetics joined with in vitro-in vivo extrapolation of ADME: a marriage under the arch of systems pharmacology. *Clin Pharmacol Ther* **92**:50–61.
- Schneck DW, Birmingham BK, Zalikowski JA, Mitchell PD, Wang Y, Martin PD, Lasseter KC, Brown CDA, Windass AS, and Raza A (2004) The effect of gemfibrozil on the pharmacokinetics of rosuvastatin. *Clin Pharmacol Ther* **75**:455–463.
- Seithel A, Karlsson J, Hilgendorf C, Björquist A, and Ungell AL (2006) Variability in mRNA expression of ABC- and SLC-transporters in human intestinal cells: comparison between human segments and Caco-2 cells. *Eur J Pharm Sci* **28**:291–299.
- Taipalensuu J, Tömbblom H, Lindberg G, Einarsson C, Sjöqvist F, Melhus H, Garberg P, Sjöström B, Lundgren B, and Artursson P (2001) Correlation of gene expression of ten drug efflux proteins of the ATP-binding cassette transporter family in normal human jejunum and in human intestinal epithelial Caco-2 cell monolayers. *J Pharmacol Exp Ther* **299**:164–170.
- Troutman MD and Thakker DR (2003a) Efflux ratio cannot assess P-glycoprotein-mediated attenuation of absorptive transport: asymmetric effect of P-glycoprotein on absorptive and secretory transport across Caco-2 cell monolayers. *Pharm Res* **20**:1200–1209.
- Troutman MD and Thakker DR (2003b) Novel experimental parameters to quantify the modulation of absorptive and secretory transport of compounds by P-glycoprotein in cell culture models of intestinal epithelium. *Pharm Res* **20**:1210–1224.
- Vildhede A, Karlgren M, Svedberg EK, Wisniewski JR, Lai Y, Norén A, and Artursson P (2014) Hepatic uptake of atorvastatin: influence of variability in transporter expression on uptake clearance and drug-drug interactions. *Drug Metab Dispos* **42**:1210–1218.
- von Richter O, Glavinas H, Krajcsi P, Liehner S, Siewert B, and Zech K (2009) A novel screening strategy to identify ABCB1 substrates and inhibitors. *Naunyn-Schmiedeberg's Arch Pharmacol* **379**:11–26.
- Wacher VJ, Silverman JA, Zhang Y, and Benet LZ (1998) Role of P-glycoprotein and cytochrome P450 3A in limiting oral absorption of peptides and peptidomimetics. *J Pharm Sci* **87**:1322–1330.
- Yang J, Jamei M, Yeo KR, Tucker GT, and Rostami-Hodjegan A (2007) Prediction of intestinal first-pass drug metabolism. *Curr Drug Metab* **8**:676–684.
- Yu LX, Crison JR, and Amidon GL (1996) Compartmental transit and dispersion model analysis of small intestinal transit flow in humans. *Int J Pharm* **140**:111–118.

Address correspondence to: Dr. Matthew D. Harwood, Simcyp Ltd. (a Certara company), Blades Enterprise Centre, John Street, Sheffield S2 4SU, UK. E-mail: matthew.harwood@certara.com

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition. Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. Drug Metabolism & Disposition.

SUPPLEMENTAL INFORMATION

Table S1. Publications reporting P-gp (MDR1) and BCRP mRNA and protein expression in human jejunum obtained from literature analysis.

Protein (Gene)	mRNA/ Protein	Technique	Included for REF analysis (Yes (Y)/No (N))	Reason for Inclusion or Exclusion	Source
P-gp (MDR1)	mRNA	Slot Blot	N	Exclusion: No literature data available for Caco-2 MDR1 mRNA expression using the Slot Blot technique from this laboratory	(Fojo et al., 1987)
P-gp (MDR1) & BCRP	mRNA	RT-PCR	Y	Inclusion: Data available using the same techniques from the same laboratory for both human jejunum and Caco-2 cell monolayers	(Taipalensuu et al., 2001)
P-gp (MDR1)	mRNA/ Protein	RT-PCR & Western Blot	N	Exclusion: Data available for Caco-2 cells and jejunum MDR1 mRNA expression using this RT-PCR technique from this laboratory (Goto et al., 2003), however the Caco-2 cells were grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	(Hashida et al., 2001)
P-gp (MDR1)	Protein	Western Blot	Y	Inclusion: Data available using the same techniques from the same laboratory for both human jejunum and Caco-2 cell monolayers	(Troutman and Thakker, 2003)
P-gp (MDR1)	Protein	Western Blot	N	Exclusion: Data available for Caco-2 and jejunum MDR1 mRNA expression using this RT-PCR technique from this laboratory, however the Caco-2 cells were a non-standard phenotype having been cultured with $1\alpha,25(\text{OH})_2\text{-vitamin D}_3$ to induce CYP3A4 potentially influencing P-gp expression	(Mouly and Paine, 2003)
P-gp (MDR1)	mRNA	RT-PCR	N	Exclusion: Data available for Caco-2 cells and jejunum MDR1 mRNA expression using this RT-PCR technique	(Masuda et al., 2005)

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition. Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. Drug Metabolism & Disposition.

				from this laboratory (Goto et al., 2003), however the Caco-2 cells were grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	
P-gp (MDR1)	mRNA	RT-PCR	N	Exclusion: Data available for Caco-2 cells and jejunum MDR1 mRNA expression using this RT-PCR technique from this laboratory (Goto et al., 2003), however the Caco-2 cells were grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	(Terada et al., 2005)
P-gp (MDR1) & BCRP	mRNA	RT-PCR	N	Exclusion: Data available for Caco-2 and jejunum mRNA MDR1 expression from the same laboratory, however the Caco-2 expression data (Taipalensuu et al., 2001) are not in the same units (transcripts / μ g total RNA) as for the jejunum study by Englund et al. (2006) (relative expression to villin)	(Englund et al., 2006)
P-gp (MDR1) & BCRP	mRNA	RT-PCR	Y	Inclusion: Data available using the same techniques from the same laboratory for both human jejunum and Caco-2 cell monolayers (23d cultivated)	(Seithel et al., 2006)
P-gp (MDR1)	mRNA	RT-PCR	N	Exclusion: Data available for Caco-2 and jejunum mRNA MDR1 expression from the same laboratory, however the Caco-2 expression data (Taipalensuu et al., 2001) are not in the same units (transcripts / μ g total RNA) as for the study by Berggren et al. (2007) (percentage of integrated optical density for villin)	(Berggren et al., 2007)
P-gp (MDR1)	mRNA	RT-PCR	N	Exclusion: No literature data available for Caco-2 MDR1 mRNA expression using the RT-PCR technique from these laboratories	(Canaparo et al., 2007)
P-gp (MDR1) & BCRP	mRNA	RT-PCR	Y	Inclusion: Data available using the same techniques from the same laboratory for both human jejunum and Caco-2 cell monolayers (16d cultivated). <u>Note, it is</u>	(Hilgendorf et al., 2007)

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition. Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. Drug Metabolism & Disposition.

				<u>understood that 4 of the same jejunum samples reported in Seithel et al. (2006) are reported in the 5 samples Hilgendorf et al. (2007)</u>	
P-gp (MDR1) & BCRP	Protein	Western Blot	Y	Inclusion: Data available using the same techniques from the same laboratory for both human jejunum and Caco-2 cell monolayers	(von Richter et al., 2009)
P-gp (MDR1) & BCRP	Protein	Western Blot	N	Exclusion: No data available for Caco-2 MDR1 mRNA expression using the Western blotting technique from these laboratories	(Bruyere et al., 2010)
P-gp (MDR1)	mRNA	RT-PCR	N	Exclusion: No literature data available for Caco-2 MDR1 mRNA expression using the RT-PCR technique from this laboratory	(Ulvestad et al., 2013)
P-gp (MDR1) & BCRP	Protein	QTAP	N	Exclusion: The Caco-2 abundance data for this study were quantified from Caco-2 cells grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	(Oswald et al., 2013)
P-gp (MDR1) & BCRP	Protein	QTAP	N	Exclusion: Caco-2 cells P-gp abundances are available from the same laboratory (Oswald et al., 2013) as for this study, however these were quantified from Caco-2 cells grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	(Gröer et al., 2013)
P-gp (MDR1) & BCRP	Protein	QTAP	N	Exclusion: Caco-2 cells P-gp abundances are available from the same laboratory (Oswald et al., 2013) as for this study, however these were quantified from Caco-2 cells grown on plastic and not filter systems, therefore are not relevant for transporter assays and IVIVE	(Drozdik et al., 2014)
P-gp (MDR1) & BCRP	Protein	QTAP	Y	Inclusion: Data available using the same techniques from the same two laboratories for both human jejunum and Caco-2 cell monolayers	(Harwood et al., submitted) (currently unavailable in PubMed database)

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition.
Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. Drug Metabolism & Disposition.

TECHNICAL NOTE: RATIONALE FOR BUILDING TC

Compounds that interact with BCRP are generally low permeability compounds that generally require apical intestinal uptake transporters to gain access to the enterocyte, consequently there are no selective BCRP compounds on the market. In addition, due to the limited availability of intrinsic kinetic data ($CL_{int,T}$, or J_{max} & K_m) for compounds that interact with BCRP in Caco-2 monolayers, a theoretical compound (TC) was built to assess the impact of laboratory-specific REFs for BCRP generated by different laboratories and different methods on the same samples. TC was designed to possess high 'passive' lipoidal bilayer permeability (apparent permeability; 115×10^{-6} cm/s), so access to the binding site of BCRP via the enterocytes cytoplasm was not limited by its' permeability through the membrane. As the BCRP REFs generated by LC-MS/MS quantification were based on human jejunum samples, it was important to ensure that an TC was absorbed primarily in the jejunum, therefore after sensitivity analysis assessing the relationship between fa/C_{max} , passive permeability and intestinal BCRP $CL_{int,T}$, an intestinal $CL_{int,T}$ for BCRP was of $17 \mu\text{L}/\text{min}/\text{cm}^2$ was assigned. Finally, to reflect the negligible metabolism observed for acidic BCRP substrates such as rosuvastatin, there was no gut metabolism assigned for TC.

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition.
Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. Drug Metabolism & Disposition.

FIGURE




Figure S1. Assessing the impact of small intestinal transit time (SITT) (0.5, 3.5, 6.5 and 9.5 hours) on the regional f_a in all intestinal compartments. The passive P_{app} (Caco-2, 115.2×10^{-6} cm/s), an intestinal $CL_{int,BCRP}$ of $17 \mu\text{L}/\text{min}/\text{cm}^2$ and REF_{iBCRP} of 1.11 were used. The values are given as a mean of 100 individuals \pm standard deviation.

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition.

Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. *Drug Metabolism & Disposition*.

REFERENCES

- Berggren S, Gall C, Wollnitz N, Ekelund M, Karlbom U, Hoogstraate J, Schrenk D and Lennernas H (2007) Gene and protein expression of P-glycoprotein, MRP1, MRP2, and CYP3A4 in the small and large human intestine. *Mol Pharm* **4**:252-257.
- Bruyere A, Decleves X, Bouzom F, Ball K, Marques C, Treton X, Pocard M, Valleur P, Bouhnik Y, Panis Y, Scherrmann JM and Mouly S (2010) Effect of variations in the amounts of P-glycoprotein (ABCB1), BCRP (ABCG2) and CYP3A4 along the human small intestine on PBPK models for predicting intestinal first pass. *Mol Pharm* **7**:1596-1607.
- Canaparo R, Finnstrom N, Serpe L, Nordmark A, Muntoni E, Eandi M, Rane A and Zara GP (2007) Expression of CYP3A isoforms and P-glycoprotein in human stomach, jejunum and ileum. *Clin Exp Pharmacol Physiol* **34**:1138-1144.
- Drozdik M, Groer C, Penski J, Lapczuk J, Ostrowski M, Lai Y, Prasad B, Unadkat JD, Siegmund W and Oswald S (2014) Protein abundance of clinically relevant multidrug transporters along the entire length of the human intestine. *Mol Pharm* **11**:3547-3555.
- Englund G, Rorsman F, Ronnblom A, Karlbom U, Lazorova L, Grasjo J, Kindmark A and Artursson P (2006) Regional levels of drug transporters along the human intestinal tract: co-expression of ABC and SLC transporters and comparison with Caco-2 cells. *Eur J Pharm Sci* **29**:269-277.
- Fojo AT, Ueda K, Slamon DJ, Poplack DG, Gottesman MM and Pastan I (1987) Expression of a multidrug-resistance gene in human tumors and tissues. *Proc Natl Acad Sci U S A* **84**:265-269.
- Goto M, Masuda S, Saito H and Inui K (2003) Decreased expression of P-glycoprotein during differentiation in the human intestinal cell line Caco-2. *Biochem Pharmacol* **66**:163-170.
- Gröer C, Bruck S, Lai Y, Paulick A, Busemann A, Heidecke CD, Siegmund W and Oswald S (2013) LC-MS/MS-based quantification of clinically relevant intestinal uptake and efflux transporter proteins. *J Pharm Biomed Anal* **85**:253-261.
- Harwood MD, Achour B, Neuhoff S, Russell MR, Carlson G, Warhurst G and Rostami-Hodjegan. In vitro-in vivo extrapolation factors for intestinal P-glycoprotein and breast cancer resistance protein: Part I: A cross-laboratory comparison of transporter protein abundances and relative expression factors in human intestine and Caco-2 cells. *Drug Metab Dispos* (submitted for peer review).
- Hashida T, Masuda S, Uemoto S, Saito H, Tanaka K and Inui K (2001) Pharmacokinetic and prognostic significance of intestinal MDR1 expression in recipients of living-donor liver transplantation. *Clin Pharmacol Ther* **69**:308-316.
- Hilgendorf C, Ahlin G, Seithel A, Artursson P, Ungell AL and Karlsson J (2007) Expression of thirty-six drug transporter genes in human intestine, liver, kidney, and organotypic cell lines. *Drug Metab Dispos* **35**:1333-1340.
- Masuda S, Goto M, Okuda M, Ogura Y, Oike F, Kiuchi T, Tanaka K and Inui K (2005) Initial dosage adjustment for oral administration of tacrolimus using the intestinal MDR1 level in living-donor liver transplant recipients. *Transplant Proc* **37**:1728-1729.
- Mouly S and Paine MF (2003) P-glycoprotein increases from proximal to distal regions of human small intestine. *Pharm Res* **20**:1595-1599.

In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicting Drug Disposition.

Matthew D Harwood, Brahim Achour, Sibylle Neuhoff, Matthew R Russell, Gordon Carlson, Geoffrey Warhurst, Amin Rostami-Hodjegan. *Drug Metabolism & Disposition*.

- Oswald S, Groer C, Drozdik M and Siegmund W (2013) Mass spectrometry-based targeted proteomics as a tool to elucidate the expression and function of intestinal drug transporters. *AAPS J* **15**:1128-1140.
- Seithel A, Karlsson J, Hilgendorf C, Bjorquist A and Ungell AL (2006) Variability in mRNA expression of ABC- and SLC-transporters in human intestinal cells: comparison between human segments and Caco-2 cells. *Eur J Pharm Sci* **28**:291-299.
- Taipalensuu J, Tornblom H, Lindberg G, Einarsson C, Sjoqvist F, Melhus H, Garberg P, Sjostrom B, Lundgren B and Artursson P (2001) Correlation of gene expression of ten drug efflux proteins of the ATP-binding cassette transporter family in normal human jejunum and in human intestinal epithelial Caco-2 cell monolayers. *J Pharmacol Exp Ther* **299**:164-170.
- Terada T, Shimada Y, Pan X, Kishimoto K, Sakurai T, Doi R, Onodera H, Katsura T, Imamura M and Inui K (2005) Expression profiles of various transporters for oligopeptides, amino acids and organic ions along the human digestive tract. *Biochem Pharmacol* **70**:1756-1763.
- Troutman MD and Thakker DR (2003) Novel experimental parameters to quantify the modulation of absorptive and secretory transport of compounds by P-glycoprotein in cell culture models of intestinal epithelium. *Pharm Res* **20**:1210-1224.
- Ulvestad M, Skottheim IB, Jakobsen GS, Bremer S, Molden E, Asberg A, Hjelmessaeth J, Andersson TB, Sandbu R and Christensen H (2013) Impact of OATP1B1, MDR1, and CYP3A4 expression in liver and intestine on interpatient pharmacokinetic variability of atorvastatin in obese subjects. *Clin Pharmacol Ther* **93**:275-282.
- von Richter O, Glavinas H, Krajcsi P, Liehner S, Siewert B and Zech K (2009) A novel screening strategy to identify ABCB1 substrates and inhibitors. *Naunyn Schmiedebergs Arch Pharmacol* **379**:11-26.