

# Detection of Weak Organic Anion-Transporting Polypeptide 1B Inhibition by Probenecid with Plasma-Based Coproporphyrin in Humans<sup>§</sup>

Yueping Zhang, Vinay K. Holenarsipur, Hamza Kandoussi, Jianing Zeng, T. Thanga Mariappan, Michael Sinz, and Hong Shen

Departments of Metabolism and Pharmacokinetics (Y.Z., M.S., H.S.) and Bioanalytical Sciences (H.K., J.Z.), Bristol Myers Squibb Company, Princeton, New Jersey; and Departments of Metabolism and Pharmacokinetics (V.K.H., T.T.M.), Biocon Bristol Myers Squibb R&D Centre (BBRC), Syngene International Ltd., Biocon Park, Bommasandra IV Phase, Bangalore, India

Received April 11, 2020; accepted July 13, 2020

## ABSTRACT

Probenecid (PROB) is a clinical probe inhibitor of renal organic anion transporter (OAT) 1 and OAT3 that inhibits *in vitro* activity of hepatic drug transporters OATP1B1 and OATP1B3. It was hypothesized that PROB could potentially affect the disposition of OATP1B drug substrates. The plasma levels of the OATP1B endogenous biomarker candidates, including coproporphyrin I (CPI), CPIII, hexadecanedioate (HDA), and tetradecanedioate (TDA), were examined in 14 healthy subjects treated with PROB. After oral administration with 1000 mg PROB alone and in combination with furosemide (FSM),  $AUC_{(0-24\text{ h})}$  values were 1.39 ± 0.21-fold and 1.57 ± 0.41-fold higher than predose levels for CPI and 1.34 ± 0.16-fold and 1.45 ± 0.57-fold higher for CPIII. Despite increased systemic exposures, no decreases in CPI and CPIII renal clearance were observed (0.97 ± 0.38-fold and 1.16 ± 0.51-fold for CPI, and 1.34 ± 0.53-fold and 1.50 ± 0.69-fold for CPIII, respectively). These results suggest that the increase of CP systemic exposure is caused by OATP1B inhibition. Consistent with this hypothesis, PROB inhibited OATP1B1- and OATP1B3-mediated transport of CPI in a concentration-dependent manner, with  $IC_{50}$  values of 167 ± 42.0 and 76.0 ± 17.2 μM, respectively, in transporter-overexpressing human embryonic kidney cell assay. The inhibition potential was further confirmed by CPI and CPIII hepatocyte uptake

experiments. In contrast, administration of PROB alone did not change  $AUC_{(0-24\text{ h})}$  of HDA and TDA relative to prestudy levels, although the administration of PROB in combination with FSM increased HDA and TDA levels compared with FSM alone (1.02 ± 0.18-fold and 0.90 ± 0.20-fold vs. 1.71 ± 0.43-fold and 1.62 ± 0.40-fold). Taken together, these findings indicate that PROB displays weak OATP1B inhibitory effects *in vivo* and that coproporphyrin is a sensitive endogenous probe of OATP1B inhibition. This study provides an explanation for the heretofore unknown mechanism responsible for PROB's interaction with other xenobiotics.

## SIGNIFICANCE STATEMENT

This study suggested that PROB is a weak clinical inhibitor of OATP1B based on the totality of evidence from the clinical interaction between PROB and CP and the *in vitro* inhibitory effect of PROB on OATP1B-mediated CP uptake. It demonstrates a new methodology of utilizing endogenous biomarkers to evaluate complex drug-drug interaction, providing explanation for the heretofore unknown mechanism responsible for PROB's inhibition. It provides evidence to strengthen the claim that CP is a sensitive circulating endogenous biomarker of OATP1B inhibition.

## Introduction

It is well established that membrane-bound transport systems are primarily critical determinants in the cellular traffic and disposition of exogenous and endogenous compounds. Notably, organic anion-transporting polypeptide (OATP) 1B1 and OATP1B3 can govern the disposition of drug substrates, including the widely used 3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors (statins). Moreover,

it is increasingly recognized that endogenous biomarkers of OATP1B1 and OATP1B3—such as coproporphyrin I (CPI), CPIII, hexadecanedioate (HDA), and tetradecanedioate (TDA)—and glycochenodeoxycholate-3-sulfate are useful tools in elucidating the role of OATP1B in complex drug-drug interaction (DDI) in healthy subjects and patients (Lai et al., 2016; Shen et al., 2016, 2017, 2018, Takehara et al., 2018, Barnett et al., 2019; Yee et al., 2019; Jones et al., 2020; Mori et al., 2020).

Probenecid (PROB) is widely used as a uricosuric agent (Cunningham et al., 1981). In addition, PROB has been used as an adjunct to enhance blood levels of antibiotics such as penicillins and cephalosporins because of an inhibitory effect on the renal transporters OAT1 and

This study is supported by Bristol Myers Squibb Company.

<https://doi.org/10.1124/dmd.120.000076>

<sup>§</sup>This article has supplemental material available at [dmd.aspetjournals.org](http://dmd.aspetjournals.org).

**ABBREVIATIONS:** *AUC*, area under plasma concentration–time curve;  $AUC_{(0-24\text{ h})}$ , area under plasma concentration–time curve from time 0 to 24 hours; CCK-8, cholecystokinin octapeptide;  $CL_R$ , renal clearance; CP, coproporphyrin; CPI, coproporphyrin I; CPIII, coproporphyrin III; DDI, drug–drug interaction; E17βG, estradiol-17β-D-glucuronide; FDA, Food and Drug Administration; FSM, furosemide; HBSS, Hanks' balanced salt solution; HDA, hexadecanedioate; HEK, human embryonic kidney; LC-MS/MS, liquid chromatography–tandem mass spectrometry; MRP, multidrug resistance-associated protein; OAT, organic anion transporter; OATP, organic anion-transporting polypeptide; PROB, probenecid; RIF SV, rifampicin SV; RSV, rosuvastatin; *R*-value, ratio of victim *AUC* in the presence and absence of perpetrators; TDA, tetradecanedioate.

OAT3. In agreement, most of the DDIs involving PROB are due to its inhibition on the kidney transport of acidic drugs such as cefaclor (Welling et al., 1979), cefonicid (Pitkin et al., 1981), cefoxitin (Vlasses et al., 1980), cephadrine (Welling et al., 1979), dicloxacillin (Beringer et al., 2008), famotidine (Inotsume et al., 1990), and furosemide (Vree et al., 1995; Shen et al., 2019a). As a result, the US Food and Drug Administration (FDA) suggests PROB as an index inhibitor to assess OAT1 and OAT3 in clinical DDI studies (<https://www.fda.gov/drugs/drug-interactions-labeling/drug-development-and-drug-interactions-table-substrates-inhibitors-and-inducers>). However, knowledge of the transporters involved in PROB inhibition could be incomplete. For example, some studies showed that the action of PROB at the renal level was not sufficient to account for severalfold serum level elevations encountered for drugs such as fexofenadine, statins, and methotrexate in humans and animals (Gewirtz et al., 1984; Yasui-Furukori et al., 2005; Liu et al., 2008; Kosa et al., 2018). As an organic acid, probenecid increased the area under plasma concentration–time curve (*AUC*) of fexofenadine approximately 1.5-fold by interfering with secretion of fexofenadine by the renal tubules in humans (Yasui-Furukori et al., 2005; Liu et al., 2008). The total and renal clearance ( $CL_R$ ) values of fexofenadine were 16.0 and 6.2 L/h (Lappin et al., 2010). Given the fact that fexofenadine is 60%–70% bound to plasma proteins (Molimard et al., 2004), the active tubular secretion contributes to approximately half of the  $CL_R$ . As a result, inhibition of OAT3-mediated renal secretion clearance alone, which resulted in a decreased amount of dose excreted in urine from 11%–12% to 6%–8%, cannot explain the observed *AUC* change (Yasui-Furukori et al., 2005; Liu et al., 2008). Moreover, studies in cynomolgus monkeys and rats have demonstrated that probenecid decreased the clearance of statins and methotrexate by the hepatic route as well (Gewirtz et al., 1984; Kosa et al., 2018), contributing to prolonged elevation of circulating statins and methotrexate levels. The reduction in hepatic uptake of statins in monkey hepatocytes and biliary secretion of methotrexate in rats is thought to arise from inhibition of hepatic uptake mediated by OATP in the presence of PROB. However, a direct interaction between PROB and OATP1B in humans has not been fully tested.

Recent bioanalytical methodologies, especially metabolomics, have identified a number of endogenous biomarker candidates for the transporter proteins expressed in the organs of importance in drug disposition, including liver and kidney (Chu et al., 2018; Muller et al., 2018; Rodrigues et al., 2018; Shen, 2018). Changes in the levels of such endogenous probes were able to phenotype the mechanism of complex DDIs involving multiple elimination pathways. For example, a clinical DDI study was conducted to assess the inhibition potential of fenebrutinib using midazolam (CYP3A), simvastatin (CYP3A and OATP1B), and rosuvastatin (Breast cancer resistance protein and OATP1B) as drug substrates (Jones et al., 2020). Fenebrutinib increased the *AUC* values of all three probes 2- to 3-fold. However, there was no change in the plasma levels of CPI and CPIII, endogenous biomarkers of OATP1B1, suggesting that the DDIs were due to the inhibition of CYP3A and breast cancer resistance protein rather than OATP1B (Jones et al., 2020). The present studies were designed to evaluate the inhibitory effects of PROB on human OATP1B1 and OATP1B3 by analyzing OATP1B biomarker levels in a clinical study and evaluating coproporphyrin uptake in transfected cell lines and hepatocytes in the presence of PROB to determine whether PROB modulates OATP1B activity.

## Materials and Methods

**Reagents and Materials.** CPI and CPIII were purchased as dihydrochloride salts from Frontier Scientific (Logan, UT). HDA, TDA, carbamazepine, and high-performance liquid chromatography–grade methanol, acetonitrile, and water were purchased from Sigma Aldrich (St. Louis,

MO). Rosuvastatin (RSV), estradiol-17 $\beta$ -D-glucuronide (E17 $\beta$ G), cholecystokinin octapeptide (CCK-8), CPI-<sup>15</sup>N<sub>4</sub> sodium bisulfate salt, HDA-d4, TDA-d4, and rifampicin SV (RIF SV) were purchased from Toronto Research Chemicals (North York, ON, Canada). Water-soluble probenecid sodium salt was purchased from Life Technologies (Carlsbad, CA). CPIII-d8 was synthesized at Bristol Myers Squibb Company (Princeton, NJ). [<sup>3</sup>H]E17 $\beta$ G (38.8 Ci/mmol) and [<sup>3</sup>H]CCK-8 (92.5mCi/mmol) were purchased from PerkinElmer Life and Analytical Sciences (Waltham, MA). Charcoal stripped plasma was obtained from Bioreclamation IVT (Westbury, NY). Cryopreserved human hepatocytes from two female donors were purchased from Celsis In Vitro Technologies (Baltimore, MD) (lot NRJ) and Bioreclamation IVT (lot BXW). Human plasma containing dipotassium EDTA was obtained from Biologic Specialty Corporation (Colmar, PA). Cell culture reagents were purchased from Corning (Manassas, VA) and Life Technologies Corporation. All other reagents and chemicals used were of the highest grade commercially available.

**Clinical Drug Interaction Study between PROB and FSM.** Plasma and urine samples were collected from an open-label, single-dose, three-treatment, three-period clinical DDI study between PROB and FSM reported previously (Shen et al., 2019a). Briefly, 14 male healthy Indian subjects who had a normal medical history and physical examination participated in the study. Each subject received 1000 mg PROB alone (period 1), 40 mg FSM alone (period 2), and 40 mg FSM at 1 hour after administration of 1000 mg PROB (period 3) with a 1-week washout between treatments. Subjects fasted the night before and until at least 4 hours after administration of the drugs. PROB and FSM were administered orally with 240 ml of water. Blood samples (3 ml each) for determination of drug and transporter endogenous biomarker concentrations were obtained at predose and at 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 4.0, 6.0, 8.0, 12.0, and 24.0 hours after dosing. Urine samples were collected during 0–8, 8–16, and 16–24 hours postdose. Subjects were housed in a clinical facility 36 hours prior to dosing in period 1, and blood and urine samples were also collected over 24 hours prior to dosing to obtain the baseline levels of transporter endogenous biomarkers in the absence of drug. Plasma was separated immediately, and the plasma and urine samples were prepared into two aliquots and kept at –70°C until analysis using liquid chromatography–tandem mass spectrometry (LC-MS/MS) (Shen et al., 2019a).

**Quantification of CPI and CPIII in Plasma and Urine by LC-MS/MS.** Plasma and urine concentrations of CPI and CPIII were quantitated by using the methods developed previously (Kandoussi et al., 2018).

**Quantification of HDA and TDA in Plasma by LC-MS/MS.** The bioanalytical analyses of HDA and TDA in plasma were performed as described in detail by Santockyte et al. (2018).

**Inhibition Studies in OATP1B1- and OATP1B3-Expressing HEK293 Cells.** The in vitro uptake experiments were repeated in at least two independent experiments. There are three replicates per experiment for each condition ( $n = 3$ ). The stable OATP1B1- and OATP1B3-transfected and mock cells generated by Bristol Myers Squibb were cultured at 37°C in an atmosphere of 95% air/5% CO<sub>2</sub> and subcultured once per week (Han et al., 2010; Shen et al., 2013). OATP1B1- and OATP1B3-expressing cells used in the current study were passaged fewer than 30 times to retain consistent transporter expression and functional activity. To assess the inhibition potential of PROB toward OATP1B1 and OATP1B3, the inhibitory effect was evaluated on the uptake of CPI and radiolabeled probe substrates in the presence of increasing concentrations of PROB in transporter-expressing cells using a protocol described previously (Shen et al., 2017; Panfen et al., 2019). In brief, transporter-expressing cells were grown to confluence in poly-D-lysine-coated 24-well plates (BD Biosciences, San Jose, CA). At confluence, medium was aspirated, and cells were rinsed twice with 2 ml of prewarmed Hanks' balanced salt solution (HBSS) (catalog number 21-023-CM; Mediatech, Manassas, VA) and preincubated with 200  $\mu$ l of the transport buffer (HBSS with 10 mM HEPES, pH 7.4) containing PROB at concentrations ranging from 1 to 10,000  $\mu$ M for 30 minutes at 37°C. Uptake was then initiated by the addition of 200  $\mu$ l of the prewarmed transport buffer containing CPI (0.2  $\mu$ M) or radiolabeled probe substrate (1  $\mu$ M [<sup>3</sup>H]E17 $\beta$ G and 0.1  $\mu$ M [<sup>3</sup>H]CCK-8 for OATP1B1 and OATP1B3, respectively). The probe substrate concentrations are well below the  $K_m$  values. Incubation proceeded for 2 minutes at 37°C to ensure linearity with time (Supplemental Fig. 1). After 2 minutes, the uptake buffer was then removed, and the cell monolayers were immediately washed three times with 1.5 ml ice-cold HBSS buffer to terminate the uptake process. To analyze the concentrations of radiolabeled compounds, cells in the dried plate were lysed with

300  $\mu$ l 0.1% Triton X-100. Aliquots of 200 and 20  $\mu$ l cell lysate were used for radioactivity counting and protein concentration analysis, respectively. After the addition of 5 ml of scintillation cocktail, radioactivity was measured on a dual-channel liquid scintillation counter, Tri-Carb 3100TR liquid scintillation counter (PerkinElmer Life Sciences, Boston, MA). To analyze the concentrations of CPI, cells in the dried plate were lysed in 300  $\mu$ l 2:1 (v/v) ratio of acetonitrile and 1 M formic acid with CPI- $^{15}$ N $_4$  (internal standard). The contents were filtered through a 96-well filter plate (0.45- $\mu$ m low-binding hydrophilic polytetrafluoroethylene), and the filtrate was dried under nitrogen gas. The dried contents were reconstituted in 80  $\mu$ l of 1 M formic acid/acetonitrile (20:80, v/v), and the concentration of CPI was measured by LC-MS/MS. Throughout the entire process, appropriate precautions were taken to minimize sample exposure to ambient light. Protein concentration of cell lysates was determined with the bicinchoninic acid protein assay kit (Pierce Chemical, Rockford, IL). Cellular uptake in OATP1B1- and OATP1B3-HEK cells was normalized based on the protein amount in each well.

**Inhibition of CP Uptake in Human Hepatocytes by PROB.** To further assess the inhibitory effect of PROB on OATP1B, the uptake of CPI and CPIII in human hepatocytes in the presence of PROB was examined using a protocol described previously, with some modification (Shen et al., 2016; Zhang et al., 2019). Briefly, transporter-qualified cryopreserved human hepatocytes from two female donors were thawed according to the vendor's instructions (Bioreclamation IVT). The hepatocytes were then pooled by resuspending the cell pellet in Krebs-Henseleit buffer to give a cell density of 2 million viable cells per milliliter, which was determined by trypan blue staining ( $\geq 80\%$  post-thaw viability). After cell suspensions were prewarmed to 37°C, the uptake of CPI in hepatocytes in the presence and absence of an inhibitor (final concentration of 100, 300, and 1000  $\mu$ M PROB and 200  $\mu$ M RIF SV) was determined. The uptake study was initiated by adding an equal volume of CP solution to the hepatocyte suspensions at 37°C, resulting in a final cell density of 1 million viable cells per milliliter and a CP or RSV concentration of 1  $\mu$ M. There was a 3-minute preincubation with the inhibitors or control buffer with hepatocytes at 37°C before the initiation of uptake. At 0.25, 1, 1.5, and 5 minutes, 100- $\mu$ l reaction mixtures were removed and overlaid onto prepared 0.4-ml microcentrifuge tubes containing 50  $\mu$ l 2 M ammonium acetate (bottom layer) and 100  $\mu$ l filtration oil (top layer; 84.5:15.5 silicon oil–mineral oil mix, final density of 1.015). Samples were centrifuged immediately at 10,000g for 15 seconds using a benchtop centrifuge to pellet the cells. The tubes were placed on dry ice and then cut, and the cell pellet was digested in a 2:1 (v/v) ratio of acetonitrile and 1 M formic acid containing the internal standard at room temperature. The contents were filtered through a 96-well filter plate (0.45- $\mu$ m low-binding hydrophilic polytetrafluoroethylene), and the filtrate was dried under nitrogen gas. Finally, the dried contents were reconstituted in 80  $\mu$ l of 1 M formic acid/acetonitrile (20:80, v/v) for LC-MS/MS

analysis. Throughout the entire process, appropriate precautions were taken to minimize sample exposure to ambient light.

**LC-MS/MS Analysis of CPI and CPIII in Cell Lysates.** Cell lysate concentrations of CPI, CPIII, and RSV were measured by LC-MS/MS as described previously (Lai et al., 2016; Shen et al., 2016).

**Pharmacokinetic and Transport Analyses.** The pharmacokinetic parameters of CPI, CPIII, HDA, and TDA were derived using Phoenix WinNonlin, version 8.1 (Certara, Princeton, NJ). Maximum plasma concentration ( $C_{max}$ ) and area under the concentration-time curve from time 0 to 24 hours [ $AUC_{(0-24 h)}$ ] were obtained from plasma concentrations versus time data by performing a noncompartmental analysis with mixed log-linear trapezoidal method.  $CL_R$  was estimated by the following equation:

$$CL_R = \frac{A_{e(0-24h)}}{AUC_{(0-24h)}}$$

where  $A_{e(0-24 h)}$  is the cumulative amount excreted in the urine during the time interval from 0 to 24 hours.

The concentrations required to inhibit transport by 50% ( $IC_{50}$ ) of PROB toward OATP1B1- and OATP1B3-mediated uptake of CPI and radiolabeled probes were estimated by fitting the uptake data to the following equation using the nonlinear regression approach (Phoenix WinNonlin; Certara):


$$V = V_0 \times \left( 1 - \frac{I^h}{I^h + IC_{50}^h} \right)$$

where  $V$  is the mean transporter-mediated uptake of substrate at 2 minutes observed at the given PROB testing concentration ( $I$ ),  $V_0$  is the uptake in the absence of PROB, and  $h$  is the Hill slope factor.

The hepatic uptake rate in the absence and presence of inhibitor was calculated based on the initial rate of uptake during the linear phase (up to 1.0 or 1.5 minutes). The uptake clearance ( $CL_u$ ) was calculated by dividing the uptake rate by the initial substrate concentration. Percent  $CL_u$  inhibition was calculated using the following equation:

$$\text{Inhibition \%} = \frac{CL_u \text{ in the absence of inhibitor} - CL_u \text{ in the presence of inhibitor}}{CL_u \text{ in the absence of inhibitor}} \times 100.$$

**Statistical Analyses.** The results are expressed as means  $\pm$  S.D. in the text and tables. To test for statistically significant differences in  $C_{max}$ ,  $AUC_{(0-24 h)}$ , and  $CL_R$  among treatments (predose control, PROB, FSM, and PROB with FSM) in the clinical study, repeated measures one-way ANOVA was performed. When the  $F$  ratio showed that there were significant differences among treatments, the Tukey's method of multiple comparisons was used to determine which treatments



**Fig. 1.** Effect of 1000 mg PROB and 40 mg FSM doses on plasma concentration and  $CL_R$  of CP. The plasma concentration–time profiles of CPI (A) and CPIII (B) are shown as the mean and S.D. values obtained from 14 healthy subjects before dose (open squares) and after a single oral dose of PROB (open circles), FSM (open triangles), and coadministration of PROB and FSM (open diamonds). The  $CL_R$  of CPI (C) and CPIII (D) before and after the indicated treatment.

TABLE 1

Comparison of pharmacokinetic parameters of CPI and CPIII in healthy subjects after administration of PROB (1000 mg), FSM (40 mg), and coadministration of PROB and FSM

Data represent the means and S.D. from 9 to 14 subjects (n = 9–14).

Treatment	CPI			CPIII		
	$C_{max}$ (nM)	$AUC_{(0-24\text{ h})}$ (nM*h)	$CL_R$ (ml/min)	$C_{max}$ (nM)	$AUC_{(0-24\text{ h})}$ (nM*h)	$CL_R$ (ml/min)
Predose	1.34 ± 0.58	27.9 ± 11.5	18.6 ± 7.07	0.19 ± 0.08	3.54 ± 1.38	330 ± 210
PROB alone	1.98 ± 0.74***	41.1 ± 16.4***	16.3 ± 4.77	0.26 ± 0.11***	4.87 ± 2.22***	414 ± 238
Fold change <sup>a</sup>	1.54 ± 0.36	1.39 ± 0.21	0.97 ± 0.38	1.41 ± 0.22	1.34 ± 0.16	1.34 ± 0.53
FSM alone	1.44 ± 0.37	28.4 ± 7.27	19.0 ± 7.96	0.21 ± 0.07	3.54 ± 1.28	312 ± 191
Fold change <sup>a</sup>	1.15 ± 0.26	1.08 ± 0.27	1.10 ± 0.47	1.15 ± 0.32	1.02 ± 0.23	1.02 ± 0.50
PROB + FSM	1.96 ± 0.47**	41.3 ± 9.75**	21.2 ± 6.67	0.29 ± 0.09***	5.06 ± 2.03*	498 ± 247
Fold change <sup>a</sup>	1.60 ± 0.50	1.57 ± 0.41	1.16 ± 0.51	1.62 ± 0.46	1.45 ± 0.57	1.50 ± 0.69
Fold change <sup>b</sup>	1.42 ± 0.39	1.58 ± 0.25	1.23 ± 0.65	1.43 ± 0.26	1.34 ± 0.35	1.51 ± 0.41

<sup>a</sup>The fold-change values of  $C_{max}$ ,  $AUC_{(0-24\text{ h})}$ , and  $CL_R$  are calculated using predose levels.

<sup>b</sup>The fold-change values of  $C_{max}$ ,  $AUC_{(0-24\text{ h})}$ , and  $CL_R$  are calculated using FSM alone.

\* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ , statistically significant difference compared with the predose control.

differ. All statistical analyses were carried out using GraphPad Prism version 8 (GraphPad Software, San Diego, CA), and a  $P$  value of less than 0.05 was considered statistically significant.

## Results


**Effects of Administration of PROB on Plasma Levels of CPI and CPIII.** The effect of PROB on the pharmacokinetics of CPI and CPIII, endogenous biomarkers of OATP1B1 and OATP1B3, was evaluated in both plasma and urine after administration of PROB alone, furosemide alone, and probenecid in combination with FSM in 14 subjects. The systemic exposures and  $CL_R$  of CPI and CPIII are shown in Fig. 1 and Table 1. The table also presents the statistical comparisons for each treatment relative to predose baseline. The administration of 1000 mg PROB alone and in combination with 40 mg FSM significantly increased  $C_{max}$  and  $AUC_{(0-24\text{ h})}$  of CPI and CPIII compared with the baseline (1.34- to 1.62-fold) (Table 1) ( $P < 0.05$ ). In contrast, the  $C_{max}$  and  $AUC_{(0-24\text{ h})}$  values were similar to the basal levels after administration of FSM alone (1.02- to 1.15-fold). These changes were not statistically significant ( $P > 0.05$ ). Moreover, the  $CL_R$  values of CPI and CPIII were not significantly changed in the presence of PROB compared with the baseline control, although the mean  $CL_R$  of CPIII increased by 1.34- to 1.50-fold (Table 1) ( $P > 0.05$ ), indicating no significant inhibition of the  $CL_R$  of CPI and CPIII during PROB treatments.

**Effects of Administration of PROB on Plasma Levels of HDA and TDA.** We also examined the effects of a single dose of 1000 mg PROB, either alone or in combination with 40 mg FSM, on the plasma concentrations of HDA and TDA in healthy subjects. Figure 2 shows the arithmetic mean plasma concentrations ± S.D. of HDA and TDA before the first dose of the study (predose) and during the three treatment phases. Table 2 shows the arithmetic mean ± S.D. of  $C_{max}$ ,  $AUC_{(0-24\text{ h})}$ , and the ratios of HDA and TDA. Although pretreatment with PROB alone did not significantly alter the  $C_{max}$  or  $AUC_{(0-24\text{ h})}$  of HDA and TDA compared with baseline (0.73- to 1.02-fold) (Table 1) ( $P > 0.05$ ), the coadministration of PROB with FSM significantly increased the systemic exposures of HDA and TDA by 1.62- and 2.27-fold compared with FSM alone (Table 2) ( $P < 0.05$ ). In addition, administration of a single oral dose of 40 mg FSM alone resulted in less than a 1% change in the  $C_{max}$  and  $AUC_{(0-24\text{ h})}$  of HDA compared with the baseline controls ( $P > 0.05$ ). In contrast, administration of FSM alone significantly decreased the  $C_{max}$  and  $AUC_{(0-24\text{ h})}$  of TDA compared with the baseline (0.61- and 0.76-fold, respectively) (Table 2) ( $P < 0.05$ ). Unfortunately, the amount of HDA and TDA excreted in the urine were not analyzed in this study.

**Characterization of PROB as an In Vitro Inhibitor of OATP1B1 and OATP1B3.** To evaluate the inhibitory effect of PROB on

OATP1B1 and OATP1B3, the uptake of 0.2 μM CPI (OATP1B1 and OATP1B3), 1 μM [<sup>3</sup>H]E17βG (OATP1B1), and 0.1 μM [<sup>3</sup>H]CCK-8 (OATP1B3) was measured in the presence of increasing concentrations of PROB in transporter-expressing cells. As shown in Fig. 3, PROB inhibited OATP1B1- and OATP1B3-mediated uptake of CPI in a concentration-dependent manner, with  $IC_{50}$  values of 167 ± 42.0 and 76.0 ± 17.2 μM, respectively (Table 4). PROB showed very similar inhibition on the uptake of [<sup>3</sup>H]E17βG and [<sup>3</sup>H]CCK-8 by OATP1B1 and OATP1B3, respectively ( $IC_{50}$  values for OATP1B1 and OATP1B3 were 121 ± 7.5 and 147 ± 37.5 μM, respectively).

**Effect of PROB on CP Uptake in Hepatocytes.** To further demonstrate the inhibition of PROB on the hepatic elimination of CP,



**Fig. 2.** Effect of 1000 mg PROB and 40 mg FSM doses on plasma concentration of HDA and TDA. The plasma concentration–time profiles of HDA (A) and TDA (B) are shown as the mean and S.D. values obtained from 14 healthy subjects before dose (open squares) and after a single oral dose of PROB (open circles), FSM (open triangles), and coadministration of PROB and FSM (open diamonds).

TABLE 2

Comparison of pharmacokinetic parameters of HDA and TDA in healthy subjects after administration of PROB (1000 mg), FSM (40 mg), and coadministration of PROB and FSM

Data represent the means and S.D. from 14 subjects (n = 14).

Treatment	HDA		TDA	
	$C_{max}$ (nM)	$AUC_{(0-24\ h)}$ (nM*h)	$C_{max}$ (nM)	$AUC_{(0-24\ h)}$ (nM*h)
Predose	61.2 ± 34.2	884 ± 419	63.6 ± 47.9	787 ± 551
PROB alone	60.1 ± 48.7	891 ± 444	45.4 ± 34.3	693 ± 415
Fold change <sup>a</sup>	0.96 ± 0.29	1.02 ± 0.18	0.73 ± 0.25	0.90 ± 0.20
FSM alone	63.2 ± 52.7	890 ± 592	42.7 ± 52.8***	623 ± 585***
Fold change <sup>a</sup>	1.00 ± 0.34	0.99 ± 0.21	0.61 ± 0.17	0.76 ± 0.14
PROB + FSM	123 ± 65.7**	1404 ± 540***	72.5 ± 52.9	919 ± 516*
Fold change <sup>a</sup>	2.20 ± 1.26	1.66 ± 0.45	1.17 ± 0.49	1.22 ± 0.26
Fold change <sup>b</sup>	2.27 ± 1.06	1.71 ± 0.43	1.87 ± 0.86	1.62 ± 0.40

<sup>a</sup>The fold-change values of  $C_{max}$  and  $AUC_{(0-24\ h)}$  are calculated using predose levels.

<sup>b</sup>The fold-change values of  $C_{max}$  and  $AUC_{(0-24\ h)}$  are calculated using FSM alone.

\* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ , statistically significant difference compared with the predose control.


the uptake of 1  $\mu\text{M}$  CPI and CPIII in human hepatocytes was assessed. Figure 4 and Table 3 show the uptake of 1  $\mu\text{M}$  CPI and CPIII into human hepatocytes in suspension in the presence of increasing concentrations of PROB. The CPI and CPIII influx was sensitive to the presence of PROB, as a PROB concentration of 100  $\mu\text{M}$  reduced CPI and CPIII uptake by 12% and 33%, respectively, of the control. As the PROB concentration was raised, CPI and CPIII influx was further reduced so that at 1000  $\mu\text{M}$  PROB, uptake has been inhibited by 56% (Table 3). The uptake of the reference substrate RSV (1  $\mu\text{M}$ ) also appeared to be inhibited to a similar extent by PROB in hepatocytes (Fig. 4; Table 3).

### Discussion


The role and importance of OATP1B1 and OATP1B3 in the transport of drugs across the basolateral membranes of the hepatocytes are well recognized. It is within the last decade that the inhibition potential of a new molecular entity toward OATP1B transporter proteins has begun to be evaluated during drug discovery and development (Giacomini et al., 2010; Tweedie et al., 2013; Lee et al., 2017). For 34 small molecular drugs approved by the FDA in 2017, OATP1B together with CYP3A4 played a significant role in mediating more than half of the drug interactions with  $AUC$  changes  $\geq 5$ -fold (Yu et al., 2019). Moreover, it has become increasingly evident that endogenous biomarkers of OATP1B, for example CPI, can be used to evaluate complex DDIs for better DDI risk assessment and management based on a mechanistic understanding in healthy subjects and patients (Suzuki et al., 2019; Jones et al., 2020; Mori et al., 2020). The described findings suggest that this is the case for PROB. This drug is regarded as a clinically efficient inhibitor of OAT1 and OAT3 since the benzoic acid derivative is known to compete for active renal tubular secretion primarily with acidic drugs (Maeda et al., 2014). Moreover, PROB has recently been identified as a weak inhibitor of the OATP1B-mediated transport of organic anions using a heterologous expression system (Hirano et al., 2006; Matsushima et al., 2008; Izumi et al., 2013, 2016). Accordingly, the in vivo and in vitro inhibition potential of PROB toward OATP1B1 and OATP1B3 was investigated using the endogenous probes CPI, CPIII, HDA, and TDA.

For the first time, the results of this study showed a significant increase in plasma concentration of CPI and CPIII [ $C_{max}$  and  $AUC_{(0-24\ h)}$ ], endogenous probes of OATP1B1 and OATP1B3, in humans during PROB treatments compared with baseline levels (1.34- to 1.62-fold) (Fig. 1; Table 1). These findings suggest that PROB inhibited the elimination of the endogenous substrates of OATP1B, and this might be explained by an inhibitory effect of PROB on renal and/or hepatic

transporters, as CPI and CPIII are excreted not only in bile but also in the urine (Wolkoff et al., 1976; Lai et al., 2016; Shen et al., 2016). However, the renal clearance of CP was not significantly decreased by PROB (Table 1). When the plasma and urine findings are taken together, this suggests that the interaction occurs in the liver but not in the kidney. This is not unexpected, because CPI and CPIII are not substrates for OAT1 and OAT3 (Bednarczyk and Boiselle, 2016; Shen et al., 2017; Kunze et al., 2018), although OAT1 and OAT3 are capable of transporting small anionic endogenous substrates (e.g., *p*-aminohippurate and estrone-3-sulfate) and xenobiotics (e.g., adefovir, benzylpenicillin, and FSM) (Inui et al., 2000; Shen et al., 2019b). By contrast, OATP1B1 and



**Fig. 3.** Inhibitory effect of PROB on OATP1B1- (A) and OATP1B3-mediated transport of CPI and prototypical probe substrates. Uptake of 0.2  $\mu\text{M}$  CPI, 1  $\mu\text{M}$  E17 $\beta$ G (OATP1B1), and 1  $\mu\text{M}$  CCK-8 (OATP1B3) was assessed in stably transfected OATP1B1- and OATP1B3-HEK cells over 2 minutes in the absence (control) or presence of PROB at the indicated concentrations. Uptake changes are normalized to transporter-mediated uptake in the absence of PROB. Results represent the mean and S.D. from three independent determinations.



**Fig. 4.** Inhibitory effect of PROB on the uptake of CPI (A), CPIII (B), and RSV (C) into human hepatocytes. Open circles, squares, triangles and diamonds, and closed circles represent the uptake of 1  $\mu\text{M}$  substrate alone with 100  $\mu\text{M}$  PROB, 300  $\mu\text{M}$  PROB, 1000  $\mu\text{M}$  PROB, and 200  $\mu\text{M}$  RIF SV, respectively. Results represent the mean and S.D. from three independent determinations.

OATP1B3 were found to effectively mediate the uptake of CPI and CPIII, which is consistent with the anionic nature of these endogenous compounds (Bednarczyk and Boisselle, 2016; Shen et al., 2016, 2017). Very recently, Wiebe et al. (2020) investigated the effects of four commonly employed drug transporter inhibitors on cocktail drug pharmacokinetics. PROB treatment increased  $C_{\text{max}}$  and  $AUC$  of RSV by 328% and 123%, respectively. Although PROB decreased RSV  $CL_R$  by 78%, its inhibition of renal clearance alone could not explain the pronounced  $AUC$  increase since approximately 28% of total body clearance of RSV was via the renal route. In line with the clinical data, a monkey transporter-mediated DDI study showed that PROB increased the  $AUC$  of RSV and pitavastatin by 2.6- and 2.1-fold, respectively (Kosa et al., 2018). In addition, PROB significantly inhibited the uptake of RSV and pitavastatin in monkey hepatocytes (Kosa et al., 2018). Therefore, the interaction between PROB and statin might be attributed to the decreased hepatic uptake-mediated clearance by monkey OATP1B. Likewise, PROB had an impact on the OATP1B-mediated uptake clearance of fexofenadine in humans since its inhibition of renal organic anion transporter alone could not explain the noted  $AUC$  change (Yasui-Furukori et al., 2005; Liu et al., 2008). However, we cannot rule out that the synthesis of CPI and CPIII may be altered by PROB treatment. Interestingly, the mean renal clearance of CPIII but not CPI was increased by PROB, although the increase was not statistically significant compared with the predose level (1.34- and 1.50-fold) (Table 1) ( $P > 0.05$ ). Previously, we reported that CPIII but not CPI

was subject to active tubular secretion in the kidney of monkeys and humans (Lai et al., 2016; Shen et al., 2016). One possible explanation for the unexpected increase of CPIII  $CL_R$  is that PROB stimulated the function of transporter(s) responsible for the tubular secretion of CPIII. Such stimulation has been observed with many transporters such as multidrug resistance-associated protein (MRP) 2 (Gilibili et al., 2017). In addition, it is possible that CPIII undergoes both renal tubular secretion and reabsorption, and PROB inhibited the renal reabsorption of CPIII, resulting in the increased net renal excretion clearance of CPIII. Future work needs to be done to confirm the hypotheses. Servais et al. (2006) previously studied urinary coproporphyrin excretion in rats after inhibition of transporters by PROB and in mutant transport-deficient rats, in which MRP2 is lacking. Total urinary coproporphyrin excretion was similar in mutant transport-deficient rats and in normal rats with or without treatment by PROB, but relative urinary CPI excretion was increased in the mutant rats. Moreover, PROB is a weak inhibitor of MRP2 that is expressed on the apical surface of renal proximal tubule epithelial cells and hepatocytes. CPI and CPIII are substrates for MRP2 (Gilibili et al., 2017; Kunze et al., 2018), and the urinary CPI/(CPI + CPIII) ratio was proposed as a surrogate for MRP2 activity (Benz-de Bretagne et al., 2011, 2014). Unchanged  $CL_R$  values of CPI in the subjects administered with PROB suggested that PROB did not affect MRP2 activity in the kidney in vivo (Table 1). The increased  $CL_R$  of CPIII requires further study to identify the transporters responsible for the renal disposition of CPIII for better assessment of in vivo inhibition

TABLE 3

The hepatic uptake clearance of CP incubated in the presence and absence of PROB (100, 300, and 1000  $\mu\text{M}$ )

The hepatic uptake clearance and percent uptake clearance inhibition were determined using the data in Fig. 4.

Compound	Hepatocyte Uptake Clearance ( $\mu\text{L}/\text{min}$ per $10^6$ Cells) (%Inhibition)				
	Control	100 $\mu\text{M}$ PROB	300 $\mu\text{M}$ PROB	1000 $\mu\text{M}$ PROB	200 $\mu\text{M}$ RIF SV
CPI	2.1	1.9 (12%)	1.2 (42%)	0.93 (56%)	0.40 (81%)
CPIII	4.3	2.9 (33%)	2.0 (53%)	1.9 (56%)	0.61 (86%)
RSV	10.8	7.2 (34%)	4.9 (55%)	2.9 (73%)	0.73 (93%)

TABLE 4  
Prediction of OATP1B-mediated DDIs for PROB and FSM using *R*-value and endogenous biomarker methods

Treatment	Drug	$C_{\max}$ ( $\mu\text{M}$ ) <sup>c</sup>	$f_u$ <sup>a</sup>	OATP1B1		OATP1B3		Observed Index $C_{\max}$ and <i>AUC</i> Fold Changes			
				$IC_{50}$ ( $\mu\text{M}$ ) <sup>b</sup>	<i>R</i> -Value	$IC_{50}$ ( $\mu\text{M}$ ) <sup>b</sup>	<i>R</i> -Value	CPI	CPIII	HDA	TDA
PROB alone	1000 mg PROB	436	0.062	167 ± 42.0	1.25	76.0 ± 17.2	1.55	1.54 ( $C_{\max}$ ) 1.39 ( <i>AUC</i> )	1.41 ( $C_{\max}$ ) 1.34 ( <i>AUC</i> )	0.96 ( $C_{\max}$ ) 1.02 ( <i>AUC</i> )	0.73 ( $C_{\max}$ ) 0.90 ( <i>AUC</i> )
FSM alone	40 mg FSM	4.1	0.023	30–300	1.00–1.01	>300	1.00	1.15 ( $C_{\max}$ ) 1.08 ( <i>AUC</i> )	1.15 ( $C_{\max}$ ) 1.02 ( <i>AUC</i> )	1.00 ( $C_{\max}$ ) 0.99 ( <i>AUC</i> )	0.61 ( $C_{\max}$ ) 0.76 ( <i>AUC</i> )
PROB + FSM	1000 mg PROB	325	0.062	167 ± 42.0	1.21	76.0 ± 17.2	1.46	1.60 ( $C_{\max}$ ) 1.42 ( <i>AUC</i> )	1.62 ( $C_{\max}$ ) 1.45 ( <i>AUC</i> )	2.20 ( $C_{\max}$ ) 1.66 ( <i>AUC</i> )	1.17 ( $C_{\max}$ ) 1.22 ( <i>AUC</i> )

<sup>a</sup> $f_u$ , unbound fraction.

<sup>c</sup>Data obtained from Shen et al. (2019a).

<sup>b</sup>Concentrations required to inhibit OATP1B1- and OATP1B3-mediated transport by 50% obtained from the present study and previously reported (Ebner et al., 2015).

potential of an investigational drug using CPIII. Taken together, administration of PROB at a therapeutic dose can cause clinically relevant OATP1B inhibition.

In agreement with clinical findings, the  $IC_{50}$  of PROB in OATP1B1- and OATP1B3-HEK cells using CPI as a substrate is determined to be 167 and 76.0  $\mu\text{M}$ , respectively, in this study (Fig. 3). This is consistent with recent reports employing dichlorofluorescein, bromosulphthalein, E17 $\beta$ G, estrone-3-sulfate, pitavastatin, and fexofenadine as OATP1B1 and OATP1B3 substrates (Hirano et al., 2006; Matsushima et al., 2008; Izumi et al., 2013, 2016), in which the  $IC_{50}$  of PROB ranged from 39.8 to 227  $\mu\text{M}$ . Furthermore, 100, 300, and 1000  $\mu\text{M}$  PROB reduced the uptake of CPI, CPIII, and RSV in human hepatocytes in a concentration-dependent manner, with maximum inhibition by 56%, 56%, and 73%, respectively (Fig. 4; Table 3). The results suggested that PROB is a weak in vitro inhibitor of OATP1B1 and OATP1B3. However, because the free maximum concentration at the inlet to the liver ( $I_{\text{in,max,u}}$ ) after 1000 mg PROB administration is large [34.6–41.5  $\mu\text{M}$ , estimated by using the pharmacokinetic parameters reported previously (Shen et al., 2019a)], the predicted *AUC* ratio of a substrate drug of OATP1B in the presence and absence of PROB (*R*-value) is 1.46–1.55 using the  $IC_{50}$  value of 76.0  $\mu\text{M}$  determined using CPI as a probe substrate ( $R\text{-value} = 1 + I_{\text{in,max,u}}/IC_{50}$ ), suggesting an in vivo inhibition potential of PROB with OATP1B (Table 4). These *R*-values, estimated using the method recommended in the guidance from the EMA and FDA (<https://www.fda.gov/media/134582/download>; [https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-investigation-drug-interactions-revision-1\\_en.pdf](https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-investigation-drug-interactions-revision-1_en.pdf)), are consistent with the observed  $C_{\max}$  and *AUC* fold changes of endogenous probes CPI and CPIII. It is critical to include a sufficient number of subjects in a clinical study to detect weak drug transporter interaction. Barnett et al. (2018) demonstrated the sensitivity of CPI to identifying weak OATP1B inhibitors in an adequately powered clinical study using model-based simulations and power calculations. The analysis showed the ability to identify a weak but clinically relevant OATP1B DDI using CPI as a probe (The ratio of victim drug area under the curve in the presence of the interacting drug relative to the control (*AUCR*) > 1.25 cutoff). Use of criterion of  $\alpha = 0.01$  and power of 0.8 required a sample size of 15 subjects (Barnett et al., 2018). This conclusion was further confirmed by CPI clinical data with several weak OATP1B inhibitors (Kunze et al., 2018; Liu et al., 2018). The clinical studies with 14 and 13 subjects indicated that CPI concentration changes were predictive for a weak clinically observed DDI in which CPI *AUC* increases of 1.6- and 1.4-fold were comparable with those observed for statins as victim drugs (Kunze et al., 2018; Liu et al., 2018). Therefore, we believe that our data generated with 14 subjects is sufficient to support the conclusions made in this study.

To quantify the response of HDA and TDA, two other endogenous probes of OATP1B (Yee et al., 2016; Shen et al., 2017; Yee et al., 2019),

to PROB pretreatments in vivo, we measured the plasma HDA and TDA concentrations. The variations in plasma concentrations of HDA and TDA were larger than that of CPI and CPIII. The concentration-time profiles for HDA and TDA appear different between the PROB alone and PROB coadministration with FSM groups. Whereas the administration of PROB in combination with FSM significantly increased the  $AUC_{(0-24\text{ h})}$  of HDA and TDA by 1.71- and 1.62-fold, respectively, compared with FSM alone, the administration with PROB alone did not significantly alter the  $AUC_{(0-24\text{ h})}$  of HDA and TDA compared with prestudy levels (1.02- and 0.90-fold, respectively) (Fig. 2; Table 2). It is unclear why the PROB pretreatments show different effects on the plasma HDA and TDA levels. Unfortunately, the  $CL_R$  of HDA and TDA could not be determined in the urine. In addition, the treatment with FSM alone significantly decreased the  $AUC_{(0-24\text{ h})}$  of TDA compared with the predose control (0.76-fold) (Table 2). We cannot rule out that the synthesis of the biomarkers may be affected by PROB and/or FSM. These results suggest that plasma HDA and TDA levels may not be good surrogate endogenous probes for weak OATP1B inhibition.

In conclusion, for the first time, our results demonstrate the weak inhibitory effect on OATP1B1 and OATP1B3 by PROB in vivo. Considering the fact that PROB is an index inhibitor for clinical OAT1/3 DDI study, these findings provide an explanation for the heretofore unknown mechanism responsible for the inhibition caused by PROB.

#### Acknowledgments

The authors acknowledge the following scientists for assistance with the clinical and preclinical studies: Susan Lubin and Erika Panfen.

#### Authorship Contributions

*Participated in research design:* Zhang, Holenarsipur, Zeng, Mariappan, Sinz, Shen.

*Conducted experiments:* Zhang, Kandoussi.

*Contributed new reagents or analytic tools:* Zhang, Kandoussi, Shen.

*Performed data analysis:* Zhang, Shen.

*Wrote or contributed to the writing of the manuscript:* Zhang, Sinz, Shen.

#### References

- Barnett S, Ogungbenro K, Ménochet K, Shen H, Humphreys WG, and Galetin A (2019) Comprehensive evaluation of the utility of 20 endogenous molecules as biomarkers of OATP1B inhibition compared with rosuvastatin and coproporphyrin I. *J Pharmacol Exp Ther* **368**:125–135.
- Barnett S, Ogungbenro K, Ménochet K, Shen H, Lai Y, Humphreys WG, and Galetin A (2018) Gaining mechanistic insight into coproporphyrin I as endogenous biomarker for OATP1B-mediated drug-drug interactions using population pharmacokinetic modeling and simulation. *Clin Pharmacol Ther* **104**:564–574.
- Bednarczyk D and Boiselle C (2016) Organic anion transporting polypeptide (OATP)-mediated transport of coproporphyrins I and III. *Xenobiotica* **46**:457–466.
- Benz-de Bretagne I, Respaud R, Vourc'h P, Halimi JM, Caille A, Hulot JS, Andres CR, and Le Guellec C (2011) Urinary elimination of coproporphyrins is dependent on ABCB2 polymorphisms and represents a potential biomarker of MRP2 activity in humans. *J Biomed Biotechnol* **2011**:498757.

- Benz-de Bretagne I, Zahr N, Le Gouge A, Hulot JS, Houillier C, Hoang-Xuan K, Gyan E, Lissandre S, Choquet S, and Le Guellec C (2014) Urinary coproporphyrin I/(I + III) ratio as a surrogate for MRP2 or other transporter activities involved in methotrexate clearance. *Br J Clin Pharmacol* **78**:329–342.
- Beringer PM, Kriengkauykiat J, Zhang X, Hidayat L, Liu S, Louie S, Synold T, Burckart GJ, Rao PA, Shapiro B, et al. (2008) Lack of effect of P-glycoprotein inhibition on renal clearance of dioxacinil in patients with cystic fibrosis. *Pharmacotherapy* **28**:883–894.
- Chu X, Liao M, Shen H, Yoshida K, Zur AA, Arya V, Galetin A, Giacomini KM, Hanna I, Kusuhabara H, et al. International Transporter Consortium (2018) Clinical probes and endogenous biomarkers as substrates for transporter drug-drug interaction evaluation: perspectives from the international transporter consortium. *Clin Pharmacol Ther* **104**:836–864.
- Cunningham RF, Israeli ZH, and Dayton PG (1981) Clinical pharmacokinetics of probenecid. *Clin Pharmacokinet* **6**:135–151.
- Ebner T, Ishiguro N, and Taub ME (2015) The use of transporter probe drug cocktails for the assessment of transporter-based drug-drug interactions in a clinical setting-proposal of a four component transporter cocktail. *J Pharm Sci* **104**:3220–3228.
- Gewirtz DA, Plotkin JH, and Randolph JK (1984) Interaction of probenecid with methotrexate transport and release in the isolated rat hepatocyte in suspension. *Cancer Res* **44**:3846–3850.
- Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, Hillgren KM, et al. International Transporter Consortium (2010) Membrane transporters in drug development. *Nat Rev Drug Discov* **9**:215–236.
- Gilibili RR, Chatterjee S, Bagul P, Masure KW, Murali BV, Mariappan TT, Mandekar S, and Lai Y (2017) Coproporphyrin-I: a fluorescent, endogenous optimal probe substrate for ABC2 (MRP2) suitable for vesicle-based MRP2 inhibition assay. *Drug Metab Dispos* **45**:604–611.
- Han YH, Busler D, Hong Y, Tian Y, Chen C, and Rodrigues AD (2010) Transporter studies with the 3-O-sulfate conjugate of 17alpha-ethinylestradiol: assessment of human liver drug transporters. *Drug Metab Dispos* **38**:1072–1082.
- Hirano M, Maeda K, Shitara Y, and Sugiyama Y (2006) Drug-drug interaction between pitavastatin and various drugs via OATP1B1. *Drug Metab Dispos* **34**:1229–1236.
- Inotsume N, Nishimura M, Nakano M, Fujiyama S, and Sato T (1990) The inhibitory effect of probenecid on renal excretion of famotidine in young, healthy volunteers. *J Clin Pharmacol* **30**:50–56.
- Inui KI, Masuda S, and Saito H (2000) Cellular and molecular aspects of drug transport in the kidney. *Kidney Int* **58**:944–958.
- Izumi S, Nozaki Y, Komori T, Maeda K, Takenaka O, Kusano K, Yoshimura T, Kusuhabara H, and Sugiyama Y (2013) Substrate-dependent inhibition of organic anion transporting polypeptide 1B1: comparative analysis with prototypical probe substrates estradiol-17 $\beta$ -glucuronide, estrone-3-sulfate, and sulfobromophthalein. *Drug Metab Dispos* **41**:1859–1866.
- Izumi S, Nozaki Y, Komori T, Takenaka O, Maeda K, Kusuhabara H, and Sugiyama Y (2016) Investigation of fluorescein derivatives as substrates of organic anion transporting polypeptide (OATP) 1B1 to develop sensitive fluorescence-based OATP1B1 inhibition assays. *Mol Pharm* **13**:438–448.
- Jones NS, Yoshida K, Salphati L, Kenny JR, Durk MR, and Chinn LW (2020) Complex DDI by fenebrutinib and the use of transporter endogenous biomarkers to elucidate the mechanism of DDI. *Clin Pharmacol Ther* **107**:269–277.
- Kandoussi H, Zeng J, Shah K, Paterson P, Santockyte R, Kadiyala P, Shen H, Shipkova P, Langish R, Burrell R, et al. (2018) UHPLC-MS/MS bioanalysis of human plasma coproporphyrins as potential biomarkers for organic anion-transporting polypeptide-mediated drug interactions. *Bioanalysis* **10**:633–644.
- Kosa RE, Lazzaro S, Bi YA, Tierney B, Gates D, Modi S, Costales C, Rodrigues AD, Tremaine LM, and Varma MV (2018) Simultaneous assessment of transporter-mediated drug-drug interactions using a probe drug cocktail in cynomolgus monkey. *Drug Metab Dispos* **46**:1179–1189.
- Kunze A, Ediage EN, Dillen L, Monshouwer M, and Snoeys J (2018) Clinical investigation of coproporphyrins as sensitive biomarkers to predict mild to strong OATP1B-mediated drug-drug interactions. *Clin Pharmacokinet* **57**:1559–1570.
- Lai Y, Mandekar S, Shen H, Holenarsipur VK, Langish R, Rajanna P, Murugesan S, Gaud N, Selvam S, Date O, et al. (2016) Coproporphyrins in plasma and urine can be appropriate clinical biomarkers to recapitulate drug-drug interactions mediated by organic anion transporting polypeptide inhibition. *J Pharmacol Exp Ther* **358**:397–404.
- Lappin G, Shishikura Y, Jochemsen R, Weaver RJ, Gesson C, Houston B, Oosterhuis B, Bjerrum OJ, Rowland M, and Garner C (2010) Pharmacokinetics of fexofenadine: evaluation of a microdose and assessment of absolute oral bioavailability. *Eur J Pharm Sci* **40**:125–131.
- Lee SC, Arya V, Yang X, Volpe DA, and Zhang L (2017) Evaluation of transporters in drug development: current status and contemporary issues. *Adv Drug Deliv Rev* **116**:100–118.
- Liu L, Cheeti S, Yoshida K, Choo E, Chen E, Chen B, Gates M, Singel S, Morley R, Ware J, et al. (2018) Effect of OATP1B1/1B3 inhibitor GDC-0810 on the pharmacokinetics of pravastatin and coproporphyrin I/III in healthy female subjects. *J Clin Pharmacol* **58**:1427–1435.
- Liu S, Beringer PM, Hidayat L, Rao AP, Louie S, Burckart GJ, and Shapiro B (2008) Probenecid, but not cystic fibrosis, alters the total and renal clearance of fexofenadine. *J Clin Pharmacol* **48**:957–965.
- Maeda K, Tian Y, Fujita T, Ikeda Y, Kumagai Y, Kondo T, Tanabe K, Nakayama H, Horita S, Kusuhabara H, et al. (2014) Inhibitory effects of p-aminohippurate and probenecid on the renal clearance of adefovir and benzylpenicillin as probe drugs for organic anion transporter (OAT) 1 and OAT3 in humans. *Eur J Pharm Sci* **59**:94–103.
- Matsushima S, Maeda K, Ishiguro N, Igarashi T, and Sugiyama Y (2008) Investigation of the inhibitory effects of various drugs on the hepatic uptake of fexofenadine in humans. *Drug Metab Dispos* **36**:663–669.
- Molimard M, Diquet B, and Benedetti MS (2004) Comparison of pharmacokinetics and metabolism of desloratadine, fexofenadine, levocetirizine and mizolastine in humans. *Fundam Clin Pharmacol* **18**:399–411.
- Mori D, Ishida H, Mizuno T, Kusumoto S, Kondo Y, Izumi S, Nakata G, Nozaki Y, Maeda K, Sasaki Y, et al. (2020) Alteration in the plasma concentrations of endogenous OATP1B1-biomarkers in non-small cell lung cancer patients treated with paclitaxel. *Drug Metab Dispos* **48**:387–394.
- Müller F, Sharma A, König J, and Fromm MF (2018) Biomarkers for in vivo assessment of transporter function. *Pharmacol Rev* **70**:246–277.
- Panfen E, Chen W, Zhang Y, Sinz M, Marathe P, Gan J, and Shen H (2019) Enhanced and persistent inhibition of organic cation transporter 1 activity by preincubation of cyclosporine A. *Drug Metab Dispos* **47**:1352–1360.
- Pitkin D, Dubb J, Actor P, Alexander F, Ehrlich S, Familiar R, and Stote R (1981) Kinetics and renal handling of cefonicid. *Clin Pharmacol Ther* **30**:587–593.
- Rodrigues AD, Taskar KS, Kusuhabara H, and Sugiyama Y (2018) Endogenous probes for drug transporters: balancing vision with reality. *Clin Pharmacol Ther* **103**:434–448.
- Santockyte R, Kandoussi H, Chen W, Zheng N, Venkatarangan L, Gan J, Shen H, Bonacorsi SJ, Easter J, Burrell R, et al. (2018) LC-MS/MS bioanalysis of plasma 1, 14-tetradecanedioic acid and 1, 16-hexadecanedioic acid as candidate biomarkers for organic anion-transporting polypeptide mediated drug-drug interactions. *Bioanalysis* **10**:1473–1485.
- Servais A, Lechat P, Zahr N, Urien S, Aymard G, Jaudon MC, Deray G, and Isnard Bagnis C (2006) Tubular transporters and clearance of adefovir. *Eur J Pharmacol* **540**:168–174.
- Shen H (2018) A pharmaceutical industry perspective on transporter and CYP-mediated drug-drug interactions: kidney transporter biomarkers. *Bioanalysis* **10**:625–631.
- Shen H, Chen W, Drexler DM, Mandekar S, Holenarsipur VK, Shields EE, Langish R, Sidik K, Gan J, Humphreys WG, et al. (2017) Comparative evaluation of plasma bile acids, dehydroepiandrosterone sulfate, hexadecanedioate, and tetradecanedioate with coproporphyrins I and III as markers of OATP inhibition in healthy subjects. *Drug Metab Dispos* **45**:908–919.
- Shen H, Christopher L, Lai Y, Gong J, Kandoussi H, Garonik S, Perera W, Garimella T, and Humphreys WG (2018) Further studies to support the use of coproporphyrin I and III as novel clinical biomarkers for evaluating the potential for organic anion transporting polypeptide 1B1 and OATP1B3 inhibition. *Drug Metab Dispos* **46**:1075–1082.
- Shen H, Dai J, Liu T, Cheng Y, Chen W, Freedman C, Zhang Y, Humphreys WG, Marathe P, and Lai Y (2016) Coproporphyrins I and III as functional markers of OATP1B activity: in vitro and in vivo evaluation in preclinical species. *J Pharmacol Exp Ther* **357**:382–393.
- Shen H, Holenarsipur VK, Mariappan TT, Drexler DM, Cantone JL, Rajanna P, Singh Gautam S, Zhang Y, Gan J, Shipkova PA, et al. (2019a) Evidence for the validity of pyridoxic acid (PDA) as a plasma-based endogenous probe for OAT1 and OAT3 function in healthy subjects. *J Pharmacol Exp Ther* **368**:136–145.
- Shen H, Scialis RJ, and Lehman-McKeeman L (2019b) Xenobiotic transporters in the kidney: function and role in toxicity. *Semin Nephrol* **39**:159–175.
- Shen H, Yang Z, Mintier G, Han YH, Chen C, Balimane P, Jemal M, Zhao W, Zhang R, Kallipatti S, et al. (2013) Cynomolgus monkey as a potential model to assess drug interactions involving hepatic organic anion transporting polypeptides: in vitro, in vivo, and in vitro-to-in vivo extrapolation. *J Pharmacol Exp Ther* **344**:673–685.
- Suzuki Y, Ono H, Tanaka R, Sato F, Sato Y, Ohno K, Mimata H, and Itoh H (2019) Recovery of OATP1B activity after living kidney transplantation in patients with end-stage renal disease. *Pharm Res* **36**:59.
- Takehara I, Yoshikado T, Ishigame K, Mori D, Furihata KI, Watanabe N, Ando O, Maeda K, Sugiyama Y, and Kusuhabara H (2018) Comparative study of the dose-dependence of OATP1B inhibition by rifampicin using probe drugs and endogenous substrates in healthy volunteers. *Pharm Res* **35**:138.
- Tweedie DJ, Polli JW, Berglund EG, Huang SM, Zhang L, Poirier A, Chu X, and Feng B; International Transporter Consortium (2013) Transporter studies in drug development: experience to date and follow-up on decision trees from the International Transporter Consortium. *Clin Pharmacol Ther* **94**:113–125.
- Vlasses PH, Holbrook AM, Schrogie JJ, Rogers JD, Ferguson RK, and Abrams WB (1980) Effect of orally administered probenecid on the pharmacokinetics of cefoxitin. *Antimicrob Agents Chemother* **17**:847–855.
- Vree TB, van den Biggelaar-Marteau M, and Verwey-van Wissen CP (1995) Probenecid inhibits the renal clearance of frusemide and its acyl glucuronide. *Br J Clin Pharmacol* **39**:692–695.
- Welling PG, Dean S, Selen A, Kendall MJ, and Wise R (1979) Probenecid: an unexplained effect on cephalosporin pharmacology. *Br J Clin Pharmacol* **8**:491–495.
- Wiebe ST, Giessmann T, Hohl K, Schmidt-Gerets S, Huel E, Jambrecina A, Bader K, Ishiguro N, Taub ME, Sharma A, et al. (2020) Validation of a drug transporter probe cocktail using the prototypical inhibitors rifampin, probenecid, verapamil, and cimetidine. *Clin Pharmacokinet* DOI: 10.1007/s40262-020-00907-w [published ahead of print].
- Wolkoff AW, Wolpert E, Pascasio FN, and Arias IM (1976) Rotor's syndrome. A distinct inheritable pathophysiologic entity. *Am J Med* **60**:173–179.
- Yasui-Furukori N, Uno T, Sugawara K, and Tateishi T (2005) Different effects of three transporting inhibitors, verapamil, cimetidine, and probenecid, on fexofenadine pharmacokinetics. *Clin Pharmacol Ther* **77**:17–23.
- Yee SW, Giacomini MM, Hsueh CH, Weitz D, Liang X, Goswami S, Kinchen JM, Coelho A, Zur AA, Mertsch K, et al. (2016) Metabolomic and genome-wide association studies reveal potential endogenous biomarkers for OATP1B1. *Clin Pharmacol Ther* **100**:524–536.
- Yee SW, Giacomini MM, Shen H, Humphreys WG, Horng H, Brian W, Lai Y, Kroetz DL, and Giacomini KM (2019) Organic anion transporter polypeptide 1B1 polymorphism modulates the extent of drug-drug interaction and associated biomarker levels in healthy volunteers. *Clin Transl Sci* **12**:388–399.
- Yu J, Petrie ID, Levy RH, and Ragueneau-Majlessi I (2019) Mechanisms and clinical significance of pharmacokinetic-based drug-drug interactions with drugs approved by the U.S. Food and Drug Administration in 2017. *Drug Metab Dispos* **47**:135–144.
- Zhang Y, Panfen E, Fanher M, Sinz M, Marathe P, and Shen H (2019) Dissecting the contribution of OATP1B1 to hepatic uptake of statins using the OATP1B1 selective inhibitor estropiate. *Mol Pharm* **16**:2342–2353.

**Address correspondence to:** Dr. Hong Shen, Senior Principal Scientist, Department of Metabolism and Pharmacokinetics, Bristol Myers Squibb, Route 206 & Province Line Rd., Princeton, NJ 08543. E-mail: hong.shen1@bms.com