

The Inactivation of Human Aldehyde Oxidase 1 by Hydrogen Peroxide and Superoxide

Claudia Garrido and Silke Leimkühler

Institute of Biochemistry and Biology, Department of Molecular Enzymology, University of Potsdam, Potsdam, Germany

Received May 20, 2020; accepted June 21, 2021

ABSTRACT

Mammalian aldehyde oxidases (AOX) are molybdo-flavoenzymes of pharmacological and pathophysiologic relevance that are involved in phase I drug metabolism and, as a product of their enzymatic activity, are also involved in the generation of reactive oxygen species. So far, the physiologic role of aldehyde oxidase 1 in the human body remains unknown. The human enzyme hAOX1 is characterized by a broad substrate specificity, oxidizing aromatic/aliphatic aldehydes into their corresponding carboxylic acids, and hydroxylating various heteroaromatic rings. The enzyme uses oxygen as terminal electron acceptor to produce hydrogen peroxide and superoxide during turnover. Since hAOX1 and, in particular, some natural variants produce not only H₂O₂ but also high amounts of superoxide, we investigated the effect of both ROS molecules on the enzymatic activity of hAOX1 in more detail. We compared hAOX1 to the high-O₂⁻-producing natural variant L438V for their time-dependent inactivation with H₂O₂/O₂⁻ during substrate turnover. We show that the inactivation of the hAOX1 wild-

type enzyme is mainly based on the production of hydrogen peroxide, whereas for the variant L438V, both hydrogen peroxide and superoxide contribute to the time-dependent inactivation of the enzyme during turnover. Further, the level of inactivation was revealed to be substrate-dependent: using substrates with higher turnover numbers resulted in a faster inactivation of the enzymes. Analysis of the inactivation site of the enzyme identified a loss of the terminal sulfido ligand at the molybdenum active site by the produced ROS during turnover.

SIGNIFICANCE STATEMENT

This work characterizes the substrate-dependent inactivation of human aldehyde oxidase 1 under turnover by reactive oxygen species and identifies the site of inactivation. The role of ROS in the inhibition of human aldehyde oxidase 1 will have a high impact on future studies.

Introduction

Human aldehyde oxidase (hAOX1) is a cytosolic homodimeric molybdo-flavoenzyme that binds the molybdenum cofactor (Moco), 2 × [2Fe-2S] clusters, and FAD as prosthetic groups (Terao et al., 2020). The enzyme belongs to the xanthine oxidase (XO) family of molybdoenzymes (Hille et al., 2014). So far, no clear physiologic function has been described for hAOX1 (Garattini et al., 2007; Terao et al., 2016). hAOX1 is a phase I drug-metabolizing enzyme, and its broad substrate specificity and unpredictable interindividual variability still present a challenge to fully understand the role of this enzyme in the human body (Beedham, 2020). hAOX1-mediated metabolism has led to several failures in clinical trials and to the termination of drug discovery programs, raising the awareness for the importance of this metalloenzyme (Manevski et al., 2019).

Like XO, an enzyme involved in the catabolism of purines (Coughlan, 1980), eukaryotic AOX enzymes use oxygen as the terminal electron acceptor and produce reactive oxygen species (ROS) during turnover (Coelho et al., 2015; Foti et al., 2017). Mammalian XO, however, exists in two interconvertible forms, the oxidase form (XO), which uses O₂ as electron acceptor, and the dehydrogenase form (XDH), which uses NAD⁺ as electron acceptor (Enroth et al., 2000; Kuwabara

et al., 2003). In contrast, mammalian AOX enzymes solely exist in the oxidase form and cannot be converted to the dehydrogenase form using NAD⁺ as electron acceptor (Kurosaki et al., 2013). ROS represent prominent key molecules in physiologic and pathologic conditions in the cell (Oberley, 2002). When ROS achieve higher and unbalanced concentrations, they are a dangerous source of damage to several molecules within the cell. ROS can cause damage to the DNA, leading to an oncogenic effect, damage to lipids with consequent peroxidation, and damage to residues or cofactors of proteins, e.g., iron-sulfur clusters (Holmström and Finkel, 2014). So far, mainly XO had been implicated to generate significant amounts of O₂⁻ during the course of its catalytic activity with purine substrates (Terada et al., 1991b; Harris and Massey, 1997; Kundu et al., 2007; Lee et al., 2014).

In our previous studies, the hAOX1 enzyme was shown to produce superoxide radicals with a rate of around 10%, as compared with the amount of hydrogen peroxide produced during the overall reaction (Foti et al., 2017). This value, however, is lower than the reported rates of 16%–20% for bovine XO (Nishino and Tamura, 1991). Further, it was reported that significant alterations in the rate of superoxide anion production are present in human natural variants of hAOX1 based on single nucleotide polymorphisms (SNPs) (Foti et al., 2017). In particular, the SNP-based amino acid exchange L438V of hAOX1 in proximity to the isoalloxazine ring of the FAD cofactor resulted in an increased rate of superoxide radical production to 75%. This variant is therefore considered to be an overproducer of O₂⁻. Considering the high toxicity of superoxide in the cell, the hAOX1-L438V SNP variant is a considerable

This work was supported by the Deutsche Forschungsgemeinschaft [LE1171/8-3].
The authors declare no conflict of interest.
<https://dx.doi.org/10.1124/dmd.121.000549>

ABBREVIATIONS: AOX, aldehyde oxidase; DCPIP, 2,6-dichlorophenolindophenol; DCPIP₂, reduced DCPIP; FAD, flavin adenin dinucleotide; hAOX1, human aldehyde oxidase 1; KCN, potassium cyanide; Moco, molybdopterin cofactor; ROS, reactive oxygen species; SNP, single nucleotide polymorphism; SOD, superoxide dismutase; WT, wildtype; XDH, xanthine dehydrogenase; XO, xanthine oxidase.

candidate for pathologic roles within the human population. The fact that wild-type hAOX1 also produces a significant amount of H_2O_2 and $\text{O}_2^{\cdot-}$ radicals might therefore also be of pathophysiological interest (Kundu et al., 2007). Particularly, it is noticeable that hepatic hAOX1 is calculated to generate 24-fold larger amounts of $\text{O}_2^{\cdot-}$ than XO. These calculations are based on the relative levels of XO and AOX enzymatic activity in human liver (Krenitsky et al., 1972; Kundu et al., 2007). Thus, hAOX1 and other mammalian AOXs may represent significant sources of ROS in the cytosol of liver cells and other tissues and may play a critical role in ROS-mediated tissue injury under specific conditions (Hunt and Massey, 1992; Kundu et al., 2012). Oxidative damage has long been implicated in both the malignant phenotype and carcinogenesis (Oberley, 2002). A role of hAOX1 and XO in cancer has been assigned (Qiao et al., 2020). In a recent study on XO, it was shown that the block of ROS production slowed down tumor growth (Kusano et al., 2019). Therefore, the contribution and role of hAOX1 in ROS production that leads to cancer development needs further investigation in the future.

More than 95 years ago, Dixon (1925) showed that incubation of bovine milk XO with hydrogen peroxide resulted in an inactivation of the enzyme. This observation has been confirmed several times with XO and with the related XDH (Bergel and Bray, 1959; Betcher-Lange et al., 1979). Apart from nonspecific modes of inactivation, the various prosthetic groups in these enzymes have been suggested to be the targets for attack by hydrogen peroxide (Lynch and Fridovich, 1979; Terada et al., 1991a). However, the nature of the reaction involved has not been investigated in detail, nor has the relation between the sites of inactivation and of catalysis been clearly established. It has been concluded, however, that H_2O_2 treatment of the enzyme results in the desulfo form at the molybdenum active site (Betcher-Lange et al., 1979).

Since hAOX1 and, in particular, some variants produce not only H_2O_2 but also high amounts of superoxide, we investigated the effect of both ROS molecules on the enzymatic activity of hAOX1 in more detail. We show that the enzyme is inactivated by hydrogen peroxide or, when produced, also by superoxide during turnover, when the enzyme is in its reduced state. We identified the sulfido ligand at the active site as the target of inactivation, and the enzyme activity was restored by a chemical sulfuration procedure that restores the sulfido ligand at the Moco active site, which is essential for the activity of the enzyme.

Materials and Methods

Expression and Purification of hAOX1 and Variants. The hAOX1 wild-type protein and the L438V variant were expressed and purified as described previously, with minor modifications (Foti et al., 2016). The constructs pTHcohAOX1 (hAOX1 wild type) (Foti et al., 2016) and pTHcohAOX1-L438V (Foti et al., 2017) were transformed into *Escherichia coli* TP1000 ($\Delta mobAB$) cells (Palmer et al., 1996). For protein expression, *E. coli* cell cultures were grown at 30°C in lysogeny broth medium supplemented with 150 $\mu\text{g}/\text{ml}$ ampicillin, 1 mM sodium molybdate, and 20 μM isopropyl- β -D-thiogalactopyranoside. Cells were harvested by centrifugation after 24 hours of cell growth and resuspended in 50 mM sodium phosphate buffer, pH 8.0, containing 300 mM NaCl. After cell lysis, the crude extract containing hAOX1 was first purified using a nickel-nitrilotriacetic acid resin (QIAGEN GmbH, Hilden, Germany) and then subjected to chemical sulfuration before the final purification by a size-exclusion chromatography step using a Superdex 200 10/300 GL Column (GE Healthcare), as described previously (Foti et al., 2016).

Metal Quantification. Inductively coupled plasma optical emission spectroscopy with an Optima 2100 DV (PerkinElmer Life and Analytical Sciences, Waltham, MA) was used to measure the metal content (Neumann and Leimkühler, 2008). In total, 500 μl of purified hAOX1 (about 10 μM) and an equal volume of 65% nitric acid were mixed to wet ash the protein at 100°C overnight. The samples were diluted with 4 ml of water. The buffer used as reference was 50 mM Tris-HCl, 200 mM NaCl, EDTA 1 mM, pH 8.0. The detection was at

wavelengths of 203.845 nm, 202.031 nm, and 204.597 nm for molybdenum and 238.204 nm, 239.562 nm, and 259.939 nm for iron. A standard was used for calibration and quantification of the detected metals (Standard Solution XVI, Merck). The resulting mass concentrations were calculated and related as percentage of protein saturated with Moco and iron corresponding to the two [2Fe-2S] clusters.

Steady-State Kinetics. Steady-state enzyme kinetics were performed with purified 200 nM hAOX1 or 200 nM L438V in 50 mM Tris and 1 mM EDTA buffer, pH 8.0, at 25°C in a final volume of 500 μl . Phthalazine and benzaldehyde were used in a range of 5–200 μM , and for zoniporide, the concentrations used were from 75 μM to 600 μM . The electron acceptor was molecular oxygen in air-saturated EDTA-Tris buffer, pH 8.0. The oxidized product (benzoic acid, phenanthridone, or 2-oxo-zoniporide) was detected by UV absorbance (295 nm, 304 nm, or 315 nm, respectively). Reactions were monitored over a range of 60 seconds. Activities were calculated using the molar extinction coefficients of 1166 $\text{M}^{-1}\text{cm}^{-1}$ (at 304 nm) for phthalazine, 1321 $\text{M}^{-1}\text{cm}^{-1}$ (at 290 nm) for benzaldehyde, and 5775 $\text{M}^{-1}\text{cm}^{-1}$ (at 315 nm) for zoniporide. Mean values with standard deviations were obtained from at least three independent measurements. k_{cat} and K_M values were normalized to 100% molybdenum content. The kinetic constants were obtained using the Michaelis-Menten equation by nonlinear regression with the software Origin Pro 8.1G (Waltham, MA). The enzyme kinetics assays were performed on a Shimadzu UV-2401PC photometer at room temperature.

Inactivation of hAOX1 under Turnover Conditions. The inactivation of hAOX1 WT and the L438V variant by substrates was analyzed with each enzyme (20 μM) that was incubated in air-saturated 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, in the presence of 1.5 mM benzaldehyde, 1.5 mM zoniporide, or 1.5 mM phthalazine. A pH of 8.0 was used in our study, since the pH optimum of hAOX1 is at pH 8.0 (Foti et al., 2016). The enzyme activity was tested every 15 minutes by taking a 5- μl aliquot diluted into 500 μl of fresh in air-saturated 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, containing either 150 μM benzaldehyde, 150 μM phthalazine, or 150 μM zoniporide. In total, 2000 U/ml of catalase and/or SOD was included when required.

Activity Assays Using DCPIP as Electron Acceptor. Inactivation of hAOX1 WT and the L438V variant by substrates was monitored using 100 μM DCPIP as electron acceptor. The specific activities were measured by monitoring the product formation of reduced DCPIP, 2,6-dichlorophenolindophenol (DCPIP_H) at 600 nm using an extinction coefficient of 16100 $\text{M}^{-1}\text{cm}^{-1}$. The measurements were performed under aerobic conditions using air-saturated buffer and under anaerobic conditions using degassed buffer in an anaerobic chamber (Coy).

Reactivation of the Enzymes by Resulfuration. First, an aliquot of 1 ml of 20 μM enzyme with 150 μM benzaldehyde was prepared. The enzyme activity of the aliquot was measured at time point 0 and after 60 minutes of incubation with benzaldehyde and/or 2000 U/ml catalase. After 60 minutes, the assay mixture was exchanged into 50 mM KH_2PO_4 • KOH, 0.1 mM EDTA, pH 7.4, by using a PD-10 column.

Further, the enzyme concentration was measured by using the absorbance at 450 nm, and the activity of half of the reaction was measured and used as a non-sulfurated control. The remaining enzyme was used for chemical sulfuration and transferred to an anaerobic chamber (Coy). An adapted procedure to the chemical sulfuration protocol originally reported by Wahl and Rajagopalan (1982) was used. To 10 μM enzyme, 200 μl of 5 mM sodium dithionite, 1 mM sodium sulfite, and 12 μM methylviologen was added, and the mixture was incubated for 30 minutes on ice. Afterward, the buffer was exchanged into 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, using a PD-10 column. The activity was determined with benzaldehyde as substrate and compared with the activity of the nonsulfurated control.

Results

Steady-State Kinetics of hAOX1 and the Superoxide-Overproducing Variant L438V with Different Substrates A substrate-dependent inactivation has been reported previously for hAOX1 (Abbasi et al., 2019) and the XO/XDH enzymes (Dixon, 1925; Betcher-Lange et al., 1979). For chicken liver XDH, it had been suggested that the enzyme is inactivated by hydrogen peroxide that is produced during turnover (Betcher-Lange et al., 1979). Therefore, we wanted to

investigate the nature of the inactivation of hAOX1 by comparison of two enzyme variants, the purified hAOX1 wild-type enzyme and the natural variant L438V, which was shown to produce 75% superoxide, whereas the wild type only produces 10% superoxide during substrate turnover with oxygen as electron acceptor (Foti et al., 2017). Both enzymes were expressed in a heterologous system in *E. coli* (Foti et al., 2016). The goal was to analyze whether hydrogen peroxide or superoxide produced by the enzyme influence the substrate-dependent inactivation of the enzyme that has been observed under turnover conditions (Abbasi et al., 2019). First, we determined the steady-state kinetic constants of the two enzymes with three selected substrates for comparison: phthalazine as fast substrate, benzaldehyde as intermediate substrate, and zoniporide as slow substrate (Table 1). The kinetic constants of these substrates with oxygen as electron acceptor were not reported before for hAOX1 wild type and the L438V variant. The results show that both hAOX1 proteins have similar kinetic constants with the three substrates, which is consistent with previously published data (Foti et al., 2017) using a different substrate, and therefore both proteins can be directly compared in our study.

Wild-Type hAOX1 Is Inactivated by Hydrogen Peroxide, Whereas the L438V Variant Is Inactivated by Both Hydrogen Peroxide and Superoxide under Turnover Conditions. To analyze the substrate and time-dependent inactivation of hAOX1 wild type and the variant L438V, we monitored product formation over a time of 60 minutes with three different substrates (phthalazine, benzaldehyde, and zoniporide) using oxygen as terminal electron acceptor. Since oxygen is converted to hydrogen peroxide and superoxide at different rates by the two enzyme variants, we also included superoxide dismutase and catalase in the reaction mixtures to analyze whether the enzyme inactivation can be slowed down by ROS scavengers. Aliquots were withdrawn from the incubation mixtures every 15 minutes and assayed for activity with fresh substrate in oxygen-saturated buffer to avoid substrate limitation caused by the consumption of substrate during the incubation time. Also, interferences of the products by product inhibition are avoided this way.

When benzaldehyde was used as intermediate substrate (Fig. 1A), the hAOX1 wild-type enzyme was 80% inactivated to 20% of its initial activity during the incubation time of 60 minutes. For comparison, the control reaction without substrate also showed a loss of activity over 60 minutes, but only of 10% maximum. The presence of SOD in the incubation mixture had no positive effect on the inactivation rate of hAOX1 WT with benzaldehyde as substrate (Fig. 1A). However, when catalase was included in the incubation mixture with benzaldehyde, the enzyme showed a slower rate of inactivation of 20% maximum (Fig. 1A). The same activities were obtained when both catalase and superoxide dismutase were present (Fig. 1A). In comparison, the L438V variant was more rapidly inactivated with benzaldehyde as substrate with a complete inactivation after 30 minutes of incubation (Fig. 1B). In contrast to the hAOX1 wild-type enzyme, the inactivation of the variant was

slowed down by the inclusion of either SOD or catalase in the reaction mixture (Fig. 1B), showing that the inactivation is likely caused by both superoxide and hydrogen peroxide, since this variant produces a higher amount of superoxide than the wild-type enzyme. Similar results were obtained for both enzymes when zoniporide was used as slow substrate (Fig. 1, C and D). The wild-type enzyme was inactivated to 50% of its initial activity after an incubation time of 60 minutes. Whereas SOD had no effect on the inactivation rate, the inclusion of SOD and catalase had a minor positive effect on the inactivation rate, which was only reduced to 40% of its initial activity after 60 minutes. With zoniporide as slow substrate, the L438V variant was inactivated to 60% of its initial activity, showing that with slower substrates, the rate of inactivation is reduced (Fig. 1D). In consistency with the reaction of L348V with benzaldehyde as substrate, the rate of inactivation was only affected by the inclusion of SOD as ROS scavenger and not by catalase (Fig. 1D). When we analyzed the inactivation rate of both enzymes with phthalazine as fast substrate, a fast inactivation of the wild-type enzyme was obtained with a loss of almost 60% of its activity during the first 15 minutes (Fig. 1E). The inclusion of catalase or superoxide dismutase in the incubation mixture of the wild-type enzyme had mainly no effect on the rapid inactivation rate (Fig. 1E). Only when both catalase and superoxide dismutase were included in the incubation mixture was the inactivation rate slower and the enzyme showed a remaining 40% activity after 60 minutes of substrate turnover (Fig. 1E). Similarly, the L438V variant was rapidly inactivated during the incubation time of 60 minutes, and the rate of inactivation was only reduced a little when SOD was included in the incubation mixture. For the L438V variant, the inclusion of SOD or catalase and SOD resulted in a slower rate of inactivation, with a remaining activity of 40% after 60 minutes of incubation (Fig. 1F).

Substrate-Dependent Inactivation of hAOX1 Is Largely Reduced under Anaerobic Conditions. To get more insight into the site of inactivation of hAOX1 (Fig. 2, A and B) and the L438V variant (Fig. 2, C and D), we performed the assays using DCPIP instead of oxygen as terminal electron acceptor. DCPIP directly accepts the electrons from the Moco active site, and therefore the electron transfer via the $2 \times [2\text{Fe-2S}]$ clusters and FAD is avoided (Foti et al., 2017). When using DCPIP as electron acceptor under aerobic conditions, an inactivation of both enzymes under turnover conditions with benzaldehyde as intermediate substrate was still obtained (Fig. 2, A and C). Since the inactivation was completely diminished by the inclusion of catalase during the reaction, this shows that the oxygen present in the incubation mixture likely reacted with the enzyme as preferred electron acceptor to produce hydrogen peroxide that inactivated the enzymes (Fig. 2, A and C). Consequently, we performed the assay under anaerobic conditions to avoid a possible electron transfer to oxygen when using DCPIP as terminal electron acceptor (Fig. 2, B and D). The results show that the inactivation of both enzymes was almost completely diminished (Fig. 2, B and D). Residual amounts of

TABLE 1

Steady-state kinetic parameters hAOX1 wild type and the L438V variant with different substrates

Steady-state kinetic parameters were corrected to a molybdenum saturation of 100%. Kinetic parameters were recorded in 50 mM Tris-HCl, 200 mM NaCl, and 1 mM EDTA (pH 8.0) in the presence of 250 μM O_2 as electron acceptor using 200–400 nM enzyme. Substrate concentration were varied using 5–200 μM for benzaldehyde, 5–120 μM for phthalazine, and 75–600 μM for zoniporide. Data are mean values from three independent measurements (\pm S.D.).

Enzyme		Benzaldehyde	Phthalazine	Zoniporide
hAOX1-WT	k_{cat} (min^{-1})	14.87 \pm 1.42	123.11 \pm 1.02	0.43 \pm 0.03
	K_M (μM)	104.98 \pm 19.96	201.56 \pm 31.98	623.82 \pm 9.09
hAOX1-L438V	k_{cat} (min^{-1})	10.58 \pm 1.07	89.28 \pm 1.14	0.42 \pm 0.02
	K_M (μM)	101.48 \pm 21.01	117.06 \pm 28.92	573.41 \pm 37.51




Fig. 1. Inactivation of hAOX1 WT and the L438V variant by substrates. hAOX1 and L438V (20 μ M) were incubated in air-saturated 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, in the presence of 1.5 mM benzaldehyde (A and B), 1.5 mM zoniporide (C and D), and 1.5 mM phthalazine (E and F). The enzyme activity was tested every 15 minutes by taking a 5- μ l aliquot diluted into 500 μ l of fresh, air-saturated 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, containing either 150 μ M benzaldehyde, 150 μ M phthalazine, or 150 μ M zoniporide. In total, 2000 U/ml of catalase and SOD were included as indicated. Black dotted line, substrate; solid line, no substrate control; red line, SOD + catalase; green line, catalase; blue line, SOD.

oxygen (10 ppm) were likely still present in the reaction mixture, since catalase was still able to slightly reduce the rate of inactivation. The results conclusively show that oxygen is reacting with the enzyme very fast as the preferred electron acceptor and that the site of inactivation is the Moco active site.

Inactivated hAOX1 during Turnover Can Be Reactivated by Chemical Sulfuration. To analyze the nature of the inactivation of the Moco active site, one possible target is the terminal sulfido ligand at the molybdenum atom that is characteristic for enzymes of the XO family and essential for their catalytic activity. Since the sulfido ligand can be ligated to AOX1 by using a chemical sulfuration reaction, we analyzed whether the enzyme activity can be restored by this procedure. As a control, we chemically sulfurated the hAOX1 and L438V enzymes that were treated with cyanide, since cyanide treatment was shown to release the sulfido ligand as thiocyanide, resulting in an inactive oxo enzyme. Afterward the enzyme can be resulfurated by a chemical sulfuration procedure using dithionite and sulfide under anaerobic conditions, which restores the sulfido ligand, and enzyme activity is regained to 60% of its initial activity (Fig. 3, A and B, yellow bars). For

comparison, we chemically sulfurated the hAOX1 wild-type enzymes that were incubated with benzaldehyde (gray bars) and benzaldehyde and catalase (green bars) under turnover conditions for 60 minutes (Fig. 3A). As the cyanide-inactivated control reaction, the activity of the inactivated enzyme incubated with benzaldehyde under turnover conditions was regained to 55% of its initial activity. The chemical sulfuration had mainly no effect on the hAOX1 wild-type enzyme that contained catalase under turnover conditions, since the enzyme already showed 70% of the initial activity. For comparison, we chemically sulfurated the L438V variant. This direct comparison is intended to differentiate the inactivation site of the L438V enzyme with superoxide from the one of the wild-type enzyme with hydrogen peroxide and will reveal whether both ROS molecules target the sulfido ligand at the molybdenum atom. The results in Fig. 3B show that, after sulfuration of the cyanide and benzaldehyde inactivated L438V variant, the activity of both enzymes was restored to 60% and 40%, respectively, revealing that the sulfido ligand was also the target of inactivation of the L438V variant (Fig. 3B). No higher reactivation rate was obtained for the enzyme that was incubated in the presence of catalase under turnover conditions, which




Fig. 2. Inactivation of hAOX1 WT and the L438V variant by substrates using DCPIP as electron acceptor. hAOX1 and L438V (20 μ M) were incubated in air-saturated 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, in the presence of 1.5 mM benzaldehyde (A and C) or in oxygen-free 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, containing 150 μ M benzaldehyde and 100 μ M DCPIP (A and C) or diluted into 1 ml of oxygen-depleted 50 mM Tris-HCl, 1 mM EDTA, pH 8.0, containing 150 μ M benzaldehyde and 100 μ M DCPIP (B and D). In total, 2000 U/ml of catalase was included as indicated. black dotted line, substrate; solid line, no substrate control; green line, catalase.

shows that the sulfido ligand is the main target for enzyme inactivation of both hydrogen peroxide and superoxide.

Discussion

In this report, we show a substrate-dependent inactivation of hAOX1 and the L438V variant under turnover conditions in the presence of oxygen over time. We show that in dependence of the substrate, the enzyme inactivation of the WT enzyme can be reduced by the inclusion

of catalase in the incubation mixture. In particular, with slower substrates, catalase could prevent the damaging effect of hydrogen peroxide on the enzyme, whereas when using substrates with a high turnover rate, such as phthalazine, catalase had mainly no effect. We conclude that with fast substrates, hydrogen peroxide is produced at high rates and the inactivation of the enzyme is faster than the scavenging reaction of catalase to convert hydrogen peroxide to water and oxygen. That would provide an explanation for why, with fast substrates, no effect of




Fig. 3. Chemical sulfuration of the hAOX1 WT (A) and the L438V (B) variant after substrate-dependent inactivation. The specific activity was measured after 60 minutes of incubation of the enzyme under turnover conditions with benzaldehyde (gray bars) or with benzaldehyde and catalase (green bars). AOX1 enzymes that were inactivated with 100 μ M KCN were used as control (orange bars). After buffer exchange using PD-10 columns, the enzymes were chemically sulfurated with sulfide and dithionite, and the remaining activity was determined. KCN, potassium cyanide.

the inclusion of catalase in the incubation mixture has been observed in the past (Abbasi et al., 2019).

When the enzyme variant L438V was used, which produces high amounts of superoxide, also superoxide was damaging the enzyme, an effect that could be prevented by the inclusion of SOD in the assay. Also, in this case, SOD had a better effect to prevent enzyme inactivation with substrates with a slower turnover number in comparison with fast substrates with a high turnover number. Our results clearly show that the time-dependent inactivation under turnover conditions is based on the presence of oxygen as electron acceptor, since with other electron acceptors such as DCPIP, which takes the electrons directly from the molybdenum center (Foti et al., 2017), and additionally under anaerobic conditions, the AOX1 enzymes were not inactivated. When both DCPIP and oxygen were present as electron acceptor, the enzyme still reacted with oxygen as the preferred electron acceptor, since the electron transfer to oxygen is faster than to DCPIP (which also is a weak inhibitor of the enzyme) (Foti et al., 2017). Further, we were able to show that the site of inactivation is the terminal sulfido ligand at the molybdenum ion, since we were able to reactivate the enzyme by a chemical sulfuration reaction.

A similar inactivation has been reported for XDH and XO (Dixon, 1925; Betcher-Lange et al., 1979). Although the inactivation of XO by hydrogen peroxide has been reported already in 1925 by Dixon (1925), the nature of the reaction was mainly revealed by a study on chicken liver XDH in 1979 (Betcher-Lange et al., 1979), an enzyme that cannot be converted to the oxidase form (Rajagopalan and Handler, 1967; Nishino et al., 1989). However, XDH is nevertheless able to react with oxygen as electron acceptor, even though the turnover rate is less than 2% of that obtained with the physiologic electron acceptor NAD^+ (Nishino et al., 1989). In the study by Betcher-Lange et al. (1979), it was shown that hydrogen peroxide reacts more rapidly with the reduced enzyme than with the oxidized form of the enzyme and that the site of inactivation is the molybdenum active site and not at the FAD site. It was assumed that inactivation results from modification of the cyanolyzable sulfur (Wahl and Rajagopalan, 1982) present at the molybdenum center; the nature of the terminal sulfido ligand at the molybdenum ion had not been identified at the time of the publication in 1979. Our results on hAOX1 are fully consistent with the results of the inactivation of chicken liver XDH and show the common sensibility of these enzymes of the XO family to their reaction products hydrogen peroxide and superoxide in the reduced form. Consequently, the cyanolyzable sulfur ligand is oxidized and exchanged by an oxygen more easily in the reduced form than in the oxidized form.

In a more recent study by Abbasi et al. (2019), a nonlinear time course of hAOX1 has also been reported over incubation times of 250 minutes with different substrates. In their study, however, catalase and SOD did not influence the inactivation rate of the enzyme with different substrates. Although the authors of this study state that their buffer had an oxygen concentration of 213 μM , it can only be speculated that the actual oxygen concentration was lower at an assay temperature of 37°C. Since the authors also did not aerate their assay over the reaction time of 250 minutes, the oxygen concentration likely decreased during the long incubation time, resulting in a higher ratio of superoxide formation that more rapidly inactivates the enzyme, as reported by Lynch and Fridovich (1979) and confirmed by our studies using the L438V variant. In addition, in that study, the assays were performed at pH 7.4, which is below the pH optimum of hAOX1 of 8.0 (Coelho et al., 2015). Abbasi et al. (2019) concluded that inactivation of the enzyme during turnover is based on a slow rate of electron transfer to oxygen, which the authors propose to be the rate-limiting step of the reaction. Our results, however, do not confirm their conclusion, since the reaction of the enzyme with oxygen (taking electrons at the FAD site) was faster than the reaction of the enzyme with DCPIP (which accepts the electrons from the Moco active site). Further, the authors did not perform their assay under

anaerobic conditions with an electron acceptor other than oxygen that accepts the electrons from the FAD site (e.g., ferricyanide) to confirm their conclusions. Overall, our study confirms the time-dependent inactivation of hAOX1 under turnover conditions observed by Abbasi et al. (2019); however, we provide the evidence that the site of inactivation is the terminal sulfido ligand at the molybdenum ion, as has been previously reported for chicken liver XDH, another enzyme of the XO family (Betcher-Lange et al., 1979).

Overall, the production of hydrogen peroxide by hAOX1 has long been overlooked as a significant source for cellular ROS production. First reports, however, indicated that inhibition of hAOX1 improves the survival time of cancer patients of certain cancers (Qiao et al., 2020). In fatty liver rats treated with the AOX1 inhibitor hydralazine, liver triglyceride levels markedly decreased, indicating that AOX1 inhibitors are capable of ameliorating fatty liver and preventing liver cancer (Takeuchi et al., 2018). Raloxifene, a potent AOX1 inhibitor, is approved for the prevention of breast cancer in postmenopausal women (Provinciali et al., 2016). Raloxifene inhibits the matrix metalloproteinase-2 enzyme, which is known to be responsible for tumor invasion and the initiation of angiogenesis during the tumor growth (Ağardan et al., 2016). Inhibition of AOX1 might be another mechanism by which raloxifene reduces breast cancer (Qiao et al., 2020).

Oxidative damage has long been implicated in both the malignant phenotype and carcinogenesis (Oberley, 2002). Also, XO has been implicated to be involved in cancer development. In a recent study on XO knock-in mice, it was shown that mice that were expressing the XO-locked form showed strongly enhanced tumor growth when compared with the XDH-locked form of wild-type mice. Further, it was shown that the block of ROS production slowed down tumor growth (Kusano et al., 2019). Only mammalian XDH can be converted into the XO form, but the physiologic functions still remain elusive. Since the conversion of XDH to the XO form is not completely understood, it might be possible that ROS are also produced by hAOX1 and not only by XO in these cells. Therefore, the contribution and role of hAOX1 in ROS production that leads to cancer development needs further investigation. In summary, our study shows that hAOX1, like XDH or XO, are inhibited by ROS during substrate turnover. The inactivation of hAOX1 is thereby more rapid with substrates with a high turnover number, which cannot be prevented by catalase. When using slower substrates, the inactivation is slower based on lower levels of produced hydrogen peroxide, and consequently, the inactivation can be prevented by the inclusion of ROS scavengers. Since these enzymes produce a higher ratio of superoxide at lower concentrations of oxygen (Lynch and Fridovich, 1979), hAOX1 and XOR might be more quickly inactivated when cells get oxygen-depleted. Further, when substrates are present in amounts that are too high in the cell, hAOX1 might also get inactivated more rapidly based on a higher rate of hydrogen peroxide production. This might be a regulatory mechanism of the cell to prevent too much ROS production by hAOX1 and XOR and therefore prevent tumor growth. The role of hAOX1 in cancer and the prevention of cancer by ROS scavengers and hAOX1 inhibitors needs to be investigated in future studies.

Acknowledgments

The authors thank Angelika Lehmann, Jasmin Kurtzke, and Frank Dorendorf for their technical assistance and Mariam Esmaeeli for helpful discussions.

Authorship Contributions

Participated in research design: Leimkühler.

Conducted experiments: Garrido.

Performed data analysis: Garrido, Leimkühler.

Wrote or contributed to the writing of the manuscript: Garrido, Leimkühler.

References

- Abbasi A, Paragas EM, Joswig-Jones CA, Rodgers JT, and Jones JP (2019) Time course of aldehyde oxidase and why it is nonlinear. *Drug Metab Dispos* **47**:473–483.
- Agardan NB, Değim Z, Yölmaz Ş, Altıntaş L, and Topal T (2016) The effectiveness of raloxifene-loaded liposomes and cochleates in breast cancer therapy. *AAPS PharmSciTech* **17**:968–977.
- Beedham C (2020) Aldehyde oxidase; new approaches to old problems. *Xenobiotica* **50**:34–50.
- Bergel F and Bray RC (1959) The chemistry of xanthine oxidase. The problems of enzyme inactivation and stabilization. *Biochem J* **73**:182–192.
- Betcher-Lange SL, Coughlan MP, and Rajagopalan KV (1979) Syncatalytic modification of chicken liver xanthine dehydrogenase by hydrogen peroxide. The nature of the reaction. *J Biol Chem* **254**:8825–8829.
- Coelho C, Foti A, Hartmann T, Santos-Silva T, Leimkühler S, and Romão MJ (2015) Structural insights into xenobiotic and inhibitor binding to human aldehyde oxidase. *Nat Chem Biol* **11**:779–783.
- Coughlan MP (1980) Aldehyde oxidase, xanthine oxidase and xanthine dehydrogenase; hydroxylases containing molybdenum, iron-sulphur and flavin, in *Molybdenum and Molybdenum-Containing Enzymes* (Coughlan MP, ed) pp 119–185. Pergamon Press, Oxford.
- Dixon M (1925) Studies on xanthine oxidase: the function of catalase. *Biochem J* **19**:507–512.
- Enroth C, Eger BT, Okamoto K, Nishino T, Nishino T, and Pai EF (2000) Crystal structures of bovine milk xanthine dehydrogenase and xanthine oxidase: structure-based mechanism of conversion. *Proc Natl Acad Sci USA* **97**:10723–10728.
- Foti A, Dorendorf F, and Leimkühler S (2017) A single nucleotide polymorphism causes enhanced radical oxygen species production by human aldehyde oxidase. *PLoS One* **12**:e0182061.
- Foti A, Hartmann T, Coelho C, Santos-Silva T, Romão MJ, and Leimkühler S (2016) Optimization of the expression of human aldehyde oxidase for investigations of single-nucleotide polymorphisms. *Drug Metab Dispos* **44**:1277–1285.
- Garattini E, Fratelli M, and Terao M (2007) Mammalian aldehyde oxidases: genetics, evolution and biochemistry. *Cell Mol Life Sci* **65**:1019–1048.
- Harris CM and Massey V (1997) The reaction of reduced xanthine dehydrogenase with molecular oxygen. Reaction kinetics and measurement of superoxide radical. *J Biol Chem* **272**:8370–8379.
- Hille R, Hall J, and Basu P (2014) The mononuclear molybdenum enzymes. *Chem Rev* **114**:3963–4038.
- Holmström KM and Finkel T (2014) Cellular mechanisms and physiological consequences of redox-dependent signalling. *Nat Rev Mol Cell Biol* **15**:411–421.
- Hunt J and Massey V (1992) Purification and properties of milk xanthine dehydrogenase. *J Biol Chem* **267**:21479–21485.
- Krenitsky TA, Neil SM, Elion GB, and Hitchings GH (1972) A comparison of the specificities of xanthine oxidase and aldehyde oxidase. *Arch Biochem Biophys* **150**:585–599.
- Kundu TK, Hille R, Velayutham M, and Zweier JL (2007) Characterization of superoxide production from aldehyde oxidase: an important source of oxidants in biological tissues. *Arch Biochem Biophys* **460**:113–121.
- Kundu TK, Velayutham M, and Zweier JL (2012) Aldehyde oxidase functions as a superoxide generating NADH oxidase: an important redox regulated pathway of cellular oxygen radical formation. *Biochemistry* **51**:2930–2939.
- Kurosaki M, Bolis M, Fratelli M, Barzago MM, Pattini L, Perretta G, Terao M, and Garattini E (2013) Structure and evolution of vertebrate aldehyde oxidases: from gene duplication to gene suppression. *Cell Mol Life Sci* **70**:1807–1830.
- Kusano T, Ehrlichou D, Matsumura T, Chobaz V, Nasi S, Castelblanco M, So A, Lavanchy C, Acha-Orbea H, Nishino T, et al. (2019) Targeted knock-in mice expressing the oxidase-fixed form of xanthine oxidoreductase favor tumor growth. *Nat Commun* **10**:4904.
- Kuwabara Y, Nishino T, Okamoto K, Matsumura T, Eger BT, Pai EF, and Nishino T (2003) Unique amino acids cluster for switching from the dehydrogenase to oxidase form of xanthine oxidoreductase. *Proc Natl Acad Sci USA* **100**:8170–8175.
- Lee MC, Velayutham M, Komatsu T, Hille R, and Zweier JL (2014) Measurement and characterization of superoxide generation from xanthine dehydrogenase: a redox-regulated pathway of radical generation in ischemic tissues. *Biochemistry* **53**:6615–6623.
- Lynch RE and Fridovich I (1979) Autoinactivation of xanthine oxidase: the role of superoxide radical and hydrogen peroxide. *Biochim Biophys Acta* **571**:195–200.
- Manevski N, King L, Pitt WR, Lecomte F, and Toselli F (2019) Metabolism by aldehyde oxidase: drug design and complementary approaches to challenges in drug discovery. *J Med Chem* **62**:10955–10994.
- Neumann M and Leimkühler S (2008) Heavy metal ions inhibit molybdoenzyme activity by binding to the dithiolene moiety of molybdopterin in *Escherichia coli*. *FEBS J* **275**:5678–5689.
- Nishino T, Nishino T, Schopfer LM, and Massey V (1989) The reactivity of chicken liver xanthine dehydrogenase with molecular oxygen. *J Biol Chem* **264**:2518–2527.
- Nishino T and Tamura I (1991) The mechanism of conversion of xanthine dehydrogenase to oxidase and the role of the enzyme in reperfusion injury. *Adv Exp Med Biol* **309A**:327–333.
- Oberley TD (2002) Oxidative damage and cancer. *Am J Pathol* **160**:403–408.
- Palmer T, Santini C-L, Iobbi-Nivol C, Eaves DJ, Boxer DH, and Giordano G (1996) Involvement of the narJ and mob gene products in distinct steps in the biosynthesis of the molybdoenzyme nitrate reductase in *Escherichia coli*. *Mol Microbiol* **20**:875–884.
- Provinciani N, Suen C, Dunn BK, and DeCensi A (2016) Raloxifene hydrochloride for breast cancer risk reduction in postmenopausal women. *Expert Rev Clin Pharmacol* **9**:1263–1272.
- Qiao Y, Maiti K, Sultana Z, Fu L, and Smith R (2020) Inhibition of vertebrate aldehyde oxidase as a therapeutic treatment for cancer, obesity, aging and amyotrophic lateral sclerosis. *Eur J Med Chem* **187**:111948.
- Rajagopalan KV and Handler P (1967) Purification and properties of chicken liver xanthine dehydrogenase. *J Biol Chem* **242**:4097–4107.
- Takeuchi K, Yokouchi C, Goto H, Umehara K, Yamada H, and Ishii Y (2018) Alleviation of fatty liver in a rat model by enhancing N¹-methylnicotinamide bioavailability through aldehyde oxidase inhibition. *Biochem Biophys Res Commun* **507**:203–210.
- Terada LS, Leff JA, Guidot DM, Willingham IR, and Repine JE (1991a) Inactivation of xanthine oxidase by hydrogen peroxide involves site-directed hydroxyl radical formation. *Free Radic Biol Med* **10**:61–68.
- Terada LS, Rubinstein JD, Lesnfsky EJ, Horwitz LD, Leff JA, and Repine JE (1991b) Existence and participation of xanthine oxidase in reperfusion injury of ischemic rabbit myocardium. *Am J Physiol* **260**:H805–H810.
- Terao M, Garattini E, Romão MJ, and Leimkühler S (2020) Evolution, expression, and substrate specificities of aldehyde oxidase enzymes in eukaryotes. *J Biol Chem* **295**:5377–5389.
- Terao M, Romão MJ, Leimkühler S, Bolis M, Fratelli M, Coelho C, Santos-Silva T, and Garattini E (2016) Structure and function of mammalian aldehyde oxidases. *Arch Toxicol* **90**:753–780.
- Wahl RC and Rajagopalan KV (1982) Evidence for the inorganic nature of the cyanolyzable sulfur of molybdenum hydroxylases. *J Biol Chem* **257**:1354–1359.

Address correspondence to: Silke Leimkühler, Institute of Biochemistry and Biology, University of Potsdam, Karl-Liebknecht St. 24-25, Potsdam, 14476 Germany. E-mail: sleim@uni-potsdam.de
