

DMD#10439

**INHIBITION OF THE HUMAN LIVER MICROSOMAL AND HUMAN
CYTOCHROME P450 1A2 and 3A4 METABOLISM OF ESTRADIOL BY
DEPLOYMENT-RELATED AND OTHER CHEMICALS**

KHAWJA A. USMANI, TAEHYEON M. CHO, RANDY L. ROSE¹ AND ERNEST
HODGSON.

Arena Pharmaceuticals, Inc., 6166 Nancy Ridge Drive, San Diego CA 92121 USA, (KAU)

Department of Environmental and Molecular Toxicology, Box 7633, North Carolina State
University, Raleigh, NC 27695, USA (TMC, RLR, EH)

DMD#10439

Running title: Effects of deployment-related chemicals on estradiol metabolism

Corresponding author:

Ernest Hodgson

Department of Environmental and Molecular Toxicology

Box 7633

North Carolina State University

Raleigh, NC 27695

Phone: (919) 515-5295

Fax: (919) 513-1012

E-mail: ernest_hodgson@ncsu.edu

The number of text pages: 19

The number of tables: 1

The number of figures: 11

The number of references: 36

The number of words in the *Abstract*: 249

The number of words in the *Introduction*: 796

The number of words in the *Discussion*: 1264

DMD#10439

A list of non-standard abbreviations:

DEET - *N,N*-diethyl-*m*-toluamide

E2 - estradiol

TST - testosterone

HLM - human liver microsomes

CYP - cytochrome P450

HPLC - high performance liquid chromatography

DMD#10439

Abstract

Cytochrome P450s (CYPs) are major catalysts in metabolism of xenobiotics and endogenous substrates such as estradiol (E_2). It has previously been shown that E_2 is predominantly metabolized in humans by CYP1A2 and 3A4 with 2-hydroxyestradiol (2-OHE₂) the major metabolite. This study examines effects of deployment-related and other chemicals on E_2 metabolism by human liver microsomes (HLM) and individual CYP isoforms. Kinetic studies using HLM, CYP3A4, and CYP1A2 demonstrated similar affinities (K_m) for E_2 with respect to 2-OHE₂ production. V_{max} and CL_{int} values for HLM are 0.32 nmol/min/mg protein and 7.5 μ l/min/mg protein, those for CYP3A4 are 6.9 nmol/min/nmole CYP and 291 μ l/min/nmol CYP and those for CYP1A2 are 17.4 nmol/min/nmole CYP and 633 μ l/min/nmol CYP. Phenotyped HLM use demonstrated that individuals with high levels of CYP1A2 and CYP3A4 have the greatest potential to metabolize E_2 . Preincubation of HLM with a variety of chemicals, including those used in military deployments, resulted in varying levels of inhibition of E_2 metabolism. The greatest inhibition was observed with organophosphorus compounds, including chlorpyrifos and fonofos, with up to 80% inhibition for 2-OHE₂ production. Carbaryl, a carbamate pesticide, and naphthalene, a jet fuel component, inhibited ca. 40% of E_2 metabolism. Preincubation of CYP1A2 with chlorpyrifos, fonofos, carbaryl, or naphthalene resulted in 96, 59, 84, and 87% inhibition of E_2 metabolism, respectively. Preincubation of CYP3A4 with chlorpyrifos, fonofos, deltamethrin or permethrin resulted in 94, 87, 58 and 37% inhibition of E_2 metabolism. Chlorpyrifos inhibition of E_2 metabolism is shown to be irreversible.

DMD#10439

Introduction

The cytochrome P450 (CYP) monooxygenase system is comprised of a superfamily of heme-containing enzymes, expressed in many mammalian tissues with the highest levels found in liver, and capable of catalyzing the metabolism of a wide range of both endogenous and exogenous substrates (Nelson et al., 1996). Human CYP3A4 and CYP1A2 are two of the most important and abundant drug-metabolizing CYP isoforms in human liver microsomes. On average, CYP3A4 and CYP1A2 account for approximately 40% and 13% of the total CYP in human liver microsomes, respectively (Shimada et al, 1994; Lehman et al., 1998). Human CYP3A4 and CYP1A2 not only metabolize xenobiotics but are also responsible for the metabolism of endogenous compounds, such as steroid hormones. Both human CYP1A2 and CYP3A4 are involved in the metabolism of estradiol (E_2) and estrone (E_1) (Lee et al., 2001, 2003), whereas CYP3A4 plays a predominant role in the metabolism of testosterone (TST), androstenedione (AD), and progesterone (Waxman et al., 1988; Usmani et al., 2003).

In the human female, E_2 is the most potent primary circulating estrogen of a group of endogenous estrogen steroids which includes E_1 and estriol (E_3). During female development, E_2 stimulates the growth of female sex organs, regulates and sustains female sexual development and reproductive function, promotes hypertrophy in female breast and male muscle during puberty, initiates the synthesis of specific proteins, and controls fat deposition and distribution in subcutaneous tissues, thereby determining the characteristic female figure. Like TST in males, estrogens are the primary cause of the growth spurt, maturation of long bones and development of secondary sexual characteristics in females. In the adult, E_2 regulates events during the menstrual cycle (growth of endometrial lining), is important during pregnancy and lactation and

DMD#10439

contributes to the maintenance of sexual drive and female personality (Wilson et al., 1998; and Constanti et al., 1998). Maintaining hormonal balance relies upon a number of variables including the rate of hormone synthesis, interactions among hormones, and rates of secretion, transport, and metabolism by phase I and phase II enzymes. CYPs (phase I) are a major element in the maintenance of a proper steroid hormone levels in mammalian systems and are the sole subject of this investigation. Exposure to foreign compounds may exert changes in endocrine function both directly (hormone agonists or antagonists) or indirectly (altering circulating levels of hormones by influencing rates of hormone synthesis or metabolism) that can severely affect steroid hormone action (Wilson and LeBlanc, 1998). It follows that perturbation of the CYP system by xenobiotics may in turn affect the subsequent metabolism and disposition of E₂. Perturbations in E₂ metabolism may affect levels of circulating E₂ with possible reproductive and other consequences, including further modulation of the expression of some CYP proteins.

The uses of deployment-related chemicals are essential for the health and well-being of deployed forces during both peace keeping and wartime missions. Human hazards associated with chemicals used to protect personnel during peace and wartime are poorly understood since the definition of cause and effect relationships depends on knowledge of the mechanisms of toxic action and interaction in humans. Following the Gulf War some veterans reported illnesses which may have been the result of chemical exposures. Some studies of these veterans have concluded that significant correlations between perceived illnesses and chemical use exist (Haley and Kurt, 1997). The reported chemical exposures included the insect repellent *N,N*-diethyl-*m*-toluamide (DEET), insecticides such as permethrin and chlorpyrifos used to protect against insect borne diseases and the neuroprotective agent pyridostigmine bromide used to protect

DMD#10439

against possible nerve gas attack. It has been reported that chlorpyrifos and DEET are metabolized by human CYPs (Tang et al., 2001; Usmani et al., 2002) and that interactions of deployment-related chemicals can inhibit or induce the CYPs involved in their metabolism (Usmani et al., 2002). In a recent study examining the effects of various deployment-related chemicals on TST metabolism, Usmani et al. (2003) reported that organophosphorus pesticides are potent non-competitive and irreversible inhibitors of TST metabolism due to CYP3A4 inhibition. Other studies have reported that interaction of Gulf War related chemicals could produce greater than additive toxicity in rats and mice (McCain et al., 1997; Chaney et al., 1997), increased neurotoxicity in hens associated with increased inhibition of brain acetylcholinesterase and Neurotoxicity Target Esterase (Abou-Donia et al., 1996a,b) and neurobehavioral deficits associated with significant inhibition of brainstem acetylcholinesterase activities in rats (Abou-Donia et al., 2001). Possible detrimental interactions as a result of altered estrogen metabolism have not yet been examined relative to these chemicals nor have studies been carried out to examine the induction or inhibition by these or related compounds of human CYP-mediated metabolism of estrogens, such as E₂.

The objective of present study was to study the inhibition or activation by various deployment-related chemicals of the metabolism of E₂ by HLM, CYP1A2, and CYP3A4.

DMD#10439

Materials and Methods

Chemicals. Chlorpyrifos, fonofos, phorate, DEET, fipronil, imidacloprid, deltamethrin, permethrin, carbofuran, carbaryl, and naphthalene were purchased from ChemService (West Chester, PA). Pyridostigmine bromide was purchased from Roche (Indianapolis, IN). Testosterone (TST), 17 β -estradiol (E₂), 2-hydroxyestradiol (2-OHE₂), and 6 β -hydroxytestosterone were purchased from Steraloids (Newport, RI). HPLC grade water, methanol, acetonitrile, and tetrahydrofuran were purchased from Fisher Scientific (Pittsburgh, PA). All other chemicals, if not specified, were purchased from Sigma (St. Louis, MO). The structures of all chemicals tested as inhibitors are shown in Fig 1.

Human liver microsomes and human CYP isoforms. Pooled human liver microsomes (HLM) (pooled from 21 donors), single-donor HLM, and the human CYP isoforms, 1A2 and 3A4, expressed in baculovirus infected insect cells (Sf9) (BTI-TN-5B1-4) were purchased from BD Biosciences (Woburn, MA).

Assay conditions. All assays were conducted in triplicate and all the reactions were pre-warmed for 5 min at 37°C. All the reactions contained NADPH-regenerating system (0.25 mM NADP, 2.5 mM glucose 6-phosphate, and 2 U/ml glucose-6-phosphate dehydrogenase) and 100 mM potassium phosphate buffer (pH 7.4) containing 5 mM ascorbic acid and 3.3 mM MgCl₂. Final concentrations of HLM were 1 mg/ml while final concentrations of CYP3A4 and CYP1A2 were 50 pmol/ml. The final reaction volume was 250 μ l and all reactions were terminated after 20 min by the addition 150 μ l of methanol and vortexing. After 5 min centrifugation at 21,000 g in a microcentrifuge, the supernatants were analyzed for E₂ metabolite concentrations by HPLC. The protein concentrations and incubation times used in the assays were found to be in the linear

DMD#10439

range in preliminary experiments. No metabolites were detected when incubations were carried out in the absence of an NADPH-generating system.

***In vitro* E₂ metabolism.** Enzyme kinetic assays using HLM or the recombinant isoforms CYP1A2 or 3A4 were performed by incubation of serial concentrations of E₂ (final concentrations 3.125-200 μ M) with an NADPH-generating system in potassium phosphate buffer and incubated at 37°C for 5 min. The reactions were initiated by the addition of ice cold HLM (final protein concentration 1 mg/ml), CYP1A2 or 3A4. No measurable lag phase was observed.

To determine the metabolic activities of individual HLM, incubations of E₂ (final concentration 50 μ M) were conducted for 20 min at 37°C using 1 mg protein/ml. To differentiate between the relative concentrations of CYP1A2 and CYP3A4 to the metabolism of E₂ in individual HLM, a specific CYP3A4 inhibitor, ketoconazole (final concentration 2 μ M), was added to the reaction mixture simultaneously with 50 μ M E₂.

Preliminary inhibition studies. The effects of test chemicals (chlorpyrifos, fonofos, phorate, DEET, fipronil, imidacloprid, deltamethrin, permethrin, carbofuran, carbaryl, naphthalene, and pyridostigmine bromide) on E₂ metabolism were examined using HLM, CYP1A2, or 3A4 after preincubation with the test compounds in otherwise complete mixtures lacking only the substrate, E₂. The HLM, CYP3A4, or CYP1A2 were incubated with individual test compounds (final concentration 50 μ M) for 5 min at 37°C before adding E₂ (final concentration 50 μ M). All inhibitors, except pyridostigmine bromide, were added in 2.5 μ l of acetonitrile with an equal volume of acetonitrile being added to the control incubations. Pyridostigmine bromide was added in 2.5 μ l of water. Because of the likelihood that

DMD#10439

phosphorothioates are metabolized to irreversible CYP inhibitors, preincubation is tested as a variable with these chemicals. Preliminary experiments indicated little or no difference between pre- and co-incubation for other potential inhibitors. Only those chemicals which were capable of greater than 50% inhibition of E₂ metabolism at concentration of 50 μM were studied further.

Inhibition kinetics of deployment-related chemicals on E₂ metabolism by CYP3A4 and CYP1A2. The chemicals for which IC₅₀ values were below 25 μM as determined by preliminary incubation studies were analyzed further for inhibitory potential. Range finding assays were conducted for chlorpyrifos, fonofos, phorate, deltamethrin, and permethrin (0.39–200 μM) by incubating with CYP3A4 for 5 min at 37°C before adding E₂ (final concentration 50 μM). The kinetic study of E₂ using varying E₂ concentrations (6.25 – 100 μM) was performed using CYP3A4, which was preincubated with chlorpyrifos or fonofos (0.78, 1.56 or 3.125 μM) for 5 mins at 37°C.

For CYP1A2, range finding assays were conducted for chlorpyrifos and fonofos as described above. Concentrations of chlorpyrifos (0.5 and 2.0 μM) and fonofos (12.5 and 50.0 μM) were preincubated with CYP1A2 for 5 min at 37°C before adding E₂ (final concentration 3.125-100 μM) in order to measure their inhibition of CYP1A2 metabolism of E₂. In all inhibition studies involving preincubation with the inhibitor, the NADPH generating system was present during the preincubation.

Pre-incubation time and concentration-dependent inactivation by chlorpyrifos of CYP3A4 metabolism of estradiol. These studies were conducted using the same conditions as described in the assay conditions (50 μM E₂; 20 min. incubation at 37°C) but varying concentrations (0, 1, 2, 5, 10 and 50 μM of CPS) and pre-incubation times (0, 1, 2, 3, 4 and 5 min.). The generation of 2-

DMD#10439

OHE₂ (the major metabolite of E₂) was compared in this assay, conducted in duplicate. Inactivation parameters (k_{inact} and K_I) were estimated as described in previous studies (Silverman, 1995, Heydari et al, 2004). The rate constant for initial inactivation (k_{obs}) at each concentration of inhibitor (CPS) was calculated from the slopes in the regression plots ($-k_{\text{obs}}$) (natural logarithm of remaining activity as % of control activity versus pre-incubation time). The inactivation values (k_{inact} and K_I) were calculated as described in Heydari et al. (2004); $k_{\text{obs}} = k_{\text{inact}} \times [I] / (K_I + [I])$ where $[I]$ is the inhibitor concentration, k_{inact} is the maximum rate constant for inactivation and K_I is the inhibitor concentration that produces half maximal rate of inactivation.

Inhibition kinetics of carbaryl and its metabolites on E₂ metabolism by CYP1A2. Before the K_i (inhibition constant) determination was done, range finding assays were conducted for the inhibition by carbaryl and its metabolites of the metabolism of E₂ by CYP1A2. Varying concentrations (1.56–50 μM) of carbaryl, 4-hydroxycarbaryl, 5-hydroxycarbaryl, and carbaryl methylol were pre- or co-incubated with CYP1A2. Preincubation experiments with inhibitors were conducted similarly to those described above. For coincubation experiments, carbaryl or its metabolites were incubated with E₂ (final concentration 50 μM). The reactions were initiated by the addition of ice-cold CYP1A2. For Michaelis-Menten plots, carbaryl (6.25 and 12.5 μM), 4-hydroxycarbaryl (6.25 and 12.5 μM), and carbaryl methylol (12.5 and 25 μM) were coincubated with E₂ (final concentration 3.125–100 μM). The reactions were initiated by the addition of ice-cold CYP1A2.

Inhibition kinetics of naphthalene and its metabolites on E₂ metabolism by CYP1A2.

Before the K_i determination was done, range finding assays were conducted for the inhibition by

DMD#10439

naphthalene and its metabolites of the metabolism of E₂ by CYP1A2. Varying concentrations of naphthalene (3.125 – 25 μM), 1-naphthol (0.78 - 25 μM), 2-naphthol (25 μM), *trans*-1,2-dihydro-1,2-naphthalenediol (25 μM), 1,2-naphthoquinone (25 μM), 1,4-naphthoquinone (25 μM), and 1,4-dihydroxynaphthene (25 μM) were pre- or co-incubated with CYP1A2. The pre- and co-incubation experiments with naphthalene and its metabolites were conducted similarly as described above. For Michaelis-Menten plots, naphthalene (1.56, 3.125, and 6.25 μM) was preincubated with CYP1A2 for 5 min at 37°C before adding E₂ (final concentration 3.125-100 μM).

For Michaelis-Menten plots, 1-naphthol (1.56 and 3.125 μM) was coincubated with E₂ (final concentration 3.125-100 μM). The reactions were initiated by the addition of ice-cold CYP1A2.

To demonstrate whether 1-naphthol inhibition is reversible or irreversible, incubations with and without 1-naphthol (3.125 μM) were conducted with E₂ (final concentration 50 μM). The reactions were initiated by the addition of varying concentrations of ice-cold CYP1A2 (3.125-12.5 pmol).

Analysis of 2-OHE₂ and 6β-hydroxytestosterone by HPLC. Analysis of the E₂ metabolite 2-OHE₂ was performed with an HPLC system coupled with in-line UV detection as described previously (Suchar et al., 1995). The HPLC system consisted of a Waters 2690 separation module and a Waters UV photodiode array detector (model 2996). All system components were controlled through the Waters powerline firmware. Data was collected via a Waters system controller and analyzed using Waters Empower software. The solvent system for separation of E₂ and 2-OHE₂ consisted of acetonitrile (solvent A), 0.1% acetic acid in water

DMD#10439

(solvent B), and 0.1% acetic acid in methanol (solvent C). The solvent gradient (solvent A/solvent B/solvent C) used for eluting E_2 and 2-OHE₂ was as follows: 8 min of isocratic at 16:68:16, 7 min of a concave gradient (curve number 9) to 18:64:18, 13 min of a concave gradient (curve number 8) to 20:59:21, 10 min of a convex gradient (curve number 2) to 22:57:21, 13 min of a concave gradient (curve number 8) to 58:21:21, followed by a 0.1 min step to 92:5:3 and a 3.9 min isocratic period at 92:5:3. The gradient was returned to the initial condition (16:68:16) for 2 min and held for 3 min before analyzing the next sample. The flow rate was 1.2 ml/min. E_2 and 2-OHE₂ were separated by an Ultracarb 5 ODS column (150 x 4.6 mm, Phenomenex, Rancho Palos Verdes, CA) at 30°C and detected at 280 nm. The limits of detection for 2-OHE₂ was approximately 0.04 μM. Concentrations of 2-OHE₂ were obtained from the chromatographic peak area from a standard curve (0.15 μM – 20.0 μM).

TST and its metabolite, 6β-hydroxytestosterone, were separated using the method described by Usmani et al. (2003). E_2 and its metabolites eluted after 6β-hydroxytestosterone and TST.

Data Analysis and statistics. The apparent K_m and V_{max} parameters were calculated using the SigmaPlot Enzyme Kinetics Module for Windows, version 1.1 (SPSS Inc., Chicago, IL). The K_i values were estimated by nonlinear regression analysis by fitting different models of enzyme inhibition to the kinetic data using SigmaPlot Enzyme Kinetics Module for Windows, version 1.1 (SPSS Inc., Chicago, IL). The mode of inhibition was established by comparing the statistical results including the r^2 information criterion values of different inhibition models and by selecting the one with the best fit. Furthermore, the mode of inhibition was established by comparing the 95% confidence intervals that did not overlap for K_m and V_{max} values. Significant

DMD#10439

differences between data sets were determined by student t-test method using SigmaPlot for Windows, version 8.0 (SPSS Inc., Chicago, IL).

DMD#10439

Results

Enzyme kinetics. HLM, CYP3A4, and CYP1A2 all displayed similar K_m values for metabolism of E_2 to 2-OHE₂ (Table 1). V_{max} and CL_{int} values for HLM were 0.32 nmol/min/mg protein and 7.5 μ l/min/ mg protein, those for CYP3A4 were 6.9 nmol/nmole CYP and 291 μ l/min/ nmol CYP and those for CYP1A2 were 17.4 nmol/min/nmole CYP and 633 μ l/min/ nmol CYP.

E_2 metabolism in single-donor HLM. Incubations of E_2 with individual donor HLM (6 male and 11 female) demonstrated as much as 36-fold variability in metabolism among individuals (Fig. 2). Phenotype data based on metabolic activities of these individuals was used to derive correlations between CYP isoform content and E_2 metabolic activity. CYP3A4 was the isoform with the best correlation ($r^2 = 0.87$). All other correlations between E_2 hydroxylation activity and specific CYP isoforms were less than 0.40. To determine the CYP1A2 contribution to E_2 metabolism among individual HLM, ketoconazole (2 μ M), a specific CYP3A4 inhibitor, was used to inhibit CYP3A4 activity. Preliminary observations with ketoconazole indicated that 2 μ M inhibited less than 5% of CYP1A2 activity (data not shown). In the presence of ketoconazole, variation among individuals was reduced to 12-fold and the correlation of CYP1A2 with E_2 metabolism increased to 0.81 (Fig. 2).

Inhibition of E_2 metabolism. The effects of various deployment-related and other chemicals on E_2 (50 μ M) metabolism were investigated by preincubating 50 μ M concentrations of each chemical with pooled HLM (Fig. 3). The organophosphorus compounds, chlorpyrifos fonofos, and phorate all significantly inhibited E_2 metabolism to 2-OHE₂, although inhibition by chlorpyrifos and fonofos was profound (ca. 80%) compared to phorate (ca. 16.5%).

DMD#10439

Preincubation of pooled HLM with carbaryl, a carbamate, and naphthalene, a jet fuel component, also resulted in significant inhibition (ca. 40%) of E₂ metabolism to 2-OHE₂. Fipronil, a phenyl pyrazole insecticide also significantly inhibited E₂ metabolism, though to a lesser degree (ca. 18%).

To further explore the inhibition of E₂ metabolism observed in HLM, assays were also conducted using CYP3A4 and CYP1A2. Preincubation of CYP3A4 with a variety of chemicals resulted in varying levels of activation and inhibition of E₂ metabolism (Fig. 4). E₂ metabolism was significantly inhibited by the organophosphorus compounds, chlorpyrifos and fonofos, with up to ca. 90% inhibition of E₂ metabolism to 2-OHE₂. Preincubation of CYP3A4 with pyrethroids, deltamethrin and permethrin resulted in ca. 58 and 37% inhibition of E₂ metabolism to 2-OHE₂, respectively. In contrast, preincubation of CYP3A4 with DEET, an insect repellent, and carbofuran, a carbamate, resulted in the production of small, but not significantly greater, levels of 2-OHE₂.

Similarly, preincubation of CYP1A2 resulted in varying levels of activation and inhibition of E₂ metabolism (Fig. 5). Preincubation of chlorpyrifos and fonofos resulted in ca. 96% and 59% inhibition of E₂ metabolism, respectively. In contrast with CYP3A4, preincubation of CYP1A2 with carbaryl and naphthalene resulted in significant (ca. 85%) inhibition of E₂ metabolism. Some chemicals, such as fipronil and deltamethrin appeared to produce small, but not significantly higher, levels of 2-OHE₂.

Mechanism of inhibition. To investigate the type of inhibition of CYP3A4 by chlorpyrifos and fonofos in the production of 2-OHE₂, a major E₂ metabolite, different concentrations of chlorpyrifos and fonofos were preincubated for 5 min with CYP3A4 before

DMD#10439

adding varying concentrations of E₂. Michaelis-Menten plots showed that the V_{max} values were significantly reduced without affecting K_m values, indicative of a non-competitive inhibition of E₂ metabolism by chlorpyrifos and fonofos (Fig. 6).

In a similar manner, inhibition of CYP1A2 metabolism of E₂ by chlorpyrifos and fonofos involved a 5 minute preincubation prior to the addition of varying concentrations of E₂. As shown for CYP3A4 (Fig. 5), Michaelis-Menton plots for CYP1A2 also indicated non-competitive inhibition (Fig. 7).

The production of 2-OHE₂ by CYP3A4 was significantly inhibited by CPS in a time and concentration-dependent manner (Fig. 8A). Inactivation parameters (k_{inact} and K_I) determined for CYP3A4 metabolism of estradiol by CPS were 0.3 min⁻¹ and 4.9 μM, respectively (Fig. 8B). Even at a pre-incubation of 0', the metabolic activity of CYP3A4 for estradiol was, apparently, significantly decreased by CPS, in a dose-dependent manner, indicating a very rapid initial rate of inhibition..

For IC₅₀ values, varying concentrations of carbaryl and its metabolites were pre- and co-incubated with CYP1A2. Under preincubation conditions but not with coincubation, carbaryl, 4-hydroxycarbaryl, and carbaryl methylol caused small increases in the production of 2-OHE₂ at the low concentrations (1.56, 3.125, and 6.25 μM) whereas at high concentrations (12.5 - 50 μM) inhibition of E₂ metabolism was observed (data not shown). Under preincubation conditions but not with coincubation, 5-hydroxycarbaryl caused small increases in the production of 2-OHE₂ at all concentrations (1.56 - 50 μM) (data not shown). Under coincubation conditions, carbaryl, 4-hydroxycarbaryl, 5-hydroxycarbaryl, and carbaryl methylol all caused inhibition of E₂ metabolism at all concentrations tested (1.56 - 50 μM) (data not shown). To determine the K_i

DMD#10439

value and investigate the type of inhibition of CYP1A2 metabolism of E₂ by carbaryl and its metabolites, different concentrations of carbaryl, 4-hydroxycarbaryl, and carbaryl methylol were coincubated with CYP1A2 and varying concentrations of E₂. Michaelis-Menten plots showed that the V_{\max} values were significantly reduced without affecting K_m values, indicative of a non-competitive inhibition of E₂ metabolism by carbaryl and its metabolites (Fig. 9). The inhibition constants (K_i) indicated that carbaryl is the most potent inhibitor of E₂ metabolism with a K_i value of 7.7 μM , 4-hydroxycarbaryl is the second best inhibitor with a K_i value of 10.2 μM , while carbaryl methylol is a weak inhibitor with a K_i value of 29.0 μM .

In the determination of IC₅₀ values, varying concentrations of naphthalene and its metabolites were pre- and co-incubated with CYP1A2. Preincubation of naphthalene increased the inhibition of E₂ metabolism by nearly 2-fold in comparison with the coincubation of naphthalene with E₂, indicating that a reactive metabolite is involved (data not shown). No differences were observed in the IC₅₀ values when 1-naphthol was pre- or co-incubated with E₂ (data not shown). 2-Naphthol, 1,2-naphthoquinone, 1,4-naphthoquinone, and 1,4-dihydronaphthalene (25 μM) inhibited E₂ (50 μM) metabolism ca. 85 – 100%, whereas trans-1,2-dihydro-1,2-naphthalenediol had no inhibitory effect on E₂ metabolism (data not shown). To determine the K_i value and to investigate the type of inhibition of CYP1A2 by naphthalene on E₂ metabolism, naphthalene was preincubated for 5 min with CYP1A2 before adding varying concentrations of E₂. The Michaelis-Menten plot showed that the V_{\max} values were significantly reduced without affecting K_m values, indicative of a non-competitive inhibition of E₂ metabolism by naphthalene (Fig. 10). To investigate the type of inhibition of CYP1A2 by 1-naphthol on E₂ metabolism, 1-naphthol was coincubated with CYP1A2 and varying concentrations of E₂. 1-

DMD#10439

Naphthol, a predominant naphthalene metabolite, was shown to be a noncompetitive inhibitor of E₂ metabolism (Fig. 11). Further investigation of noncompetitive reversible or nonreversible inhibition data revealed that the inhibition of 2-OHE₂ by 1-naphthol is nonreversible (data not shown).

DMD#10439

Discussion

CYP-dependent hydroxylation is a major pathway of oxidative metabolism of E_2 in mammalian liver. The studies carried out using human CYP isoforms provided further insight into the range of E_2 hydroxylation reactions that can be catalyzed by human CYP enzymes. Human CYP1A2 and CYP3A4 are the major isoforms responsible for E_2 metabolism and 2-OH E_2 is the major E_2 metabolite, as previously demonstrated (Yamazaki et al., 1998; Lee et al., 2001, 2003). The phenotyped HLM data and the mean metabolic intrinsic clearance rates, as estimated by V_{\max}/K_m , corroborates earlier findings (Yamazaki et al., 1998; Lee et al., 2001, 2003) that both CYP3A4 and CYP1A2 play major roles in the metabolism of E_2 in human liver and the content of these two CYPs in HLM determine which isoform is more important. However, CYP3A4 may play a more important role than CYP1A2 because on average CYP3A4 accounts for approximately 40% of the total CYP in HLM. Our kinetic data presented in this study for CYP1A2 and CYP3A4 are in general agreement with the previous reports showing that CYP1A2 is more active than CYP3A4 in E_2 metabolism (Yamazaki et al., 1998; Badawi et al., 2001).

CYP3A4 is one of the most important and abundant CYP isoforms in human liver and has broad substrate specificity. CYP3A4 not only metabolizes xenobiotics but is also responsible for the metabolism of endogenous compounds such as steroid hormones, including TST and E_2 (Usmani et al., 2003; Lee et al., 2003). Because TST and E_2 are predominantly metabolized by CYP3A4, both can potentially compete for the same catalytic site of CYP3A4. The usual interaction between two different substrates for the same enzyme is competitive inhibition. But because the K_m values for TST and E_2 are quite different ($K_m = 108 \mu\text{M}$ and 24

DMD#10439

μM for TST and E_2 , respectively), interactions between these substrates may be complicated due to the allosteric characteristics of CYP3A4 (Shimada and Guengerich 1989; Lee et al., 1995; Usmani et al., 2003; Williams et al., 2004). The recent demonstration of a relatively large substrate binding catalytic site by crystal structure elucidation is consistent with the capacity of CYP3A4 to accommodate large molecules and possibly more than one substrate (Yano et al., 2004).

Endogenous steroids, such as E_2 , always exist in vivo and considerable amounts of these steroids are metabolized by the CYPs expressed in the human liver, where foreign compounds are mainly metabolized. Since TST and E_2 both have important, and different effects in vivo, the inhibition of the metabolism of both of these steroid hormones must be considered in detail. If xenobiotics substantially alter enzymes such as CYP3A4 and CYP1A2, they may affect the rate of E_2 metabolism, ultimately disrupting E_2 homeostasis. Preincubation of pooled HLM, CYP3A4, and CYP1A2 with organophosphorus compounds, such as chlorpyrifos and fonofos, resulted in the extensive inhibition of the production of 2-OHE₂, a major E_2 metabolite. Chlorpyrifos and fonofos inhibited 2-OHE₂ formation noncompetitively and are among the most potent inhibitors of the CYP-dependent oxidation of E_2 yet described. Organophosphorus pesticides, such as chlorpyrifos and fonofos are activated by a CYP-catalyzed desulfuration reaction (Fukuto, 1990). The sulfur atom released from these pesticides in this reaction is highly reactive and is believed to bind immediately to the heme iron of P450 and inhibit its activity (Norman et al, 1974, Halpert et al., 1980, Neal, 1980, Neal and Halpert, 1982. Butler and Murray, 1997). Our previous study demonstrated that organophosphorus pesticides are potent

DMD#10439

non-competitive and irreversible inhibitors of TST metabolism by HLM and by CYP3A4 (Usmani et al., 2003).

Enzyme activity is often modulated by inhibition or induction, either condition modifying the extent to which xenobiotics or endogenous substrates are metabolized (Guengerich, 1997; Szklarz and Halpert, 1998). Inhibition may, in some interactions, be more serious than enzyme induction since inhibition occurs more rapidly than induction (Guengerich, 1997). Preincubation of CYP3A4 with chlorpyrifos and fonofos resulted in almost complete inhibition of E₂ metabolism to 2-OHE₂. Preincubation of CYP1A2 with chlorpyrifos resulted in almost complete inhibition of 2-OHE₂ formation, whereas fonofos was not as potent an inhibitor. The kinetics of inactivation of E₂ metabolism by CPS confirms that CPS is a mechanism based inactivator of CYP3A4 in these studies. Pre-incubation of CPS in the presence of an NADPH-generating system increased the inhibitory effect in a time and concentration-dependent manner. The significant reduction of 3A4 activity toward E₂ metabolism by CPS at time 0' suggests that the binding affinity of the inhibitor (CPS) is higher than that of the substrate (E₂) with the CYP3A4 isoform and that the initial rate of inhibition is rapid. The relatively low K_I value (high inhibitory potency) of CPS toward CYP3A4 metabolism of E₂ appears to confirm the strong binding affinity of CPS with this isoform. The possibility exists, therefore, that inhibition of CYP3A4 and CYP1A2 by these chemicals could lead to higher levels of E₂ and alter hormonal properties. The *in vivo* importance of these observations will require further studies.

Carbaryl is a widely used anticholinesterase carbamate insecticide. In an *in vitro* study, 5-hydroxycarbaryl, 4-hydroxycarbaryl, and carbaryl methylol were identified as the major metabolites of carbaryl produced by HLM (Tang et al., 2002). CYP isoform data indicated that

DMD#10439

CYP1A1, 1A2, 2B6, 2C19, and 3A4 are the most active isoforms in human metabolism of carbaryl (Tang et al., 2002). In our preliminary inhibition studies we found that preincubation of pooled HLM and CYP1A2 with carbaryl significantly inhibited E_2 metabolism. Because carbaryl is metabolized by CYP1A2 while also demonstrating significant inhibition of E_2 metabolism we expected competitive interactions between E_2 and carbaryl or its metabolites. However, carbaryl and its metabolites inhibited formation of 2-OHE₂ noncompetitively in our coincubation studies. The K_i values indicated that carbaryl and 4-hydroxycarbaryl were much better inhibitors of 2-OHE₂ than carbaryl methylol. While there is no obvious explanation of the slight activation of E_2 at low concentrations of carbaryl, 4-hydroxycarbaryl and carbaryl methylol, it may be noted that this is a common phenomenon in the oxidation of TST by human liver microsomes (Usmani et al., 2003).

The polycyclic aromatic hydrocarbon naphthalene is an industrial chemical, an environmental pollutant, and a component of jet fuel and has considerable toxicological importance because of widespread human exposure and its potential to form toxic and carcinogenic metabolites in humans (Wilson et al., 1996). We recently demonstrated that CYP1A2 is the most efficient of the 15 different isoforms tested for their ability to metabolize naphthalene to its two major metabolites in HLM, 1-naphthol and 1,2-dihydro-1,2-naphthalenediol (Cho et al., 2006). In this study, preincubations of naphthalene with pooled HLM and CYP1A2 significantly inhibited E_2 metabolism. Because naphthalene is metabolized by CYP1A2 and also significantly inhibited E_2 metabolism we expected a competitive interaction between E_2 and naphthalene or its metabolite. However, this study demonstrated noncompetitive inhibition of E_2 metabolism. Coincubation of 1-naphthol, a well known

DMD#10439

cytotoxic metabolite of naphthalene, resulted in noncompetitive and irreversible inhibition of E₂ metabolism.

In conclusion, the hydroxylation of E₂ by CYP3A4 and CYP1A2 isoforms indicates important functions for these enzymes other than detoxification of xenobiotics. The deployment-related and other chemicals used in this study, including pesticides, caused a marked modification of CYP mediated E₂ metabolism *in vitro*. Organophosphorus pesticides were very potent inhibitors of the production of 2-OHE₂ and inhibited E₂ metabolism noncompetitively. Carbaryl and some of its metabolites inhibited E₂ metabolism non-competitively. Naphthalene inhibited E₂ metabolism noncompetitively, an effect probably due to 1-naphthol, a naphthalene metabolite. 1-Naphthol inhibits E₂ metabolism noncompetitively and irreversibly. It should be noted that *in vivo* toxicokinetic data are not available for these chemicals in humans and thus the effects noted indicate only the potential for *in vivo* effects.

DMD#10439

References

- Abou-Donia MB, Wilmarth KR, Abdel-Rehman AA, Jensen FK, Oehme FW and Kurt TL (1996a) Increased neurotoxicity following concurrent exposure to pyridostigmine bromide, DEET and chlorpyrifos. *Fund Appl Toxicol* **34**:201-222.
- Abou-Donia MB, Wilmarth KR, Jensen FK, Oehme FW and Kurt TL (1996b) Neurotoxicity resulting from coexposure to pyridostigmine bromide, DEET and permethrin: implications of Gulf War chemical exposure. *Toxicol Environ Health* **48**:35-56.
- Abou-Donia MB, Goldstein LB, Jones KH, Abdel-Rehman AA, Damodaran TV, Dechkovskaia AM, Bullman SL, Amir BE, and Khan WA (2001) Locomotor and sensorimotor performance deficit in rats following exposure to pyridostigmine bromide, DEET, and permethrin, alone and in combination. *Toxicol Sci* **60**:305-314.
- Badawi AF, Cavalieri EL and Rogan EG (2001) Role of human cytochrome P450 1A1, 1A2, 1B1, and 3A4 in the 2-, 4-, and 16 α -hydroxylation of 17 β -estradiol. *Metabolism* **50**:1001-1003.
- Butler AM and Murray M (1997) Biotransformation of parathion in human liver: participation of CYP3A4 and its inactivation during microsomal parathion oxidation. *J Pharmacol Exp Ther* **280**: 966-973.
- Constanti A, Bartke A, and Khardori R (1998) Basic Endocrinology. Howard Academic Publishers, Amsterdam.
- Chaney LA, Rockhold RW, Mozingo JR, and Hume AS (1997) Potentiation of pyridostigmine bromide toxicity in mice by selected adrenergic agents and caffeine. *Vet Human Toxicol* **39**:214-219.

DMD#10439

- Cho TM, Rose, RL, and Hodgson, E. (2006) In vitro metabolism of naphthalene by human liver microsomal cytochrome P450 enzymes. *Drug Metabol Disp* 34: 176-183.
- Fukuto TR (1990) Mechanism of action of organophosphorus and carbamate insecticides. *Environ Health Perspect* **87**:245-254.
- Guengerich FP (1997) Role of cytochrome P450 enzymes in drug-drug interactions. *Adv Pharmacol* **43**:7-35.
- Haley RW and Kurt TL (1997) Self reported exposure to neurotoxic chemical combinations in the Gulf War: a cross-sectional epidemiological study. *JAMA* **277**:231-237.
- Halpert J, Hammond D and Neal RA (1980) Inactivation of purified rat liver cytochrome P-450 during the metabolism of parathion (diethyl p-nitrophenyl phosphorothionate). *J. Biol Chem* **255**: 1080-1089.
- Heydari A, Yeo KR, Lennard MS, Ellis SW, Tucker, GT and Rostami-Hodjegan A (2004) Mechanism-based inactivation of CYP2D6 by methylenedioxymethamphetamine. Short communication. *Drug Metab Disp* **32**: 1213-1217.
- Lee AJ, Kosh JW, Conney AH and Zhu BT (2001) Characterization of the NADPH-dependent metabolism of 17 β -estradiol to multiple metabolites by human liver microsomes and selectively expressed human cytochrome P450 3A4 and 3A5. *J Pharmacol Exp Ther* **298**:420-432.
- Lee AJ, Cai XM, Thomas PE, Conney AH and Zhu BT (2003) Characterization of the oxidative metabolites of 17 β -estradiol and estrone formed by 15 selectively expressed human cytochrome P450 isoforms. *Endocrinology* **144**:3382-3398.

DMD#10439

- Lee CA, Kadwell SH, Kost TA and Serabjit-Singh CJ (1995) CYP 3A4 expressed by insect cells infected with a recombinant baculovirus containing both CYP 3A4 and human NADPH-cytochrome P450 reductase is catalytically similar to human liver microsomal CYP 3A4. *Arch Biochem Biophys* **319**:157-167.
- Lehmann JM, McKee DD, Watson MA, Willson TM, Moore JT, and Kliewer SA (1998) The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J Clin Invest* **102**:1016-1023.
- McCain WF, Lee R, Johnson MS, Whaley JE, Ferguson JW, Beall P, and Leach G (1997) Acute oral toxicity study of pyridostigmine bromide, permethrin and DEET in the laboratory rat. *J Toxicol Environ Health* **50**:113-124.
- Neal RA (1980) Microsomal metabolism of thiono-sulfur compounds: mechanisms and toxicological significance. *Rev Biochem Toxicol* **2**:131-171.
- Neal RA and Halpert J (1982) Toxicology of thiono-sulfur compounds. *AnnRev Pharmacol Toxicol* **22**: 321-339.
- Nelson DR, Koymans L, Kamataki T, Stegeman JJ, Feyereisen R, Waxman DJ, Waterman MR, Gotoh O, Coon MJ, Estabrook RW, Gunsalus IC and Nebert DW (1996) P450 superfamily: Update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics* **6**:1-42.
- Norman BJ, Poore RE and Neal RA (1974) Studies of the binding of sulfur released in the mixed-function oxidase catalyzed metabolism of diethyl p-nitrophenyl phosphorothionate (parathion) to diethyl p-nitrophenyl phosphate (paraoxon). *Biochem Pharmacol* **23**: 1733-1744.

DMD#10439

- Shimada T, Yamazaki H, Mimura M, Inui Y, and Guengerich FP (1994) Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: Studies with liver microsomes of 30 Japanese and 30 Caucasians. *J Pharmacol Exp Ther* 270:414-423.
- Shimada T and Guengerich FP (1989) Evidence for cytochrome P450_{NF}, the nifedipine oxidase, being the principle enzyme involved in the bioactivation of aflatoxins in human liver. *Proc Natl Acad Sci USA* 86:462-465.
- Silverman RB (1955) Mechanism-based enzyme inactivators, in: *Enzyme Kinetics and Mechanism, Part D: Developments in Enzyme Dynamics* (Purich DL, ed), 249: 240-283, Academic Press, London.
- Suchar LA, Chang RL, Rosen RT, Lech J and Conney AH (1995) High-performance liquid chromatography separation of hydroxylated estradiol metabolites: formation of estradiol metabolites by liver microsomes from male and female rats. *J Pharmacol Exp Ther* 272:197-206.
- Szklarz GD and Halpert JR (1998) Molecular basis of P450 inhibition and activation: implications for drug development and drug therapy. *Drug Metab Dispos* 26:1179-1184.
- Tang J, Cao Y, Rose RL and Hodgson E (2002) In vitro metabolism of carbaryl by human cytochrome P450 and its inhibition by chlorpyrifos. *Chem.-Biol. Interact.* 141:229-241.
- Usmani KA, Rose RL, Goldstein JA, Taylor WG, Brimfield AA and Hodgson E (2002) In vitro human metabolism and interactions of repellent N,N-diethyl-m-toluamide. *Drug Metab Dispos* 30:289-294.

DMD#10439

- Usmani KA, Rose RL and Hodgson E (2003) Inhibition and activation of the human liver microsomal and human cytochrome P450 3A4 metabolism of testosterone by deployment-related chemicals. *Drug Metab Dispos* **31**:384-391.
- Waxman DJ, Attisano C, Guengerich FP, and Lapenson DP (1988) Human liver microsomal steroid metabolism: identification of the major microsomal steroid hormone 6 β -hydroxylase cytochrome P-450 enzyme. *Arch Biochem Biophys* **263**:424-436.
- William PA, Cosme J, Vinkovic DM, Ward A, Angove HC, Day PJ, Vonrhein C, Tickle IJ and Jhoti H (2004) Crystal structures of human cytochrome P450 3A4 bound to metyrapone and progesterone. *Science* **305**:683-686.
- Wilson AS, Davis CD, Williams DP, Buckpitt AR, Pirmohamed M and Park BK (1996) Characterisation of the toxic metabolite(s) of naphthalene. *Toxicology* **114**:233-242.
- Wilson VS and LeBlanc GA (1998) Endosulfan elevates testosterone biotransformation and clearance in CD-1 mice. *Toxicol Appl Pharmacol* **148**:158-168.
- Wilson JD, Foster DW, Kronenberg HM and Larsen PR (1998) Williams Textbook of Endocrinology. Saunders, Philadelphia.
- Yano JK, Wester MR, Schoch GA, Griffin KJ, Stout CD and Johnson EF (2004) The structure of human microsomal cytochrome P450 3A4 determined by X-ray crystallography to 2.05- \AA resolution. *J Biol Chem* **279**:38091-38094.
- Yamazaki H, Shaw PM, Guengerich FP and Shimada T (1998) Roles of cytochromes P450 1A2 and 3A4 in the oxidation of estradiol and estrone in human liver microsomes. *Chem Res Toxicol* **11**:659-665.

DMD#10439

Unnumbered footnote:

This research was supported by US Army Cooperative Agreement DAMD 17-00-2-0008. Part of the studies were presented at the 7th International ISSX meeting in Vancouver, Canada, 2004 and part were presented at the 44th Annual meeting of SOT in New Orleans, 2005.

Numbered footnote:

¹Deceased, Dr Rose died in a tragic accident on May 23rd, 2006.

The person to receive reprint requests:

Dr. Ernest Hodgson

Department of Environmental and Molecular Toxicology

Box 7633

North Carolina State University

Raleigh, NC 27695

DMD#10439

Legends for Figures

Figure 1. Structures of all chemicals tested as inhibitors of estradiol metabolism.

Figure 2. Metabolism of E₂ to 2-OHE₂ by single-donor HLM in the presence and absence of ketoconazole. Solid bars = no ketoconazole and open bars = 2 μM ketoconazole.

Figure 3. Effects of deployment-related and other chemicals on E₂ metabolism by pooled HLM.

* Statistically significant different when compared with control ($P < 0.01$).

Figure 4. Effects of deployment-related and other chemicals on E₂ metabolism by CYP3A4.

* Statistically significant different when compared with control ($P < 0.01$).

◆ Statistically significant different when compared with control ($P < 0.05$).

Figure 5. Effects of deployment-related and other chemicals on E₂ metabolism by CYP1A2.

* Statistically significant different when compared with control ($P < 0.01$).

Figure 6. Michaelis-Menten plots for the inhibition of CYP3A4 mediated E₂ hydroxylation by chlorpyrifos (A) and fonofos (B). Kinetic constants for the uninhibited (no inhibitor present) enzyme are shown in Table 1.

Figure 7. Michaelis-Menten plots for the inhibition of CYP1A2 mediated E₂ hydroxylation by chlorpyrifos (A) and fonofos (B). Kinetic constants for the uninhibited (no inhibitor present) enzyme are shown in Table 1.

Figure 8. Time and concentration-dependent inactivation of CYP3A4 metabolism of E₂ by CPS.

A. Correlation between preincubation time with various concentrations of CPS and CYP3A4 activity. B. Plot of inactivation rate constant as a function of CPS concentration to determine k_{inact} and K_I values.

DMD#10439

Figure 9. Michaelis-Menten plots for the inhibition of CYP1A2 mediated E₂ hydroxylation by carbaryl (A), 4-hydroxycarbaryl (B), and carbaryl methylol (C). Kinetic constants for the uninhibited (no inhibitor present) enzyme are shown in Table 1.

Figure 10. Michaelis-Menten plot for the inhibition of CYP1A2 mediated E₂ hydroxylation by naphthalene. Kinetic constants for the uninhibited (no inhibitor present) enzyme are shown in Table 1.

Figure 11. Michaelis-Menten plot for the inhibition of CYP1A2 mediated E₂ hydroxylation by 1-naphthol.


DMD#10439

Table 1. Kinetic parameters for 2-hydroxylation of E₂ by pooled HLM, CYP3A4, and CYP1A2


	K_m	V_{max}	V_{max}/K_m
	μM	nmol/min/mg protein	$\mu\text{l}/\text{min}/\text{mg}$ protein
HLM	42.8 ± 9.40	0.32 ± 0.02	7.5
		nmol/min/nmol CYP	$\mu\text{l}/\text{min}/\text{nmol}$ CYP
CYP3A4	23.7 ± 3.00	6.9 ± 0.35	291
CYP1A2	27.5 ± 2.16	17.4 ± 0.54	633

Values are expressed as mean \pm S.E.M. ($n = 3$).


Fig. 1




Permethrin




Deltamethrin




Carbaryl




Carbofuran




Pyridostigmine




Naphthalene




Fipronil




Phorate



Chlorpyrifos



Fonofos



Imidacloprid

Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.




Figure 8.





Figure 9.




Fig. 10.




Figure 11.

