

Review

Breast Cancer Resistance Protein (ABCG2) in Clinical Pharmacokinetics and Drug Interactions: Practical Recommendations for Clinical Victim and Perpetrator Drug-Drug Interaction Study Design

Caroline A. Lee, Meeghan A. O'Connor, Tasha K. Ritchie, Aleksandra Galetin, Jack A. Cook, Isabelle Ragueneau-Majlessi, Harma Ellens, Bo Feng, Mitchell E. Taub, Mary F. Paine, Joseph W. Polli, Joseph A. Ware, and Maciej J. Zamek-Gliszczynski

Drug Metabolism and Pharmacokinetics, QPS LLC, RTP, NC (CAL)

Drug Metabolism and Pharmacokinetics, Boehringer-Ingelheim Pharmaceuticals, Inc., Ridgefield, CT (MAO'C, MET)

Centre for Applied Pharmacokinetic Research, Manchester Pharmacy School, The University of Manchester, Manchester, UK (AG)

Pharmacokinetics and Drug Metabolism, Pfizer Inc., Groton, CT (BF)

Clinical Pharmacology, Global Innovative Pharma Business, Pfizer Inc., Groton, CT (JAC)

School of Pharmacy, University of Washington, Seattle, WA (IRM, TKR)

Drug Metabolism and Pharmacokinetics, GlaxoSmithKline, RTP, NC (MJZG, JWP) and King of Prussia, PA (HE)

College of Pharmacy, Washington State University, Spokane, WA (MFP)

Clinical Pharmacology, Genentech, South San Francisco, CA (JAW)

Current Affiliation:

Drug Metabolism and Pharmacokinetics, Ardea Bioscience Inc., San Diego, CA (CAL)

Running Title: BCRP clinical DDI study design recommendations (46/60 characters)

Corresponding Author:

Maciej J. Zamek-Gliszczyński, Ph.D.

GlaxoSmithKline, Drug Metabolism and Pharmacokinetics

5 Moore Drive, P.O. Box 13398

RTP, NC 27709-3398, USA

e-mail: maciej.x.zamek-gliszczyński@gsk.com

Tel: 1-919-483-6366

Review Format

Abstract: 247/250 words

Body Text: (review: no word limit)

References: (review: no reference limit)

Number of Tables: 7

Number of Figures: 1

Abbreviations: AUC, area under the concentration-time curve; BCRP (ABCG2), Breast Cancer Resistance Protein; DDI, drug-drug interaction; EMA, European Medicines Agency; FDA, Food and Drug Administration; IR, immediate release; ITC, International Transporter Consortium; MRP, multidrug resistance associated protein; NAT, *N*-acetyltransferase; OATP, organic anion transporting polypeptide; P-gp, P-glycoprotein; SNP, single nucleotide polymorphism; TKIs, tyrosine kinase inhibitors; PPIs, proton pump inhibitors;

ABSTRACT (247/250 words)

BCRP limits intestinal absorption of low-permeability substrate drugs and mediates biliary excretion of drugs and metabolites. Based on clinical evidence of BCRP-mediated drug-drug interactions (DDIs) and the c.421C>A functional polymorphism affecting drug efficacy and safety, both the FDA and EMA recommend preclinical evaluation, and when appropriate, clinical assessment of BCRP-mediated DDIs. Although many BCRP substrates and inhibitors have been identified in vitro, clinical translation has been confounded by overlap with other transporters and metabolic enzymes. Regulatory recommendations for BCRP-mediated clinical DDI studies are challenging, as consensus is lacking on the choice of the most robust and specific human BCRP substrates and inhibitors and optimal study design. This review proposes a path forward based on a comprehensive analysis of available data. Oral sulfasalazine (1000 mg, IR tablet) is the best available clinical substrate for intestinal BCRP, oral rosuvastatin (20 mg) for both intestinal and hepatic BCRP, and intravenous rosuvastatin (4 mg) for hepatic BCRP. Oral curcumin (2000 mg) and lapatinib (250 mg) are the best available clinical BCRP inhibitors. To interrogate the worst-case clinical BCRP DDI scenario, study subjects harboring the BCRP c.421C/C reference genotype are recommended. In addition, if sulfasalazine is selected as the substrate, subjects having the rapid acetylator phenotype are recommended. In the case of rosuvastatin, subjects with the OATP1B1 c.521T/T genotype are recommended, together with monitoring of rosuvastatin cholesterol lowering effect at baseline and DDI phase. A proof-of-concept clinical study is being planned by a collaborative consortium to evaluate the proposed BCRP DDI study design.

Introduction

Breast cancer resistance protein (BCRP, ABCG2) is an ATP-binding cassette efflux transporter comprised of a homodimer of two half subunits (Giacomini et al., 2010). BCRP is expressed in the small intestine, liver, blood-brain barrier, testis, placenta and mammary glands (Endres et al., 2006) and limits systemic and organ exposure of relevant substrates. BCRP transports endogenous and exogenous substrates with diverse physicochemical properties, which generally include a positive and/or negative charge, but otherwise span a large range of molecular weight, lipophilicity, and permeability (Mao and Unadkat, 2005; Ni et al., 2010; Robey et al., 2011). However, the impact of BCRP on pharmacokinetics is the most pronounced for compounds with low passive permeability (Poirier et al., 2014).

From a drug disposition perspective, BCRP functions primarily as an apical efflux pump in enterocytes, attenuating intestinal absorption of low-permeability substrate drugs, and as a canalicular efflux pump, transporting substrates from hepatocytes into bile. Unlike intestinal BCRP, which is rate-determining in the absorption of low-permeability substrate drugs, in the liver, uptake is generally rate-determining in hepatic clearance of drugs that predominantly rely on carrier-facilitated transport into the liver prior to excretion into bile via BCRP. However, when a basolateral efflux mechanism is operational (e.g. MRP4), BCRP biliary excretion can become rate-determining in systemic drug clearance when it is impaired to a sufficiently large extent (Pfeifer et al., 2013a; Pfeifer et al., 2013b). Such shifting of the rate-determining step in hepatobiliary clearance is a kinetically complex phenomenon and is best described with both systemic pharmacokinetic data, as well as a measure of hepatic drug exposure, which in humans can be assessed directly (e.g. imaging of a metabolically-stable drug) or indirectly using a hepatic pharmacodynamic biomarker (e.g. cholesterol for statins).

Pharmacogenetic studies identified a c.421C>A (p.Q141, rs2231142) single nucleotide polymorphism (SNP) of BCRP that results in reduced activity (Giacomini et al., 2013). In vitro studies demonstrated that this BCRP variant exhibits maintained mRNA expression, but 50-70% reduced protein expression and function due to enhanced susceptibility to proteasomal degradation (Kondo et al., 2004; Furukawa et al., 2009). Several marketed drugs including sulfasalazine, rosuvastatin, and atorvastatin have increased systemic exposure due to impaired BCRP activity in carriers of this functional polymorphism (Giacomini et al., 2013). For example, rosuvastatin demonstrated 2.4-fold higher exposure in homozygous BCRP c.421A/A individuals compared to homozygous c.421C/C individuals (Keskitalo et al., 2009b). In addition to changes in the pharmacokinetic profile, clinical studies involving BCRP c.421A/A homozygous carriers demonstrated an increased incidence of gefitinib side effects (Cusatis et al., 2006).

BCRP has been demonstrated to be an important mediator of drug-drug interactions (DDIs) in humans (Giacomini et al., 2010). For example, the herbal product curcumin increased sulfasalazine oral exposure 3.2-fold. This clinical DDI has been attributed to BCRP inhibition, because it could be replicated in wild-type, but not *Bcrp*-knockout mice, clearly proving *Bcrp* to be the DDI mechanism (Shukla et al., 2009; Kusuhara et al., 2012).

The clinical evidence of BCRP inhibition as a mechanism of DDIs with potential safety and efficacy consequences prompted the International Transporter Consortium (ITC) to highlight BCRP as an important transporter to evaluate during drug development (Giacomini et al., 2010). Subsequently, the FDA and EMA recommended that all investigational drugs should be evaluated as substrates (victims) or inhibitors (perpetrators) of BCRP, and when warranted, that victim and perpetrator DDI potential should be characterized in humans (FDA, 2012; EMA

2012). The exception to the BCRP substrate evaluation is when the drug candidate meets biopharmaceutical classification system 1 criteria, where intestinal solubility and permeability are too high for oral absorption to be attenuated by BCRP efflux (FDA, 2012; EMA 2012). These regulatory recommendations are reasonable and justified by overall clinical evidence supporting BCRP as a contributor to, although typically not the sole site of, victim and perpetrator DDIs.

Many drugs from various therapeutic areas have been identified as substrates or inhibitors of BCRP *in vitro*, yet clinical DDIs attributed directly and specifically to BCRP are limited due to overlap with other transport, as well as metabolic, pathways (Poirier et al., 2014). For example, the 8-fold increase in atorvastatin exposure during tipranavir and ritonavir co-administration was attributed mechanistically to inhibition of BCRP, OATP1B1 and CYP3A4 (Pham et al., 2009); however, quantification of the relative contribution of each of these pathways to the overall DDI is challenging. Since well-characterized clinical BCRP substrate probes and inhibitors are lacking, DDIs attributed solely to BCRP are rare. As a consequence, a consensus is lacking on how to optimally evaluate clinical BCRP victim and perpetrator DDI potential.

Recognizing the challenges faced by the pharmaceutical industry regarding the clinical evaluation of BCRP-mediated drug disposition and DDIs, the intent of this review is 1) to identify the best available clinical probes and inhibitors that can be used to understand the contribution of BCRP to intestinal, hepatic, or overall systemic drug disposition in humans, and 2) to propose a clinical study design to test the validity of the selected substrate/inhibitor pairs to evaluate BCRP-mediated DDIs, as well as delineate organ-specific BCRP contributions in humans.

Clinical Relevance of BCRP Polymorphisms

Impaired BCRP activity is associated with a non-synonymous SNP, c.421C>A (Furukawa et al., 2009). Clinically, this polymorphism is linked to increased exposure to rosuvastatin, sunitinib, an active metabolite of leflunomide, sulfasalazine, flavopiridol, diflomotecan and gefitinib (Giacomini et al., 2013). For example, homozygous carriers of the variant allele (c.421A/A) showed up to a 3.5-fold increase in sulfasalazine oral exposure compared to homozygous carriers of the reference allele (c.421C/C) (Yamasaki et al., 2008; Kusuhara et al., 2012). Clinical examples of elevated statin exposure in homozygous carriers of the c.421C>A SNP range from 1.7-fold increase for both atorvastatin and fluvastatin to 2.4-fold increase for rosuvastatin (Keskitalo et al., 2009a; Keskitalo et al., 2009b).

In general, the effect of the BCRP c.421C>A variant on the pharmacokinetics of most BCRP substrates has been relatively modest; however, a major limitation of many of these studies is the low number of subjects homozygous for this SNP due to low population frequency of the recessive allele. For example, intravenous diflomotecan exhibited 3-fold higher intravenous exposure in c.421C/A (n=5) versus c.421C/C carriers (n=15) but no change in AUC following oral drug administration (Sparreboom et al., 2004), while rosuvastatin exposure was 1.8-fold higher in a combined cohort of c.421C/A and A/A (n=5 C/A and n=2 A/A) versus c.421C/C carriers (n=7) (Zhang et al., 2006) (Table 1). In contrast, no effect of the c.421C/A SNP was observed on the oral pharmacokinetics of the following BCRP substrates: nitrofurantoin, lamivudine, pravastatin, pitavastatin, and intravenous irinotecan (Ferrante et al., 1991; de Jong et al., 2004; Han et al., 2007; Ieiri et al., 2007; Jada et al., 2007; Adkison et al., 2008; Keskitalo et al., 2009a; Zhou et al., 2013). Lack of effect of BCRP c.421C>A on pravastatin exposure (Keskitalo 2009a) stands in contrast to rosuvastatin (which has comparable

low passive permeability to pravastatin) and supports the hypothesis that P-gp and MRP2 play more prominent roles than BCRP in pravastatin absorption.

The above examples highlight the critical importance of thoughtful design of BCRP pharmacogenetic studies when intended to be used in lieu of traditional DDI inhibitor studies to delineate the victim DDI potential of a BCRP substrate drug. As demonstrated by the above examples, the number of study subjects in typical DDI studies is not high enough to randomly enroll enough carriers of the BCRP c.421A/A and c.421C/A genotypes, making a post-hoc comparison of BCRP genotype effects impossible. As such, when BCRP pharmacogenetic studies are intended to be used as a substitute for a DDI study with a BCRP inhibitor, subjects must be genotyped during enrollment, and the study should be powered with an equal and adequate number of subjects in each of the three groups (C/C, C/A, A/A). The two sulfasalazine BCRP pharmacogenetic studies by Yamasaki et al. and Adkison et al. provide good examples of how to proactively investigate victim DDI potential of a BCRP substrate drug in a properly designed study (Yamasaki et al., 2008; Adkison et al., 2010) (Table 2).

Recently, several null allelic mutations in the BCRP gene were reported to correlate with the Junior(a-) blood group antigen (Saison et al., 2012; Zelinski et al., 2012). The Junior(a-) phenotype has been associated with clinically relevant adverse events, including transfusion reactions and fetal anemia. This phenotype has been detected at a low frequency in individuals of northern European and Romani ancestry and in Bedouin Arabs, but at a relatively higher frequency in Asians, particularly Japanese (0.02-1.7%) (Zelinski et al., 2012; Tanaka et al., 2014). All of these mutations result in non-functional protein, thereby identifying individuals with the Junior(a-) blood group as a population lacking BCRP activity.

As BCRP deficiency has recently been elucidated as the mechanism underlying the Junior(a-) phenotype, these populations have not yet been used to study BCRP-mediated drug disposition in humans. Initial investigations into the safety and efficacy implications of BCRP deficiency demonstrated conflicting results with respect to elevated blood concentrations of uric acid (no change and hyperuricemia) (Matsuo et al., 2009; Saison et al., 2012); however, uric acid disposition is complex and involves several transporters besides BCRP, thus these conflicting findings are not entirely surprising (Reginato et al., 2012).

Theoretically, the Junior(a-) population may provide an opportunity to tease out the contribution of BCRP by comparing the effect to healthy controls; however, such a clinical study paradigm remains to be established. In practice, these studies would have to be conducted in Japan, where the prevalence of the Junior(a-) phenotype is sufficiently high and the phenotype and study subject recruitment is enabled by tracking of this phenotype by blood banks.

Potential Clinical BCRP Probe Substrates

Based on studies with BCRP-transfected polarized cell lines and membrane vesicles generated from BCRP over-expressing cells, many marketed drugs and dietary/natural constituents have been identified as substrates of BCRP. A list of 81 substrates was evaluated based on reported efflux ratios >2 in cell monolayers and/or reported K_m values $<10 \mu\text{M}$ (Table 3). These substrates have a broad range of physicochemical properties, with passive permeability ranging from low to high ($0.07\text{-}21.8 \times 10^{-6} \text{ cm/s}$). Many BCRP substrates are extensively metabolized, whereas others are excreted unchanged. The contribution of other transporters to the disposition of these substrates is common (Poirier et al., 2014). In contrast to the numerous BCRP substrates identified in vitro, the number of potential clinical substrates is limited. Theoretically, the ideal clinical probe substrate should be selective for BCRP with minimal contribution from other transporters and metabolizing enzymes in order to minimize the complexity in data interpretation and enable extrapolation to other BCRP substrate drugs. Analogous to digoxin safety studies for P-glycoprotein inhibitor drugs, if the candidate BCRP probe substrate has a narrow therapeutic window and/or identified safety concerns, then the DDI will have direct clinical relevance by informing dose adjustment for the substrate drug. An example of the latter is the case of statins where increased exposure can result in myopathy (Egan and Colman, 2011). Several potential clinical BCRP substrates for consideration are described below.

Sulfasalazine. Sulfasalazine, an anti-inflammatory and immunomodulatory drug used to treat ulcerative colitis, rheumatoid arthritis, and Crohn's disease (Azulfidine Prescribing Information, 2011), has been proposed as a clinical probe for intestinal BCRP. Sulfasalazine is restricted largely to the gastrointestinal tract due to low permeability, low solubility, and efficient intestinal BCRP efflux. As such, oral bioavailability at therapeutic doses is low ($<10\%$) and

highly variable. The low fraction of a sulfasalazine oral dose that is absorbed into the systemic circulation is mostly excreted unchanged in urine (via glomerular filtration) and bile (Azulfidine Prescribing Information, 2011; Giacomini et al., 2010).

Sulfasalazine is metabolized extensively by bacteria in the distal small intestine and colon to sulfapyridine, which is relatively well absorbed and cleared predominantly by *N*-acetyltransferase (NAT) 2. Inhibition of intestinal BCRP reduces sulfapyridine formation by increasing sulfasalazine absorption, which in turn decreases sulfasalazine exposure to intestinal bacteria. As such, the metabolite-to-parent AUC ratio is a more sensitive indicator of intestinal BCRP modulation in humans than parent sulfasalazine AUC, but only in intermediate and rapid acetylator phenotypes (Yamasaki et al., 2008). Specifically in NAT2 rapid acetylators, the range of BCRP clinical effect is ~10-fold as measured by the decrease in sulfapyridine-to-sulfasalazine AUC ratio, while the clinical magnitude of parent sulfasalazine AUC increase is ≤ 3.5 -fold (Yamasaki et al., 2008).

A wealth of preclinical data support sulfasalazine as an essentially specific probe for intestinal BCRP, although a potentially small contribution from P-gp and MRP2 has been proposed (Zaher et al., 2006; Dahan and Amidon, 2009; Zamek-Gliszczyński et al., 2012). Overall, Bcrp-knockout rodent data support BCRP limiting sulfasalazine oral bioavailability by at least an order of magnitude, while MRP2 and perhaps P-gp attenuate intestinal absorption by a lesser extent (≤ 2 to 3-fold). However, the fact that sulfasalazine intestinal absorption is not strictly limited by BCRP cannot be ignored in the interpretation of sulfasalazine interaction data, especially considering possible overlap between BCRP and P-gp, as well as BCRP and MRP2, inhibitors.

The possibility of uptake transporter(s) contributing to sulfasalazine absorption has been proposed, and intestinal OATP2B1 has been shown to transport sulfasalazine in vitro (Kusuhara et al., 2012; Tomaru et al., 2013). Intestinal uptake could be an important consideration concerning interpretation of DDI data should a decrease in sulfasalazine exposure be observed. Furthermore, saturation of uptake transporter(s) may underlie pharmacokinetic differences between sub-therapeutic and therapeutic doses, which cannot be explained by drug solubility alone (Kusuhara et al., 2012).

Topotecan. Topotecan is a topoisomerase inhibitor for second-line treatment of patients with metastatic ovarian carcinoma and small cell lung cancer, available in both intravenous and oral formulations (HYCAMTIN[®] Prescribing Information (06/18/2014; http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/200199s0031bl.pdf and http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/020981s0061bl.pdf). Although oral absorption is limited by intestinal BCRP (Jonker et al., 2000; Kruijtzter et al., 2002), topotecan is a substrate for other efflux transporters as well, including P-gp and MRP2 (Lin et al., 2011). Nonetheless, in P-gp-knockout mice, the nonspecific BCRP and P-gp inhibitor, elacridar, increased topotecan oral exposure 6-fold, thus demonstrating a major role of Bcrp in limiting topotecan oral absorption (Jonker et al., 2000). Co-administration of elacridar with topotecan to cancer patients increased plasma exposure 2- to 3-fold (Kruijtzter et al., 2002), which is the upper limit of the possible increase in humans, where baseline bioavailability is 40% (HYCAMTIN[®] Prescribing Information (06/18/2014; http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/020981s0061bl.pdf). The effect of BCRP c.421C>A SNP on the pharmacokinetics of topotecan showed 1.3-fold higher oral bioavailability in c.421C/A versus c.421C/C individuals, a modest effect that would be expected

to increase in patients who are c.421A/A homozygous but were not included in this study (Sparreboom et al., 2005). Although topotecan has some attractive characteristics as a BCRP probe substrate, due to its safety profile as a cytotoxic agent, topotecan is impractical for routine use in healthy volunteer BCRP DDI studies.

Statins. The physicochemical properties of statins, such as atorvastatin, cerivastatin, fluvastatin, pitavastatin, rosuvastatin, and pravastatin, are similar with respect to molecular weight, pKa, and anionic charge, but differ in terms of lipophilicity. Consequently, the involvement of a variety of metabolizing enzymes and drug transporters in statin disposition differs based on a statin's lipophilic versus hydrophilic character (Elsby et al., 2012; Shitara et al., 2013). Likewise, passive permeability varies with lipophilicity, and on one extreme is low for pravastatin and rosuvastatin, while on the other is high for cerivastatin and fluvastatin (Jones et al., 2012; Menochet et al., 2012; Li et al., 2014). With respect to clearance pathways, cerivastatin, fluvastatin, simvastatin lactone, and atorvastatin are extensively metabolized, whereas pravastatin, rosuvastatin, and pitavastatin undergo minimal metabolism and are cleared mainly through biliary and urinary excretion (Elsby et al., 2012). All statins to a varying degree are substrates for hepatic uptake transporters and some are BCRP substrates (Elsby et al. 2012).

Studies in Bcrp-knockout mice have demonstrated an important role for Bcrp in the hepatobiliary disposition of pitavastatin and rosuvastatin (Hirano et al, 2005; Kitamura et al., 2008). Following intravenous infusion to Bcrp-knockout mice, the biliary excretion rate of both drugs was reduced by approximately an order of magnitude relative to wild-type mice. In contrast, the systemic clearance of pitavastatin was not affected by Bcrp knockout, whereas that of rosuvastatin was reduced by approximately 2-fold. Likewise clinically, the BCRP c.421 C>A polymorphism had a negligible effect on pitavastatin pharmacokinetics (Ieiri et al., 2007).

Pitavastatin is a sensitive clinical probe for OATP1B1 DDI study (Prueksaritanont et al., 2014), but it is not recommended as a clinical probe in BCRP DDI study because of a minimal role of this transporter in its absorption, disposition and clearance. Although the most pronounced effects on C_{max} is reported for fluvastatin and rosuvastatin, the high permeability of fluvastatin combined with less compelling BCRP pharmacogenetic effects and DDI potential relative to rosuvastatin preclude consideration of this statin as a sensitive probe for intestinal BCRP-mediated DDI studies (Hirano et al., 2005; Ieiri et al., 2007; Elsby et al., 2012).

Rosuvastatin has limited ability to permeate cellular membranes via passive diffusion due to low lipophilicity ($\text{LogD}_{7.4} = -0.33$) (Kitamura et al., 2008; Menochet et al., 2012). Intestinal absorption is ≤ 3.2 -fold limited by BCRP in both preclinical species and humans (Karibe et al., 2014; Prueksaritanont et al., 2014). Rosuvastatin is primarily cleared by the liver (72%) and to a lesser extent by the kidneys (28%) (Elsby et al., 2012). Several hepatic uptake transporters (OATP1B1, OATP1B3, and sodium-taurocholate co-transporting polypeptide) contribute to its distribution in the liver, while BCRP and MRP2 secrete rosuvastatin into the bile, such that it can undergo extensive enterohepatic recirculation (Ho et al., 2006; Kitamura et al., 2008; van de Steeg et al., 2013). In addition to glomerular filtration, the tubular secretion of rosuvastatin is mediated by organic anion transporter 3 uptake and efflux by MRP2.

Based on pharmacogenetic assessment of all the statins, the disposition of rosuvastatin is affected primarily by BCRP polymorphisms in vivo, as summarized in Table 1. Because the BCRP c.421 C>A variant does not represent a complete knockout of BCRP activity, it is likely that more profound changes may be observed if rosuvastatin was used in DDI studies with a potent BCRP inhibitor or in the Junior(a-) population, which lacks BCRP altogether. Moreover, reduced biliary transport of rosuvastatin via BCRP (due to either inhibition or a polymorphism)

may lead to more pronounced changes in hepatic concentrations compared to systemic concentrations. Taken together, the evidence for BCRP as a major contributor to rosuvastatin disposition relative to the other statins is the strongest. However, the use of rosuvastatin as a BCRP DDI probe requires an appropriate clinical study design and careful consideration of perpetrator properties, as discussed below.

Clinical Evidence-Based Recommendations for BCRP Substrate Selection

Intestinal Probe Substrate. Four clinical studies have employed oral sulfasalazine as an intestinal BCRP probe to investigate the pharmacokinetic effects of BCRP inhibitors and the c.421C>A SNP in humans (Urquhart et al., 2008; Yamasaki et al., 2008; Adkison et al., 2010; Kusuhara et al., 2012) (Table 2 and Table 4). Although the number of human DDI studies is limited, important lessons can be drawn for future study design, specifically with respect to subject population, dose, formulation, and study endpoints. These considerations are discussed in detail below.

Population variability in sulfasalazine exposure can be large, as high as 81-fold (Adkison et al., 2010) (Table 4). As such, it is critical that clinical sulfasalazine interaction studies use a crossover design. To maximize the observed human intestinal BCRP interaction effects, study subjects with high BCRP and NAT2 activities should be enrolled, i.e., BCRP c.421C/C homozygous carriers and rapid NAT2 acetylators (Yamasaki et al., 2008). Rapid NAT2 acetylators provide the greatest sensitivity (~10-fold) for determination of clinical DDIs in terms of the sulfapyridine/sulfasalazine AUC ratio, followed by intermediate acetylators (~3-4-fold) versus slow acetylators, who show no apparent change in the ratio (Yamasaki et al., 2008). Notably, an unexpected increase in the sulfapyridine/sulfasalazine AUC ratio, or a decrease not commensurate with the increase in oral sulfasalazine exposure, may be indicative of NAT2 inhibition, which should be further investigated. In summary, intestinal BCRP DDI studies with sulfasalazine should enroll BCRP c.421C/C homozygous, rapid acetylator subjects and measure both parent and sulfapyridine oral exposure. As discussed in the preceding section, the sulfapyridine/sulfasalazine AUC ratio decrease appears to be a more sensitive measure of intestinal BCRP inhibition than parent drug exposure increase, and it is recommended that the DDI be reported using both endpoints.

The choice of sulfasalazine dose and formulation used for intestinal BCRP interaction studies also merits careful consideration (Table 2). BCRP pharmacogenetic and inhibition effects on oral sulfasalazine exposure were readily apparent and statistically significant following oral administration of a 2000 mg dose as immediate release (IR) tablets (Yamasaki et al., 2008; Kusuhara et al., 2012). In contrast, following oral administration of 500 mg sulfasalazine enteric coated tablets, systemic pharmacokinetics were not altered by BCRP pharmacogenetics or by oral co-administration of the BCRP inhibitor, pantoprazole (Adkison et al., 2010). The exact mechanistic reasons for why sulfasalazine administered as enteric coated tablets is not sensitive to BCRP modulation remain unknown. One plausible explanation is that this type of formulation releases drug further down the GI tract, where relative BCRP expression is up to 2-fold lower (Englund et al., 2006; Enokizono et al., 2007b; MacLean et al., 2008) and much less concentrated per unit of surface area (Pang, 2003), such that sulfasalazine is released past BCRP in the upper small intestine (Adkison et al., 2010). Although the extended release formulation appears to be the most likely culprit for the lack of BCRP impairment effect in the Adkison et al. (2010) study, because three independent clinical studies using sulfasalazine immediate release tablets or suspension showed consistent pharmacokinetic changes (Urquhart et al., 2008; Yamasaki et al., 2008; Kusuhara et al., 2012), further work is needed to provide an unequivocal mechanistic explanation for the negative findings of Adkison et al. (2010). Other formulations have been evaluated, including an oral suspension prepared from crushed IR tablets. At a therapeutic dose of 1000 mg, oral exposure was 1.6-fold higher following administration of the suspension versus IR tablet but was not statistically significant (Urquhart et al., 2008; Kusuhara et al., 2012). However, following oral administration of a 0.1 mg suspension (microdose) versus a 2000 mg IR tablet, the dose-normalized exposure was an order of

magnitude higher for the microdose suspension compared to the IR tablet, an effect possibly attributed primarily to potential saturation of the speculated intestinal absorptive transporter(s) at the therapeutic dose and secondarily to solubility effects (Kusuhara et al., 2012). Since the apparent fraction absorbed is greater for the microdose suspension, the magnitude of the intestinal BCRP DDI is not as large as that observed for the therapeutic IR tablet (AUC ratio of 2.0 vs. 3.2) (Kusuhara et al., 2012).

Total daily doses of up to 2000 mg appear to be reasonably well tolerated in most subjects; however, gastric side effects are common, with about one-third of subjects experiencing these side effects to some degree and increasing with dose level (Azulfidine Prescribing Information, 2011; Kusuhara et al., 2012). Serious side effects such as Stevens-Johnson syndrome can occur and have been observed following a single dose. To avoid unnecessary adverse event complications, DDI studies involving sulfasalazine should exclude subjects with known “sulfa” allergy, viral infections, a weakened immune system, a history or family history of Stevens-Johnson syndrome, or with the HLA-B *1502 allele. Based on currently available clinical experience, 1000 mg administered orally as IR tablets is the most optimal and practical sulfasalazine dosing regimen for intestinal BCRP DDI studies. Considering the maximal (<4-fold) clinical increase in sulfasalazine exposure due to BCRP inhibition, clinical exposure following administration of the 1000 mg dose would not exceed the exposure observed following administration of the highest approved dose level of 4000 mg/day. Finally, while the subject genotyping recommendations for robust response with sulfasalazine as a marker of intestinal BCRP activity (BCRP c.421 C/C; NAT2 rapid acetylator phenotype, non-carriers of HLA-B *1502 allele) might seem onerous, these are all routine clinical genotyping tests, which add

insignificant cost to the overall study and do not present a serious enrollment hurdle, because the predominant genotype is recommended for all three genes.

Systemic Probe Substrate. Oral rosuvastatin represents a sensitive probe to assess the magnitude of inhibition of both intestinal and hepatic BCRP. Allowing for assessment of the worst-case DDI scenario, a rosuvastatin BCRP DDI study should enroll subjects homozygous for the BCRP reference allele c.421 C/C (most prevalent in non-Asian ethnicities) and exclude any carriers of OATP1B1 c.521T/C or C/C. Consideration of CYP2C9 polymorphism (rosuvastatin undergoes minimal metabolism via this enzyme), is not of relevance because CYP2C9*1/*3 or *3/*3 was reported to have no effect on rosuvastatin (or its metabolite) PK or PD (Table 1). Alternatively, a clinical study should consider the genotype of the subjects enrolled and investigate the impact of multiple genetic covariates (e.g., co-existence of OATP1B1 and BCRP polymorphisms in the same subject) together with ethnicity data on the BCRP DDI sensitivity. An adequately powered study in terms of sample size would allow for identification of multiple high-risk combination patterns that could cause an increase in rosuvastatin exposure in conjunction with reduced BCRP activity, as recently reported for simvastatin (Tsamandouras et al., 2014).

Considering the complexity of rosuvastatin disposition and potential for enterohepatic recycling, it is evident that interpretation of increased rosuvastatin exposure in the DDI study following oral administration may be confounded by multiple factors. A key factor is the inability to differentiate between the contribution of intestinal and hepatic BCRP to the overall magnitude of a rosuvastatin-based interaction. Perpetrators that are potent BCRP as well as OATP inhibitors represent an additional issue of concern in that either or both can result in elevated blood levels of rosuvastatin. BCRP DDI study design involving oral and intravenous

rosuvastatin formulations in the same subjects would allow delineation of the contribution of intestinal and hepatic BCRP. Based on the modeling and simulation work for compounds with comparable disposition to rosuvastatin (Watanabe et al., 2009; Chu et al., 2013b), the expectation is that the inhibition of hepatic BCRP may not cause significant changes in the plasma exposure, whereas drug accumulation in the hepatocytes and increase in liver exposure would occur. To account for this, cholesterol lowering effect of rosuvastatin will also be monitored in the control and inhibitor phase of the clinical studies, as a direct reflection of any changes in rosuvastatin liver exposure.

Rosuvastatin is commonly administered at an oral dose of 5–40 mg/day, with the majority of the clinical DDIs or pharmacogenetic studies conducted using a 10 or 20 mg dose (Simonson et al., 2004; Keskitalo et al., 2009b; Allred et al., 2011). Although a rosuvastatin intravenous formulation is not available currently, clinical data after infusion of 8 mg rosuvastatin have been reported as safe and well tolerated (Martin et al., 2003).

Rosuvastatin administered orally at a dose of 20 mg represents a good clinical probe for both hepatic and intestinal BCRP function (Table 1). The selected oral dose should be sufficient to capture BCRP DDI, while minimizing any potential safety concerns associated with increased exposure at the upper limit of known clinical DDIs. Selection of a 20 mg oral rosuvastatin dose for the clinical DDI study can theoretically be paired with a 4 mg intravenous dose, reflecting the reported 20% oral bioavailability (Martin et al., 2003). In the case of intravenous rosuvastatin, inhibition of hepatic BCRP would interrupt enterohepatic recycling; monitoring of the changes in rosuvastatin pharmacodynamic effect relative to the control data would allow assessment of its contribution. Rosuvastatin DDI studies should be conducted in a cross-over design, ideally in subjects with the highest BCRP and OATP activity, and report rosuvastatin pharmacokinetics

and cholesterol lowering effect at baseline and in the presence of a perpetrator. Comparison of the increase in rosuvastatin exposure after intravenous and oral administration and any changes in the PD effect will enable differentiation of the magnitude of hepatic and intestinal BCRP interaction.

Potential Clinical BCRP Inhibitors

Identifying a drug or diet-derived/natural product that can be used as a selective BCRP inhibitor in a clinical DDI study is challenging, as currently available compounds are not selective for BCRP and inhibit other transporters and drug metabolizing enzymes (Leggas et al., 2006; Lainey et al., 2012; Fujita et al., 2013; Jamei et al., 2014; Poirier et al., 2014; Shukla et al., 2014). Upon careful consideration of inhibition data generated from various in vitro and in vivo studies, there is a high likelihood that any known BCRP inhibitor will inhibit other membrane transporters or metabolic enzymes. Given these limitations, the most potent and selective BCRP inhibitors were identified that complement the proposed BCRP substrates for their potential use in clinical DDI studies. Such inhibitors were selected through a comprehensive literature search using the University of Washington Drug Interaction Database (<http://www.druginteractioninfo.org>) and the University of California, San Francisco-FDA TransPortal (<http://bts.ucsf.edu/fdatransportal>).

Identifying Potential BCRP Inhibitors. Known BCRP inhibitors with IC_{50} (or K_i) values $<10 \mu\text{M}$ were identified that could potentially be used in a clinical DDI study (Table 5). Compounds for which the IC_{50} could not be obtained from the literature or considered impractical for use in a clinical study (e.g., cyclosporine) were not considered further. C_{max} values for the identified BCRP inhibitors were used to determine $[I_1]/IC_{50}$ values, which were then used to predict the potency of systemic BCRP inhibition (FDA, EMA 2012). Similarly, maximum intestinal concentrations, estimated as the highest approved dose/250 mL, were used to calculate $[I_2]/IC_{50}$ values, which were used to predict the potency of intestinal BCRP inhibition (Supplemental Table 1) (FDA, EMA 2012). If only a K_i was available, the IC_{50} was assumed to be equivalent to $2xK_i$ when substrate concentration is equal to the K_m (Zhang et al.,

2008). Drugs for which literature data supported inhibition of BCRP *in vivo* were included in the final assessment of potential inhibitors of systemic and intestinal BCRP. To undergo further consideration, a compound had to meet our criteria set at $[I_1]/IC_{50} > 1$ or $[I_2]/IC_{50} > 20$ for a high likelihood of clinical inhibition, which had to be supported by literature evidence that the compound either inhibits or is a BCRP substrate *in vivo* (i.e. a competitive inhibitor), BCRP inhibitors were rank ordered from highest to lowest $[I_1]/IC_{50}$ (Table 6) and $[I_2]/IC_{50}$ (Table 7).

In addition to a rank ordering of systemic and intestinal BCRP inhibition potential, a detailed assessment of the inhibition potential of other transporters and drug metabolizing enzymes was considered (Tables 6 and 7). The comprehensiveness of these tables reflects the difficulty encountered when selecting a viable substrate-inhibitor pair, as multiple inhibitory interactions occurring simultaneously will confound data interpretation concerning specific BCRP involvement in substrate pharmacokinetics. Such potential nonselective interactions were carefully considered, after which most identified BCRP inhibitors were eliminated. The final selected BCRP inhibitors have inherent pros and cons, which are detailed below.

Sulfasalazine. Sulfasalazine is a BCRP substrate with limited absorption due to efficient intestinal BCRP efflux into the lumen, as well as low permeability and low solubility (Azulfidine Prescribing Information, 2011). Being a substrate, sulfasalazine may act as a competitive inhibitor of BCRP, with reported IC_{50} values as low as 0.46 μM (Elsby et al., 2011). The C_{max} range for a single 3 to 4 gram dose of sulfasalazine is 38-79 μM (Goodman et al., 2001), resulting in a maximum $[I_1]/IC_{50}$ value of 170 (Table 6), and an $[I_2]/IC_{50}$ value of 87,000 (Table 7). As such, sulfasalazine could be used as a BCRP inhibitor in a clinical study when paired with a BCRP substrate such as rosuvastatin. However, sulfasalazine also inhibits OATP-mediated uptake, in particular OATP1B1 [$IC_{50} = 0.56 \mu\text{M}$ (Karlgrén et al., 2012b)]. As OATP1B1 is one

of the primary transporters responsible for taking up rosuvastatin into the liver (Simonson et al., 2004; Ho et al., 2006; Pasanen et al., 2007), and the inhibitory potency towards OATP1B1 is similar to that of BCRP, OATP1B1 becomes a major confounding factor. Therefore, sulfasalazine was excluded from consideration for use as a clinical BCRP inhibitor.

Elacridar. Elacridar was originally designed and developed as a reversal agent of P-glycoprotein and BCRP to overcome multidrug resistance. Although, elacridar increases the accumulation of several anticancer agents in preclinical tumor models (Hyafil et al., 1993; Agarwal et al., 2012), it is not a systemically-potent efflux inhibitor at chronically-tolerated doses in humans (Kalvass et al., 2013). However in the intestine, elacridar has been shown to increase topotecan bioavailability via inhibition of BCRP (Kruijtzter et al., 2002; Kuppens et al., 2007). These observations are consistent with elacridar ranking low in terms of systemic BCRP inhibition ($[I_1]/IC_{50} = 1.1$, Table 6), but high with respect to intestinal BCRP inhibition ($[I_2]/IC_{50} = 9,200$, Table 7). The use of elacridar as a clinical intestinal BCRP inhibitor is precluded by the fact that this compound is not currently approved by the FDA or EMA for use in humans. Furthermore, elacridar is an even more potent inhibitor of P-gp than BCRP ($[I_2]/IC_{50} = 180,000$) and would not be suitable for elucidating the intestinal BCRP contribution to limiting the absorption of dual substrates.

Proton Pump Inhibitors. Multiple proton pump inhibitors (PPIs) inhibit BCRP in vitro; however, pantoprazole and rabeprazole are the only two that meet the aforementioned criteria for consideration as clinical inhibitors (Tables 6 and 7); however, clinical and/or preclinical proof-of-concept studies have not been reported. Despite clear advantages to using an inhibitor with an acceptable safety profile such as pantoprazole (Remes-Troche et al., 2014) or rabeprazole (Zouboulis-Vafiadis et al., 2014), these drugs are not as potent BCRP inhibitors as the tyrosine

kinase inhibitors with respect to the $[I_2]/IC_{50}$ values (Table 7) where preclinical proof of concept has been demonstrated (Zaher et al., 2006). Similarly, pantoprazole marginally meets the pre-defined criterion for systemic BCRP inhibition, with an $[I_1]/IC_{50}$ value of 1.2 (Table 6). As such, although using a PPI may minimize the safety risks, other drugs are expected to offer a superior in vivo inhibition profile, rendering data interpretation more straightforward. Finally, any observed DDIs with PPIs are fundamentally confounded by increased gastrointestinal pH (Ware et al., 2013), and a separate intestinal pH control arm must be incorporated into the clinical study with a PPI, which is not a BCRP inhibitor to control for this effect (Adkison et al., 2010).

Natural Products. Several diet-derived/natural product constituents such as curcumin, daidzein, genistein, chrysin, 7,8-benzoflavone and glycyrrhetic acid have been shown to inhibit human BCRP in vitro (Zhang et al., 2005a; Yoshida et al., 2008; Merino et al., 2010; Tamaki et al., 2010). In addition, a mixture of the soy isoflavones genistein and daidzein was shown to inhibit Bcrp in wild-type but not Bcrp-knockout mice, supporting specificity for Bcrp inhibition (Merino et al., 2010). Of these constituents, only curcumin has been evaluated clinically as a BCRP inhibitor. Curcumin is the principal curcuminoid in turmeric, a commonly used spice derived from the rhizomes of *Curcuma longa*, a member of the ginger family. Curcuminoids are polyphenols that render turmeric yellow in color (Sharma et al., 2005). Curcumin is a potent inhibitor of BCRP, with a K_i of 0.7 μ M determined with membrane vesicles over-expressing human BCRP and sulfasalazine as the probe substrate (Kusuhara et al., 2012). Based on the $[I_2]/IC_{50}$, curcumin is predicted to inhibit intestinal BCRP in vivo. Several clinical studies involving oral curcumin have been conducted, with doses ranging from 2000 to 8000 mg (Cheng et al., 2001; Kusuhara et al., 2012). Curcumin was well tolerated, with no signs of toxicity observed during a phase I study in which the subjects were administered oral doses of 8000

mg/day for 3 months (Cheng et al., 2001). At 2000 mg/day, curcumin increased sulfasalazine exposure 3.2-fold, which was attributed to inhibition of intestinal BCRP (Kusuhara et al., 2012). Because curcumin is considered a food or dietary supplement, and thus not subject to strict manufacturing controls as drugs, the purity and reproducibility with respect to constituent composition, stability, and dissolution characteristics is not assured. Accordingly, a natural products chemistry laboratory should be consulted for sourcing and analyzing the test product.

Tyrosine kinase inhibitors. Tyrosine kinase inhibitors (TKIs) are a relatively new generation of anti-cancer drugs that are generally well tolerated. Several TKIs (erlotinib, gefitinib, lapatinib, nilotinib, and sunitinib) are potent BCRP inhibitors with acceptable safety margins to be considered for use in healthy volunteers. One clear advantage to using a TKI as a clinical BCRP inhibitor is that, with the exception of sunitinib, TKIs are predicted to inhibit intestinal as well as systemic BCRP. Based on the rank ordering of inhibitory potency for TKIs, lapatinib is the most potent when considering both intestinal and systemic inhibition of BCRP. Although in vitro data suggest that lapatinib may inhibit P-gp, OATP1B1, and CYP2C8 concomitantly, these interactions are based on a high therapeutic dose of 1250 mg. However, at a sub-therapeutic dose of 250 mg, based on $[I_1]/IC_{50}$ and $[I_2]/IC_{50}$ values (22 and 40,000, respectively), lapatinib is predicted to effectively inhibit BCRP (Table 6 and 7). Although lapatinib carries a black box warning for hepatotoxicity rarely observed during chronic use in cancer patients (TYKERB® product label, 10/18/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/022059s016s017lbl.pdf), it is well tolerated acutely, as doses up to 7000 mg/day have shown no dose limiting toxicities (Chien et al., 2014). At clinical doses, the diarrhea associated with patient treatment would be the only

potential concern; however, this adverse effect typically resolves within 24-48 hours of treatment cessation (MacFarlane and Gelmon, 2011; Chien et al., 2014).

Clinical Evidence-Based Recommendations for BCRP Inhibitor Selection.

Considering the totality of available information, two compounds emerge as potential clinical BCRP inhibitors for use in DDI studies. To discern the effect of intestinal-specific BCRP inhibition, a single 2000 mg dose of curcumin is proposed to extensively inhibit intestinal, but not systemic, BCRP (Tables 6-7). Due to poor absorption, curcumin is not detectable in plasma at this proposed dose (Garcea et al., 2005; Kusuhara et al., 2012), rendering the compound a practical choice for intestinal-specific BCRP inhibition. The second proposed BCRP inhibitor is lapatinib. Compared to a food product such as curcumin, the benefit of using a marketed drug to assess DDIs is that, among other things, a natural products chemistry laboratory need not be consulted. The recommended sub-therapeutic oral dose of 250 mg should have minimal adverse effects and possibly attenuate the impact of inhibition of other transporters and metabolic enzymes (Castellino et al., 2012). At this dose, lapatinib is predicted to inhibit both intestinal and systemic BCRP based on $[I_1]/IC_{50}$ and $[I_2]/IC_{50}$ values (Tables 6-7). One potential concern is that lapatinib may also inhibit OATP1B1, as rosuvastatin uptake is dependent on this transporter (the $[I_1]/IC_{50}$ is 0.29, which is marginally above the cut off value of 0.1). However, lapatinib is highly protein bound (>99%) so circulating levels of free drug available to inhibit OATP1B1 systemically will be significantly less than the total C_{max} reported in Table 6. Taking into account the concentration of lapatinib in the intestine and in the systemic circulation, lapatinib is likely to inhibit BCRP but not OATP1B1.

Recommendations for Clinical BCRP Drug Interaction Study Design

This comprehensive review of potential clinical BCRP probe substrates and inhibitors identified two substrates and two inhibitors to assess BCRP-mediated DDIs in humans. The two recommended probe substrates are sulfasalazine (1000 mg, IR, oral) and rosuvastatin (20 mg oral and 4 mg, intravenous), and the two inhibitors are curcumin (2000 mg, oral) and lapatinib (250 mg, oral). The combination of oral sulfasalazine with either inhibitor provides a substrate-inhibitor pair to assess the extent of a potential intestinal BCRP DDI. The dynamic range of sulfasalazine will provide a measure of the highest extent of an interaction due to inhibitor co-administration. Further, the combination of oral and intravenous rosuvastatin with both inhibitors will enable deconvolution of intestinal versus systemic BCRP contribution to rosuvastatin pharmacokinetics. Based on these aims, a clinical study design is proposed to validate these approaches (Figure 1).

Sulfasalazine Intestinal BCRP DDI. Oral sulfasalazine (1000 mg, IR, single dose) is recommended to evaluate intestinal BCRP-mediated DDIs. The study will be an open-label, 3-period cross-over study involving 10 subjects genotyped for the BCRP reference allele and phenotyped for rapid NAT2 acetylator status prior to enrollment. Sulfasalazine will be administered alone and in combination with oral curcumin (2000 mg, single dose) or lapatinib (250 mg, single dose) (Figure 1). Subjects will undergo three treatments with at least a 1-week washout between treatments as follows: sulfasalazine alone, sulfasalazine plus curcumin, and sulfasalazine plus lapatinib. Pharmacokinetics of both sulfasalazine and sulfapyridine will be determined, with the DDI reported as both an increase in sulfasalazine exposure and decrease in the sulfapyridine/sulfasalazine AUC ratio.

Rosuvastatin Intestinal and Hepatic BCRP DDI. Oral rosuvastatin (20 mg, single dose) is recommended to evaluate intestinal and hepatic BCRP DDIs, whereas intravenous rosuvastatin (4 mg, single dose) is recommended to evaluate hepatic BCRP DDIs. Each rosuvastatin administration route will include an open-label, 3-period cross-over study involving subjects genotyped for the BCRP and OATP1B1 reference alleles prior to enrollment. Rosuvastatin will be administered alone and with oral curcumin (2000 mg, single dose) or lapatinib (250 mg, single dose) (Figure 1). Treatments include intravenous rosuvastatin alone, intravenous rosuvastatin plus oral curcumin, intravenous rosuvastatin plus oral lapatinib, oral rosuvastatin alone, oral rosuvastatin plus oral curcumin, and oral rosuvastatin plus oral lapatinib. Inclusion of cholesterol monitoring will provide insight towards hepatic accumulation of rosuvastatin due to inhibition of hepatic BCRP. Changes in rosuvastatin exposure relative to the control will be the DDI study endpoint.

Conclusions

The regulatory recommendation to evaluate the contribution of BCRP to drug disposition and DDIs in clinical studies has posed challenges across the pharmaceutical industry. These challenges primarily reflect the lack of selective and specific human BCRP substrates and inhibitors, as well as a lack of established and accepted optimal clinical study designs. As described throughout this review, BCRP substrates and inhibitors suitable for human use were identified through a comprehensive analysis of available pharmacogenetics and DDI data. Two substrates were selected to probe intestinal and/or hepatic BCRP-mediated DDIs. Oral sulfasalazine (1000 mg, IR tablets) was deemed the best available probe for intestinal BCRP, oral rosuvastatin (20 mg) as the best available probe for both intestinal and hepatic BCRP, and intravenous rosuvastatin (4 mg) as the best available probe for hepatic BCRP. Oral curcumin (2000 mg) and oral lapatinib (sub-efficacious 250 mg dose) were deemed the best available BCRP inhibitors. Optimal study designs that would elicit the worst-case clinical BCRP DDI scenarios are proposed (Figure 1). The pharmacokinetic data obtained from the proposed 9-arm clinical study should provide the pharmaceutical industry and regulatory agencies insight into the extent of BCRP-mediated clinical DDIs, as well as validate clinical study designs that enable optimal assessment of intestinal and combined intestinal and hepatic BCRP-mediated DDIs.

Author Contributions

Participated in research design: Lee, O'Connor, Galetin, Cook, Ellens, Feng, Taub, Paine, Polli, Ware, and Zamek-Gliszczynski

Contributed new reagents: NA

Conducted experiments: NA

Performed data analysis: O'Connor, Ragueneau-Majlessi, Ritchie

Wrote or contributed to the writing of the manuscript: Lee, O'Connor, Ritchie, Galetin, Cook, Feng, Ellens, Ragueneau-Majlessi, Taub, Paine, Polli, Ware, and Zamek-Gliszczynski

References

European Medicines Agency (EMA). 2012. Guideline on the Investigation of Drug Interactions.

EMA website [online]:

http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/07/WC500129606.pdf.

FDA Guidance for Industry 2012, Drug Interaction Studies — Study Design, Data Analysis, Implications for Dosing, and Labeling Recommendations. FDA website [online]:

<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM292362.pdf>.

Pfizer, Inc. (2011) Azulfidine (sulfasalazine) prescribing information.

Abodunde OA, LevakaVeera RR, Desai R, Nweke N, and Berrou M (2013) Colchicine toxicity precipitated by interaction with sunitinib. *J Clin Pharm Ther* **38**:243-245.

Adkison KK, Vaidya SS, Lee DY, Koo SH, Li L, Mehta AA, Gross AS, Polli JW, Humphreys JE, Lou Y, and Lee EJ (2010) Oral sulfasalazine as a clinical BCRP probe substrate: pharmacokinetic effects of genetic variation (C421A) and pantoprazole coadministration. *J Pharm Sci* **99**:1046-1062.

Adkison KK, Vaidya SS, Lee DY, Koo SH, Li L, Mehta AA, Gross AS, Polli JW, Lou Y, and Lee EJ (2008) The ABCG2 C421A polymorphism does not affect oral nitrofurantoin pharmacokinetics in healthy Chinese male subjects. *Br J Clin Pharmacol* **66**:233-239.

Agarwal S, Mittapalli RK, Zellmer DM, Gallardo JL, Donelson R, Seiler C, Decker SA, Santacruz KS, Pokorny JL, Sarkaria JN, Elmquist WF, and Ohlfest JR (2012) Active efflux of Dasatinib from the brain limits efficacy against murine glioblastoma: broad

implications for the clinical use of molecularly targeted agents. *Mol Cancer Ther* **11**:2183-2192.

Ahmed-Belkacem A, Pozza A, Munoz-Martinez F, Bates SE, Castanys S, Gamarro F, Di Pietro A, and Perez-Victoria JM (2005) Flavonoid structure-activity studies identify 6-prenylchrysin and tectochrysin as potent and specific inhibitors of breast cancer resistance protein ABCG2. *Cancer Res* **65**:4852-4860.

Ai L, Zhu L, Yang L, Ge G, Cao Y, Liu Y, Fang Z, and Zhang Y (2014) Selectivity for inhibition of nilotinib on the catalytic activity of human UDP-glucuronosyltransferases. *Xenobiotica* **44**:320-325.

Allred AJ, Bowen CJ, Park JW, Peng B, Williams DD, Wire MB, and Lee E (2011) Eltrombopag increases plasma rosuvastatin exposure in healthy volunteers. *Br J Clin Pharmacol* **72**:321-329.

Ando T, Kusuhara H, Merino G, Alvarez AI, Schinkel AH, and Sugiyama Y (2007) Involvement of breast cancer resistance protein (ABCG2) in the biliary excretion mechanism of fluoroquinolones. *Drug Metab Dispos* **35**:1873-1879.

Appiah-Opong R, Commandeur JN, van Vugt-Lussenburg B, and Vermeulen NP (2007) Inhibition of human recombinant cytochrome P450s by curcumin and curcumin decomposition products. *Toxicology* **235**:83-91.

Beery E, Rajnai Z, Abonyi T, Makai I, Bansaghi S, Erdo F, Sziraki I, Heredi-Szabo K, Kis E, Jani M, Marki-Zay J, Toth GK, and Krajcsi P (2012) ABCG2 modulates chlorothiazide permeability--in vitro-characterization of its interactions. *Drug Metab Pharmacokinet* **27**:349-353.

- Bence AK, Anderson EB, Halepota MA, Doukas MA, DeSimone PA, Davis GA, Smith DA, Koch KM, Stead AG, Mangum S, Bowen CJ, Spector NL, Hsieh S, and Adams VR (2005) Phase I pharmacokinetic studies evaluating single and multiple doses of oral GW572016, a dual EGFR-ErbB2 inhibitor, in healthy subjects. *Invest New Drugs* **23**:39-49.
- Bhullar J, Natarajan K, Shukla S, Mathias TJ, Sadowska M, Ambudkar SV, and Baer MR (2013) The FLT3 inhibitor quizartinib inhibits ABCG2 at pharmacologically relevant concentrations, with implications for both chemosensitization and adverse drug interactions. *PLoS One* **8**:e71266.
- Bliesath H, Huber R, Steijnmans VW, Koch HJ, Kunz K, and Wurst W (1996a) Pantoprazole does not interact with nifedipine in man under steady-state conditions. *Int J Clin Pharmacol Ther* **34**:51-55.
- Bliesath H, Huber R, Steijnmans VW, Koch HJ, Kunz K, and Wurst W (1996b) Pantoprazole does not interact with nifedipine in man under steady-state conditions. *Int J Clin Pharmacol Ther* **34**:S81-85.
- Bliesath H, Huber R, Steijnmans VW, Koch HJ, Wurst W, and Mascher H (1996c) Lack of pharmacokinetic interaction between pantoprazole and diclofenac. *Int J Clin Pharmacol Ther* **34**:S76-80.
- Bliesath H, Huber R, Steijnmans VW, Koch HJ, Wurst W, and Mascher H (1996d) Lack of pharmacokinetic interaction between pantoprazole and diclofenac. *Int J Clin Pharmacol Ther* **34**:152-156.
- Bullock KE, Petros WP, Younis I, Uronis HE, Morse MA, Blobe GC, Zafar SY, Gockerman JP, Lager JJ, Truax R, Meadows KL, Howard LA, O'Neill MM, Broadwater G, Hurwitz HI,

- and Bendell JC (2011) A phase I study of bevacizumab (B) in combination with everolimus (E) and erlotinib (E) in advanced cancer (BEE). *Cancer Chemother Pharmacol* **67**:465-474.
- Busti AJ, Bain AM, Hall RG, 2nd, Bedimo RG, Leff RD, Meek C, and Mehvar R (2008) Effects of atazanavir/ritonavir or fosamprenavir/ritonavir on the pharmacokinetics of rosuvastatin. *J Cardiovasc Pharmacol* **51**:605-610.
- Castellino S, O'Mara M, Koch K, Borts DJ, Bowers GD, and MacLauchlin C (2012) Human metabolism of lapatinib, a dual kinase inhibitor: implications for hepatotoxicity. *Drug Metab Dispos* **40**:139-150.
- Chen ZS, Robey RW, Belinsky MG, Shchaveleva I, Ren XQ, Sugimoto Y, Ross DD, Bates SE, and Kruh GD (2003) Transport of methotrexate, methotrexate polyglutamates, and 17beta-estradiol 17-(beta-D-glucuronide) by ABCG2: effects of acquired mutations at R482 on methotrexate transport. *Cancer Res* **63**:4048-4054.
- Cheng AL, Hsu CH, Lin JK, Hsu MM, Ho YF, Shen TS, Ko JY, Lin JT, Lin BR, Ming-Shiang W, Yu HS, Jee SH, Chen GS, Chen TM, Chen CA, Lai MK, Pu YS, Pan MH, Wang YJ, Tsai CC, and Hsieh CY (2001) Phase I clinical trial of curcumin, a chemopreventive agent, in patients with high-risk or pre-malignant lesions. *Anticancer Res* **21**:2895-2900.
- Chien AJ, Munster PN, Melisko ME, Rugo HS, Park JW, Goga A, Auerback G, Khanafshar E, Ordovas K, Koch KM, and Moasser MM (2014) Phase I dose-escalation study of 5-day intermittent oral lapatinib therapy in patients with human epidermal growth factor receptor 2-overexpressing breast cancer. *J Clin Oncol* **32**:1472-1479.
- Choi MK and Song IS (2012) Characterization of efflux transport of the PDE5 inhibitors, vardenafil and sildenafil. *J Pharm Pharmacol* **64**:1074-1083.

- Chu X, Cai X, Cui D, Tang C, Ghosal A, Chan G, Green MD, Kuo Y, Liang Y, Maciolek CM, Palamanda J, Evers R, and Prueksaritanont T (2013a) In vitro assessment of drug-drug interaction potential of boceprevir associated with drug metabolizing enzymes and transporters. *Drug Metab Dispos* **41**:668-681.
- Chu X, Korzekwa K, Elsby R, Fenner K, Galetin A, Lai Y, Matsson P, Moss A, Nagar S, Rosania GR, Bai JP, Polli JW, Sugiyama Y, Brouwer KL, and International Transporter C (2013b) Intracellular drug concentrations and transporters: measurement, modeling, and implications for the liver. *Clin Pharmacol Ther* **94**:126-141.
- Crowe A and Wright C (2012) The impact of P-glycoprotein mediated efflux on absorption of 11 sedating and less-sedating antihistamines using Caco-2 monolayers. *Xenobiotica* **42**:538-549.
- Cusatis G, Gregorc V, Li J, Spreafico A, Ingersoll RG, Verweij J, Ludovini V, Villa E, Hidalgo M, Sparreboom A, and Baker SD (2006) Pharmacogenetics of ABCG2 and adverse reactions to gefitinib. *J Natl Cancer Inst* **98**:1739-1742.
- Custodio JM, Wang H, Hao J, Lepist EI, Ray AS, Andrews J, Ling KH, Cheng A, Kearney BP, and Ramanathan S (2014) Pharmacokinetics of cobicistat boosted-elvitegravir administered in combination with rosuvastatin. *J Clin Pharmacol* **54**:649-656.
- Dahan A and Amidon GL (2009) Small intestinal efflux mediated by MRP2 and BCRP shifts sulfasalazine intestinal permeability from high to low, enabling its colonic targeting. *Am J Physiol Gastrointest Liver Physiol* **297**:G371-377.
- de Jong FA, Marsh S, Mathijssen RH, King C, Verweij J, Sparreboom A, and McLeod HL (2004) ABCG2 pharmacogenetics: ethnic differences in allele frequency and assessment of influence on irinotecan disposition. *Clin Cancer Res* **10**:5889-5894.

- Dickens D, Owen A, Alfirevic A, and Pirmohamed M (2013) ABCB1 single nucleotide polymorphisms (1236C>T, 2677G>T, and 3435C>T) do not affect transport activity of human P-glycoprotein. *Pharmacogenet Genomics* **23**:314-323.
- Dong PP, Fang ZZ, Zhang YY, Ge GB, Mao YX, Zhu LL, Qu YQ, Li W, Wang LM, Liu CX, and Yang L (2011) Substrate-dependent modulation of the catalytic activity of CYP3A by erlotinib. *Acta Pharmacol Sin* **32**:399-407.
- Duursema L, Muller FO, Schall R, Middle MV, Hundt HK, Groenewoud G, Steinijs VW, and Bliesath H (1995) Lack of effect of pantoprazole on the pharmacodynamics and pharmacokinetics of warfarin. *Br J Clin Pharmacol* **39**:700-703.
- Egan A and Colman E (2011) Weighing the benefits of high-dose simvastatin against the risk of myopathy. *N Engl J Med* **365**:285-287.
- Elsby R, Hilgendorf C, and Fenner K (2012) Understanding the critical disposition pathways of statins to assess drug-drug interaction risk during drug development: it's not just about OATP1B1. *Clin Pharmacol Ther* **92**:584-598.
- Elsby R, Smith V, Fox L, Stresser D, Butters C, Sharma P, and Surry DD (2011) Validation of membrane vesicle-based breast cancer resistance protein and multidrug resistance protein 2 assays to assess drug transport and the potential for drug-drug interaction to support regulatory submissions. *Xenobiotica* **41**:764-783.
- Endres CJ, Hsiao P, Chung FS, and Unadkat JD (2006) The role of transporters in drug interactions. *Eur J Pharm Sci* **27**:501-517.
- Englund G, Lundquist P, Skogastierna C, Johansson J, Hoogstraate J, Afzelius L, Andersson TB, and Projean D (2014) Cytochrome p450 inhibitory properties of common efflux transporter inhibitors. *Drug Metab Dispos* **42**:441-447.

- Englund G, Rorsman F, Ronnblom A, Karlbom U, Lazorova L, Grasjo J, Kindmark A, and Artursson P (2006) Regional levels of drug transporters along the human intestinal tract: co-expression of ABC and SLC transporters and comparison with Caco-2 cells. *Eur J Pharm Sci* **29**:269-277.
- Enokizono J, Kusuhara H, and Sugiyama Y (2007a) Involvement of breast cancer resistance protein (BCRP/ABCG2) in the biliary excretion and intestinal efflux of troglitazone sulfate, the major metabolite of troglitazone with a cholestatic effect. *Drug Metab Dispos* **35**:209-214.
- Enokizono J, Kusuhara H, and Sugiyama Y (2007b) Regional expression and activity of breast cancer resistance protein (Bcrp/Abcg2) in mouse intestine: overlapping distribution with sulfotransferases. *Drug Metab Dispos* **35**:922-928.
- Farrell TL, Poquet L, Dew TP, Barber S, and Williamson G (2012) Predicting phenolic acid absorption in Caco-2 cells: a theoretical permeability model and mechanistic study. *Drug Metab Dispos* **40**:397-406.
- Ferrante E, Grasso S, Corbo GM, and Ciappi G (1991) [The measurement of bronchial hyperreactivity for military service fitness]. *Recenti Prog Med* **82**:520-522.
- Fujita Y, Noguchi K, Suzuki T, Katayama K, and Sugimoto Y (2013) Biochemical interaction of anti-HCV telaprevir with the ABC transporters P-glycoprotein and breast cancer resistance protein. *BMC Res Notes* **6**:445.
- Furukawa T, Wakabayashi K, Tamura A, Nakagawa H, Morishima Y, Osawa Y, and Ishikawa T (2009) Major SNP (Q141K) variant of human ABC transporter ABCG2 undergoes lysosomal and proteasomal degradations. *Pharm Res* **26**:469-479.

- Garcea G, Berry DP, Jones DJ, Singh R, Dennison AR, Farmer PB, Sharma RA, Steward WP, and Gescher AJ (2005) Consumption of the putative chemopreventive agent curcumin by cancer patients: assessment of curcumin levels in the colorectum and their pharmacodynamic consequences. *Cancer Epidemiol Biomarkers Prev* **14**:120-125.
- Giacomini KM, Balimane PV, Cho SK, Eadon M, Edeki T, Hillgren KM, Huang SM, Sugiyama Y, Weitz D, Wen Y, Xia CQ, Yee SW, Zimdahl H, Niemi M, and International Transporter C (2013) International Transporter Consortium commentary on clinically important transporter polymorphisms. *Clin Pharmacol Ther* **94**:23-26.
- Giacomini KM, Huang SM, Tweedie DJ, Benet LZ, Brouwer KL, Chu X, Dahlin A, Evers R, Fischer V, Hillgren KM, Hoffmaster KA, Ishikawa T, Keppler D, Kim RB, Lee CA, Niemi M, Polli JW, Sugiyama Y, Swaan PW, Ware JA, Wright SH, Yee SW, Zamek-Gliszczynski MJ, and Zhang L (2010) Membrane transporters in drug development. *Nat Rev Drug Discov* **9**:215-236.
- Gong IY, Mansell SE, and Kim RB (2013) Absence of both MDR1 (ABCB1) and breast cancer resistance protein (ABCG2) transporters significantly alters rivaroxaban disposition and central nervous system entry. *Basic Clin Pharmacol Toxicol* **112**:164-170.
- Goodman LS, Gilman A, Brunton LL, Lazo JS, and Parker KL (2005) *Goodman & Gilman's the pharmacological basis of therapeutics*. . McGraw-Hill, New York, N.Y.
- Goodman LS, Hardman JG, Limbird LE, and Gilman AG (2001) *Goodman & Gilman's the pharmacological basis of therapeutics*. . McGraw-Hill, New York, N.Y.
- Gugler R, Hartmann M, Rudi J, Brod I, Huber R, Steinijs VW, Bliesath H, Wurst W, and Klotz U (1996) Lack of pharmacokinetic interaction of pantoprazole with diazepam in man. *Br J Clin Pharmacol* **42**:249-252.

- Haagsma CJ, Russel FG, Vree TB, Van Riel PL, and Van de Putte LB (1996) Combination of methotrexate and sulphasalazine in patients with rheumatoid arthritis: pharmacokinetic analysis and relationship to clinical response. *Br J Clin Pharmacol* **42**:195-200.
- Han JY, Lim HS, Yoo YK, Shin ES, Park YH, Lee SY, Lee JE, Lee DH, Kim HT, and Lee JS (2007) Associations of ABCB1, ABCC2, and ABCG2 polymorphisms with irinotecan-pharmacokinetics and clinical outcome in patients with advanced non-small cell lung cancer. *Cancer* **110**:138-147.
- Han YH, Busler D, Hong Y, Tian Y, Chen C, and Rodrigues AD (2010) Transporter studies with the 3-O-sulfate conjugate of 17alpha-ethinylestradiol: assessment of human liver drug transporters. *Drug Metab Dispos* **38**:1072-1082.
- Hartmann M, Huber R, Bliesath H, Steinijans VW, Koch HJ, Wurst W, and Kunz K (1995) Lack of interaction between pantoprazole and digoxin at therapeutic doses in man. *Int J Clin Pharmacol Ther* **33**:481-485.
- Hartmann M, Huber R, Bliesath H, Steinijans VW, Koch HJ, Wurst W, and Kunz K (1996) Lack of interaction between pantoprazole and digoxin at therapeutic doses in man. *Int J Clin Pharmacol Ther* **34**:S67-71.
- He X, Mo L, Li ZY, Tan ZR, Chen Y, and Ouyang DS (2012) Effects of curcumin on the pharmacokinetics of talinolol in human with ABCB1 polymorphism. *Xenobiotica* **42**:1248-1254.
- Hemauer SJ, Patrikeeva SL, Wang X, Abdelrahman DR, Hankins GD, Ahmed MS, and Nanovskaya TN (2010) Role of transporter-mediated efflux in the placental biodisposition of bupropion and its metabolite, OH-bupropion. *Biochem Pharmacol* **80**:1080-1086.

- Hirano M, Maeda K, Matsushima S, Nozaki Y, Kusuhara H, and Sugiyama Y (2005)
Involvement of BCRP (ABCG2) in the biliary excretion of pitavastatin. *Mol Pharmacol* **68**:800-807.
- Ho RH, Tirona RG, Leake BF, Glaeser H, Lee W, Lemke CJ, Wang Y, and Kim RB (2006)
Drug and bile acid transporters in rosuvastatin hepatic uptake: function, expression, and pharmacogenetics. *Gastroenterology* **130**:1793-1806.
- Hofman J, Kucera R, Cihalova D, Klimes J, Ceckova M, and Staud F (2013) Olomoucine II, but not purvalanol A, is transported by breast cancer resistance protein (ABCG2) and P-glycoprotein (ABCB1). *PLoS One* **8**:e75520.
- Huber R, Bliesath H, Hartmann M, Steinijans VW, Koch H, Mascher H, and Wurst W (1998)
Pantoprazole does not interact with the pharmacokinetics of carbamazepine. *Int J Clin Pharmacol Ther* **36**:521-524.
- Hyafil F, Vergely C, Du Vignaud P, and Grand-Perret T (1993) In vitro and in vivo reversal of multidrug resistance by GF120918, an acridonecarboxamide derivative. *Cancer Res* **53**:4595-4602.
- Ieiri I, Suwannakul S, Maeda K, Uchimaru H, Hashimoto K, Kimura M, Fujino H, Hirano M, Kusuhara H, Irie S, Higuchi S, and Sugiyama Y (2007) SLCO1B1 (OATP1B1, an uptake transporter) and ABCG2 (BCRP, an efflux transporter) variant alleles and pharmacokinetics of pitavastatin in healthy volunteers. *Clin Pharmacol Ther* **82**:541-547.
- Imai Y, Asada S, Tsukahara S, Ishikawa E, Tsuruo T, and Sugimoto Y (2003) Breast cancer resistance protein exports sulfated estrogens but not free estrogens. *Mol Pharmacol* **64**:610-618.

- Imai Y, Tsukahara S, Asada S, and Sugimoto Y (2004) Phytoestrogens/flavonoids reverse breast cancer resistance protein/ABCG2-mediated multidrug resistance. *Cancer Res* **64**:4346-4352.
- Itagaki F, Homma M, Yuzawa K, Nishimura M, Naito S, Ueda N, Ohkohchi N, and Kohda Y (2004) Effect of lansoprazole and rabeprazole on tacrolimus pharmacokinetics in healthy volunteers with CYP2C19 mutations. *J Pharm Pharmacol* **56**:1055-1059.
- Jada SR, Lim R, Wong CI, Shu X, Lee SC, Zhou Q, Goh BC, and Chowbay B (2007) Role of UGT1A1*6, UGT1A1*28 and ABCG2 c.421C>A polymorphisms in irinotecan-induced neutropenia in Asian cancer patients. *Cancer Sci* **98**:1461-1467.
- Jamei M, Bajot F, Neuhoff S, Barter Z, Yang J, Rostami-Hodjegan A, and Rowland-Yeo K (2014) A mechanistic framework for in vitro-in vivo extrapolation of liver membrane transporters: prediction of drug-drug interaction between rosuvastatin and cyclosporine. *Clin Pharmacokinet* **53**:73-87.
- Jani M, Makai I, Kis E, Szabo P, Nagy T, Krajcsi P, and Lespine A (2011) Ivermectin interacts with human ABCG2. *J Pharm Sci* **100**:94-97.
- Jani M, Szabo P, Kis E, Molnar E, Glavinas H, and Krajcsi P (2009) Kinetic characterization of sulfasalazine transport by human ATP-binding cassette G2. *Biol Pharm Bull* **32**:497-499.
- Jin HE, Song B, Kim SB, Shim WS, Kim DD, Chong S, Chung SJ, and Shim CK (2013) Transport of gemifloxacin, a 4th generation quinolone antibiotic, in the Caco-2 and engineered MDCKII cells, and potential involvement of efflux transporters in the intestinal absorption of the drug. *Xenobiotica* **43**:355-367.
- Jones HM, Barton HA, Lai Y, Bi YA, Kimoto E, Kempshall S, Tate SC, El-Kattan A, Houston JB, Galetin A, and Fenner KS (2012) Mechanistic pharmacokinetic modeling for the

- prediction of transporter-mediated disposition in humans from sandwich culture human hepatocyte data. *Drug Metab Dispos* **40**:1007-1017.
- Jonker JW, Smit JW, Brinkhuis RF, Maliepaard M, Beijnen JH, Schellens JH, and Schinkel AH (2000) Role of breast cancer resistance protein in the bioavailability and fetal penetration of topotecan. *J Natl Cancer Inst* **92**:1651-1656.
- Juan H, Terhaag B, Cong Z, Bi-Kui Z, Rong-Hua Z, Feng W, Fen-Li S, Juan S, Jing T, and Wen-Xing P (2007) Unexpected effect of concomitantly administered curcumin on the pharmacokinetics of talinolol in healthy Chinese volunteers. *Eur J Clin Pharmacol* **63**:663-668.
- Kalvass JC, Polli JW, Bourdet DL, Feng B, Huang SM, Liu X, Smith QR, Zhang LK, Zamek-Gliszczynski MJ, and International Transporter C (2013) Why clinical modulation of efflux transport at the human blood-brain barrier is unlikely: the ITC evidence-based position. *Clin Pharmacol Ther* **94**:80-94.
- Karibe T, Hagihara-Nakagomi R, Abe K, Imaoka T, Mikkaichi T, Yasuda S, Hirouchi M, Watanabe N, Okudaira N, and Izumi T (2014) Evaluation of the Usefulness of Breast Cancer Resistance Protein (BCRP) Knockout Mice and BCRP Inhibitor-Treated Monkeys to Estimate the Clinical Impact of BCRP Modulation on the Pharmacokinetics of BCRP Substrates. *Pharm Res*.
- Karlgren M, Ahlin G, Bergstrom CA, Svensson R, Palm J, and Artursson P (2012a) In vitro and in silico strategies to identify OATP1B1 inhibitors and predict clinical drug-drug interactions. *Pharm Res* **29**:411-426.
- Karlgren M, Vildhede A, Norinder U, Wisniewski JR, Kimoto E, Lai Y, Haglund U, and Artursson P (2012b) Classification of inhibitors of hepatic organic anion transporting

- polypeptides (OATPs): influence of protein expression on drug-drug interactions. *J Med Chem* **55**:4740-4763.
- Kawahara H, Noguchi K, Katayama K, Mitsuhashi J, and Sugimoto Y (2010) Pharmacological interaction with sunitinib is abolished by a germ-line mutation (1291T>C) of BCRP/ABCG2 gene. *Cancer Sci* **101**:1493-1500.
- Kenny JR, Mukadam S, Zhang C, Tay S, Collins C, Galetin A, and Khojasteh SC (2012) Drug-drug interaction potential of marketed oncology drugs: in vitro assessment of time-dependent cytochrome P450 inhibition, reactive metabolite formation and drug-drug interaction prediction. *Pharm Res* **29**:1960-1976.
- Keskitalo JE, Pasanen MK, Neuvonen PJ, and Niemi M (2009a) Different effects of the ABCG2 c.421C>A SNP on the pharmacokinetics of fluvastatin, pravastatin and simvastatin. *Pharmacogenomics* **10**:1617-1624.
- Keskitalo JE, Zolk O, Fromm MF, Kurkinen KJ, Neuvonen PJ, and Niemi M (2009b) ABCG2 polymorphism markedly affects the pharmacokinetics of atorvastatin and rosuvastatin. *Clin Pharmacol Ther* **86**:197-203.
- Kim HS, Sunwoo YE, Ryu JY, Kang HJ, Jung HE, Song IS, Kim EY, Shim JC, Shon JH, and Shin JG (2007) The effect of ABCG2 V12M, Q141K and Q126X, known functional variants in vitro, on the disposition of lamivudine. *Br J Clin Pharmacol* **64**:645-654.
- Kim MJ, Lee JW, Oh KS, Choi CS, Kim KH, Han WS, Yoon CN, Chung ES, Kim DH, and Shin JG (2013) The tyrosine kinase inhibitor nilotinib selectively inhibits CYP2C8 activities in human liver microsomes. *Drug Metab Pharmacokinet* **28**:462-467.

- Kiser JJ, Gerber JG, Predhomme JA, Wolfe P, Flynn DM, and Hoody DW (2008) Drug/Drug interaction between lopinavir/ritonavir and rosuvastatin in healthy volunteers. *J Acquir Immune Defic Syndr* **47**:570-578.
- Kitamura S, Maeda K, Wang Y, and Sugiyama Y (2008) Involvement of multiple transporters in the hepatobiliary transport of rosuvastatin. *Drug Metab Dispos* **36**:2014-2023.
- Koch HJ, Hartmann M, Bliesath H, Huber R, Steinijans VW, Mascher H, and Wurst W (1996) Pantoprazole has no influence on steady state pharmacokinetics and pharmacodynamics of metoprolol in healthy volunteers. *Int J Clin Pharmacol Ther* **34**:420-423.
- Kondo C, Suzuki H, Itoda M, Ozawa S, Sawada J, Kobayashi D, Ieiri I, Mine K, Ohtsubo K, and Sugiyama Y (2004) Functional analysis of SNPs variants of BCRP/ABCG2. *Pharm Res* **21**:1895-1903.
- Kruijtzter CM, Beijnen JH, Rosing H, ten Bokkel Huinink WW, Schot M, Jewell RC, Paul EM, and Schellens JH (2002) Increased oral bioavailability of topotecan in combination with the breast cancer resistance protein and P-glycoprotein inhibitor GF120918. *J Clin Oncol* **20**:2943-2950.
- Kuppens IE, Witteveen EO, Jewell RC, Radema SA, Paul EM, Mangum SG, Beijnen JH, Voest EE, and Schellens JH (2007) A phase I, randomized, open-label, parallel-cohort, dose-finding study of elacridar (GF120918) and oral topotecan in cancer patients. *Clin Cancer Res* **13**:3276-3285.
- Kusuhara H, Furuie H, Inano A, Sunagawa A, Yamada S, Wu C, Fukizawa S, Morimoto N, Ieiri I, Morishita M, Sumita K, Mayahara H, Fujita T, Maeda K, and Sugiyama Y (2012) Pharmacokinetic interaction study of sulphasalazine in healthy subjects and the impact of curcumin as an in vivo inhibitor of BCRP. *Br J Pharmacol* **166**:1793-1803.

- Lagas JS, van Waterschoot RA, Sparidans RW, Wagenaar E, Beijnen JH, and Schinkel AH (2010) Breast cancer resistance protein and P-glycoprotein limit sorafenib brain accumulation. *Mol Cancer Ther* **9**:319-326.
- Lainey E, Sebert M, Thepot S, Scoazec M, Bouteloup C, Leroy C, De Botton S, Galluzzi L, Fenaux P, and Kroemer G (2012) Erlotinib antagonizes ABC transporters in acute myeloid leukemia. *Cell Cycle* **11**:4079-4092.
- Leggas M, Panetta JC, Zhuang Y, Schuetz JD, Johnston B, Bai F, Sorrentino B, Zhou S, Houghton PJ, and Stewart CF (2006) Gefitinib modulates the function of multiple ATP-binding cassette transporters in vivo. *Cancer Res* **66**:4802-4807.
- Li H, Jin HE, Kim W, Han YH, Kim DD, Chung SJ, and Shim CK (2008) Involvement of P-glycoprotein, multidrug resistance protein 2 and breast cancer resistance protein in the transport of belotecan and topotecan in Caco-2 and MDCKII cells. *Pharm Res* **25**:2601-2612.
- Li J, Volpe DA, Wang Y, Zhang W, Bode C, Owen A, and Hidalgo IJ (2011a) Use of transporter knockdown Caco-2 cells to investigate the in vitro efflux of statin drugs. *Drug Metab Dispos* **39**:1196-1202.
- Li J, Wang Y, Zhang W, Huang Y, Hein K, and Hidalgo IJ (2012) The role of a basolateral transporter in rosuvastatin transport and its interplay with apical breast cancer resistance protein in polarized cell monolayer systems. *Drug Metab Dispos* **40**:2102-2108.
- Li L, Sham YY, Bikadi Z, and Elmquist WF (2011b) pH-Dependent transport of pemetrexed by breast cancer resistance protein. *Drug Metab Dispos* **39**:1478-1485.
- Li R, Bi YA, Lai Y, Sugano K, Steyn SJ, Trapa PE, and Di L (2014) Permeability comparison between hepatocyte and low efflux MDCKII cell monolayer. *Aaps J* **16**:802-809.

- Li X, Kamenecka TM, and Cameron MD (2010) Cytochrome P450-mediated bioactivation of the epidermal growth factor receptor inhibitor erlotinib to a reactive electrophile. *Drug Metab Dispos* **38**:1238-1245.
- Li XQ, Andersson TB, Ahlstrom M, and Weidolf L (2004) Comparison of inhibitory effects of the proton pump-inhibiting drugs omeprazole, esomeprazole, lansoprazole, pantoprazole, and rabeprazole on human cytochrome P450 activities. *Drug Metab Dispos* **32**:821-827.
- Lin X, Skolnik S, Chen X, and Wang J (2011) Attenuation of intestinal absorption by major efflux transporters: quantitative tools and strategies using a Caco-2 model. *Drug Metab Dispos* **39**:265-274.
- Liu Y, Ramirez J, House L, and Ratain MJ (2010) Comparison of the drug-drug interactions potential of erlotinib and gefitinib via inhibition of UDP-glucuronosyltransferases. *Drug Metab Dispos* **38**:32-39.
- Lorf T, Ramadori G, Ringe B, and Schworer H (2000) The effect of pantoprazole on tacrolimus and cyclosporin A blood concentration in transplant recipients. *Eur J Clin Pharmacol* **56**:439-440.
- MacFarlane RJ and Gelmon KA (2011) Lapatinib for breast cancer: a review of the current literature. *Expert Opin Drug Saf* **10**:109-121.
- MacLean C, Moenning U, Reichel A, and Fricker G (2008) Closing the gaps: a full scan of the intestinal expression of p-glycoprotein, breast cancer resistance protein, and multidrug resistance-associated protein 2 in male and female rats. *Drug Metab Dispos* **36**:1249-1254.

- Macpherson IR, Poondru S, Simon GR, Gedrich R, Brock K, Hopkins CA, Stewart K, Stephens A, and Evans TR (2013) A phase 1 study of OSI-930 in combination with erlotinib in patients with advanced solid tumours. *Eur J Cancer* **49**:782-789.
- Malingre MM, Beijnen JH, Rosing H, Koopman FJ, Jewell RC, Paul EM, Ten Bokkel Huinink WW, and Schellens JH (2001) Co-administration of GF120918 significantly increases the systemic exposure to oral paclitaxel in cancer patients. *Br J Cancer* **84**:42-47.
- Mao Q and Unadkat JD (2005) Role of the breast cancer resistance protein (ABCG2) in drug transport. *Aaps J* **7**:E118-133.
- Marchetti S, Pluim D, van Eijndhoven M, van Tellingen O, Mazzanti R, Beijnen JH, and Schellens JH (2013) Effect of the drug transporters ABCG2, Abcg2, ABCB1 and ABCC2 on the disposition, brain accumulation and myelotoxicity of the aurora kinase B inhibitor barasertib and its more active form barasertib-hydroxy-QPA. *Invest New Drugs* **31**:1125-1135.
- Martin PD, Warwick MJ, Dane AL, Brindley C, and Short T (2003) Absolute oral bioavailability of rosuvastatin in healthy white adult male volunteers. *Clin Ther* **25**:2553-2563.
- Matsuo H, Takada T, Ichida K, Nakamura T, Nakayama A, Ikebuchi Y, Ito K, Kusanagi Y, Chiba T, Tadokoro S, Takada Y, Oikawa Y, Inoue H, Suzuki K, Okada R, Nishiyama J, Domoto H, Watanabe S, Fujita M, Morimoto Y, Naito M, Nishio K, Hishida A, Wakai K, Asai Y, Niwa K, Kamakura K, Nonoyama S, Sakurai Y, Hosoya T, Kanai Y, Suzuki H, Hamajima N, and Shinomiya N (2009) Common defects of ABCG2, a high-capacity urate exporter, cause gout: a function-based genetic analysis in a Japanese population. *Sci Transl Med* **1**:5ra11.

- Menochet K, Kenworthy KE, Houston JB, and Galetin A (2012) Use of Mechanistic Modelling to Assess Inter-Individual Variability and Inter-species Differences in Active Uptake in Human and Rat Hepatocytes. *Drug Metab Dispos* **40**:1744-1756.
- Merino G, Alvarez AI, Pulido MM, Molina AJ, Schinkel AH, and Prieto JG (2006) Breast cancer resistance protein (BCRP/ABCG2) transports fluoroquinolone antibiotics and affects their oral availability, pharmacokinetics, and milk secretion. *Drug Metab Dispos* **34**:690-695.
- Merino G, Perez M, Real R, Egado E, Prieto JG, and Alvarez AI (2010) In vivo inhibition of BCRP/ABCG2 mediated transport of nitrofurantoin by the isoflavones genistein and daidzein: a comparative study in Bcrp1 (-/-) mice. *Pharm Res* **27**:2098-2105.
- Messersmith WA, Laheru DA, Senzer NN, Donehower RC, Grouleff P, Rogers T, Kelley SK, Ramies DA, Lum BL, and Hidalgo M (2004) Phase I trial of irinotecan, infusional 5-fluorouracil, and leucovorin (FOLFIRI) with erlotinib (OSI-774): early termination due to increased toxicities. *Clin Cancer Res* **10**:6522-6527.
- Middle MV, Muller FO, Schall R, Groenewoud G, Hundt HK, Huber R, Bliesath H, and Steinijans VW (1995) No influence of pantoprazole on the pharmacokinetics of phenytoin. *Int J Clin Pharmacol Ther* **33**:304-307.
- Middle MV, Muller FO, Schall R, Groenewoud G, Hundt HK, Huber R, Bliesath H, and Steinijans VW (1996) No influence of pantoprazole on the pharmacokinetics of phenytoin. *Int J Clin Pharmacol Ther* **34**:S72-75.
- Midgley RS, Kerr DJ, Flaherty KT, Stevenson JP, Pratap SE, Koch KM, Smith DA, Versola M, Fleming RA, Ward C, O'Dwyer PJ, and Middleton MR (2007) A phase I and

- pharmacokinetic study of lapatinib in combination with infusional 5-fluorouracil, leucovorin and irinotecan. *Ann Oncol* **18**:2025-2029.
- Milton DT, Riely GJ, Azzoli CG, Gomez JE, Heelan RT, Kris MG, Krug LM, Pao W, Pizzo B, Rizvi NA, and Miller VA (2007) Phase 1 trial of everolimus and gefitinib in patients with advanced nonsmall-cell lung cancer. *Cancer* **110**:599-605.
- Minematsu T and Giacomini KM (2011) Interactions of tyrosine kinase inhibitors with organic cation transporters and multidrug and toxic compound extrusion proteins. *Mol Cancer Ther* **10**:531-539.
- Mizuno N, Takahashi T, Kusuhara H, Schuetz JD, Niwa T, and Sugiyama Y (2007) Evaluation of the role of breast cancer resistance protein (BCRP/ABCG2) and multidrug resistance-associated protein 4 (MRP4/ABCC4) in the urinary excretion of sulfate and glucuronide metabolites of edaravone (MCI-186; 3-methyl-1-phenyl-2-pyrazolin-5-one). *Drug Metab Dispos* **35**:2045-2052.
- Molina JR, Kaufmann SH, Reid JM, Rubin SD, Galvez-Peralta M, Friedman R, Flatten KS, Koch KM, Gilmer TM, Mullin RJ, Jewell RC, Felten SJ, Mandrekar S, Adjei AA, and Erlichman C (2008) Evaluation of lapatinib and topotecan combination therapy: tissue culture, murine xenograft, and phase I clinical trial data. *Clin Cancer Res* **14**:7900-7908.
- Moody DE, Liu F, and Fang WB (2013) In vitro inhibition of methadone and oxycodone cytochrome P450-dependent metabolism: reversible inhibition by H2-receptor agonists and proton-pump inhibitors. *J Anal Toxicol* **37**:476-485.
- Nagasaka Y, Oda K, Iwatsubo T, Kawamura A, and Usui T (2012) Effects of aripiprazole and its active metabolite dehydroaripiprazole on the activities of drug efflux transporters

- expressed both in the intestine and at the blood-brain barrier. *Biopharm Drug Dispos* **33**:304-315.
- Nakatomi K, Yoshikawa M, Oka M, Ikegami Y, Hayasaka S, Sano K, Shiozawa K, Kawabata S, Soda H, Ishikawa T, Tanabe S, and Kohno S (2001) Transport of 7-ethyl-10-hydroxycamptothecin (SN-38) by breast cancer resistance protein ABCG2 in human lung cancer cells. *Biochem Biophys Res Commun* **288**:827-832.
- Ni Z, Bikadi Z, Rosenberg MF, and Mao Q (2010) Structure and function of the human breast cancer resistance protein (BCRP/ABCG2). *Curr Drug Metab* **11**:603-617.
- Noguchi K, Kawahara H, Kaji A, Katayama K, Mitsuhashi J, and Sugimoto Y (2009) Substrate-dependent bidirectional modulation of P-glycoprotein-mediated drug resistance by erlotinib. *Cancer Sci* **100**:1701-1707.
- Oberoi RK, Mittapalli RK, and Elmquist WF (2013) Pharmacokinetic assessment of efflux transport in sunitinib distribution to the brain. *J Pharmacol Exp Ther* **347**:755-764.
- Ohura K, Nozawa T, Murakami K, and Imai T (2011) Evaluation of transport mechanism of prodrugs and parent drugs formed by intracellular metabolism in Caco-2 cells with modified carboxylesterase activity: temocapril as a model case. *J Pharm Sci* **100**:3985-3994.
- Oostendorp RL, van de Steeg E, van der Kruijssen CM, Beijnen JH, Kenworthy KE, Schinkel AH, and Schellens JH (2009) Organic anion-transporting polypeptide 1B1 mediates transport of Gimatecan and BNP1350 and can be inhibited by several classic ATP-binding cassette (ABC) B1 and/or ABCG2 inhibitors. *Drug Metab Dispos* **37**:917-923.
- Pan Y, Chothe PP, and Swaan PW (2013) Identification of novel breast cancer resistance protein (BCRP) inhibitors by virtual screening. *Mol Pharm* **10**:1236-1248.

- Pang KS (2003) Modeling of intestinal drug absorption: roles of transporters and metabolic enzymes (for the Gillette Review Series). *Drug Metab Dispos* **31**:1507-1519.
- Pasanen MK, Fredrikson H, Neuvonen PJ, and Niemi M (2007) Different effects of SLCO1B1 polymorphism on the pharmacokinetics of atorvastatin and rosuvastatin. *Clin Pharmacol Ther* **82**:726-733.
- Pauli-Magnus C, Rekersbrink S, Klotz U, and Fromm MF (2001) Interaction of omeprazole, lansoprazole and pantoprazole with P-glycoprotein. *Naunyn Schmiedebergs Arch Pharmacol* **364**:551-557.
- Pavek P, Merino G, Wagenaar E, Bolscher E, Novotna M, Jonker JW, and Schinkel AH (2005) Human breast cancer resistance protein: interactions with steroid drugs, hormones, the dietary carcinogen 2-amino-1-methyl-6-phenylimidazo(4,5-b)pyridine, and transport of cimetidine. *J Pharmacol Exp Ther* **312**:144-152.
- Pfeifer ND, Bridges AS, Ferslew BC, Hardwick RN, and Brouwer KL (2013a) Hepatic basolateral efflux contributes significantly to rosuvastatin disposition II: characterization of hepatic elimination by basolateral, biliary, and metabolic clearance pathways in rat isolated perfused liver. *J Pharmacol Exp Ther* **347**:737-745.
- Pfeifer ND, Yang K, and Brouwer KL (2013b) Hepatic basolateral efflux contributes significantly to rosuvastatin disposition I: characterization of basolateral versus biliary clearance using a novel protocol in sandwich-cultured hepatocytes. *J Pharmacol Exp Ther* **347**:727-736.
- Pham PA, la Porte CJ, Lee LS, van Heeswijk R, Sabo JP, Elgadi MM, Piliero PJ, Barditch-Crovo P, Fuchs E, Flexner C, and Cameron DW (2009) Differential effects of tipranavir plus

ritonavir on atorvastatin or rosuvastatin pharmacokinetics in healthy volunteers.

Antimicrob Agents Chemother **53**:4385-4392.

Planting AS, Sonneveld P, van der Gaast A, Sparreboom A, van der Burg ME, Luyten GP, de Leeuw K, de Boer-Dennert M, Wissel PS, Jewell RC, Paul EM, Purvis NB, Jr., and Verweij J (2005) A phase I and pharmacologic study of the MDR converter GF120918 in combination with doxorubicin in patients with advanced solid tumors. *Cancer Chemother Pharmacol* **55**:91-99.

Poirier A, Portmann R, Cascais AC, Bader U, Walter I, Ullah M, and Funk C (2014) The need for human breast cancer resistance protein substrate and inhibition evaluation in drug discovery and development: why, when, and how? *Drug Metab Dispos* **42**:1466-1477.

Pollex EK, Anger G, Hutson J, Koren G, and Piquette-Miller M (2010) Breast cancer resistance protein (BCRP)-mediated glyburide transport: effect of the C421A/Q141K BCRP single-nucleotide polymorphism. *Drug Metab Dispos* **38**:740-744.

Polli JW, Humphreys JE, Harmon KA, Castellino S, O'Mara MJ, Olson KL, John-Williams LS, Koch KM, and Serabjit-Singh CJ (2008) The role of efflux and uptake transporters in [N-{3-chloro-4-[(3-fluorobenzyl)oxy]phenyl}-6-[5-([2-(methylsulfonyl)ethyl]amino)methyl]-2-furyl]-4-quinazolinamine (GW572016, lapatinib) disposition and drug interactions. *Drug Metab Dispos* **36**:695-701.

Prueksaritanont T, Chu X, Evers R, Klopfer SO, Caro L, Kothare PA, Dempsey C, Rasmussen S, Houle R, Chan G, Cai X, Valesky R, Fraser IP, and Stoch SA (2014) Pitavastatin is a more sensitive and selective organic anion-transporting polypeptide 1B clinical probe than rosuvastatin. *Br J Clin Pharmacol* **78**:587-598.

- Ranchon F, Vantard N, Gouraud A, Schwiertz V, Franchon E, Pham BN, Vial T, You B, Bouafia F, Salles G, and Rioufol C (2011) Suspicion of drug-drug interaction between high-dose methotrexate and proton pump inhibitors: a case report - should the practice be changed? *Chemotherapy* **57**:225-229.
- Reese MJ, Savina PM, Generaux GT, Tracey H, Humphreys JE, Kanaoka E, Webster LO, Harmon KA, Clarke JD, and Polli JW (2013) In vitro investigations into the roles of drug transporters and metabolizing enzymes in the disposition and drug interactions of dolutegravir, a HIV integrase inhibitor. *Drug Metab Dispos* **41**:353-361.
- Reginato AM, Mount DB, Yang I, and Choi HK (2012) The genetics of hyperuricaemia and gout. *Nat Rev Rheumatol* **8**:610-621.
- Remes-Troche JM, Sobrino-Cossio S, Soto-Perez JC, Teramoto-Matsubara O, Morales-Arambula M, Orozco-Gamiz A, Tamayo de la Cuesta JL, and Mateos G (2014) Efficacy, safety, and tolerability of pantoprazole magnesium in the treatment of reflux symptoms in patients with gastroesophageal reflux disease (GERD): a prospective, multicenter, post-marketing observational study. *Clin Drug Investig* **34**:83-93.
- Reyner EL, Sevidal S, West MA, Clouser-Roche A, Freiwald S, Fenner K, Ullah M, Lee CA, and Smith BJ (2013) In vitro characterization of axitinib interactions with human efflux and hepatic uptake transporters: implications for disposition and drug interactions. *Drug Metab Dispos* **41**:1575-1583.
- Robert F, Sandler A, Schiller JH, Liu G, Harper K, Verkh L, Huang X, Ilagan J, Tye L, Chao R, and Traynor AM (2010) Sunitinib in combination with docetaxel in patients with advanced solid tumors: a phase I dose-escalation study. *Cancer Chemother Pharmacol* **66**:669-680.

- Robey RW, Ierano C, Zhan Z, and Bates SE (2011) The challenge of exploiting ABCG2 in the clinic. *Curr Pharm Biotechnol* **12**:595-608.
- Saison C, Helias V, Ballif BA, Peyrard T, Puy H, Miyazaki T, Perrot S, Vayssier-Taussat M, Waldner M, Le Pennec PY, Cartron JP, and Arnaud L (2012) Null alleles of ABCG2 encoding the breast cancer resistance protein define the new blood group system Junior. *Nat Genet* **44**:174-177.
- Saito H, Hirano H, Nakagawa H, Fukami T, Oosumi K, Murakami K, Kimura H, Kouchi T, Konomi M, Tao E, Tsujikawa N, Tarui S, Nagakura M, Osumi M, and Ishikawa T (2006) A new strategy of high-speed screening and quantitative structure-activity relationship analysis to evaluate human ATP-binding cassette transporter ABCG2-drug interactions. *J Pharmacol Exp Ther* **317**:1114-1124.
- Sharma RA, Gescher AJ, and Steward WP (2005) Curcumin: the story so far. *Eur J Cancer* **41**:1955-1968.
- Shitara Y, Maeda K, Ikejiri K, Yoshida K, Horie T, and Sugiyama Y (2013) Clinical significance of organic anion transporting polypeptides (OATPs) in drug disposition: their roles in hepatic clearance and intestinal absorption. *Biopharm Drug Dispos* **34**:45-78.
- Shukla S, Kouanda A, Silverton L, Talele TT, and Ambudkar SV (2014) Pharmacophore modeling of nilotinib as an inhibitor of ATP-binding cassette drug transporters and BCR-ABL kinase using a three-dimensional quantitative structure-activity relationship approach. *Mol Pharm* **11**:2313-2322.
- Shukla S, Zaher H, Hartz A, Bauer B, Ware JA, and Ambudkar SV (2009) Curcumin inhibits the activity of ABCG2/BCRP1, a multidrug resistance-linked ABC drug transporter in mice. *Pharm Res* **26**:480-487.

- Si M, Zhao J, Li X, Tian JG, Li YG, and Li JM (2013) Reversion effects of curcumin on multidrug resistance of MNNG/HOS human osteosarcoma cells in vitro and in vivo through regulation of P-glycoprotein. *Chin Med J (Engl)* **126**:4116-4123.
- Simonson SG, Raza A, Martin PD, Mitchell PD, Jarcho JA, Brown CD, Windass AS, and Schneck DW (2004) Rosuvastatin pharmacokinetics in heart transplant recipients administered an antirejection regimen including cyclosporine. *Clin Pharmacol Ther* **76**:167-177.
- Sparreboom A, Gelderblom H, Marsh S, Ahluwalia R, Obach R, Principe P, Twelves C, Verweij J, and McLeod HL (2004) Diflomotecan pharmacokinetics in relation to ABCG2 421C>A genotype. *Clin Pharmacol Ther* **76**:38-44.
- Sparreboom A, Loos WJ, Burger H, Sissung TM, Verweij J, Figg WD, Nooter K, and Gelderblom H (2005) Effect of ABCG2 genotype on the oral bioavailability of topotecan. *Cancer Biol Ther* **4**:650-658.
- Stewart CF, Tagen M, Schwartzberg LS, Blakely LJ, Tauer KW, and Smiley LM (2014) Phase I dosage finding and pharmacokinetic study of intravenous topotecan and oral erlotinib in adults with refractory solid tumors. *Cancer Chemother Pharmacol* **73**:561-568.
- Sugimoto H, Matsumoto S, Tachibana M, Niwa S, Hirabayashi H, Amano N, and Moriwaki T (2011) Establishment of in vitro P-glycoprotein inhibition assay and its exclusion criteria to assess the risk of drug-drug interaction at the drug discovery stage. *J Pharm Sci* **100**:4013-4023.
- Suzuki K, Doki K, Homma M, Tamaki H, Hori S, Ohtani H, Sawada Y, and Kohda Y (2009) Co-administration of proton pump inhibitors delays elimination of plasma methotrexate in high-dose methotrexate therapy. *Br J Clin Pharmacol* **67**:44-49.

- Swaisland HC, Ranson M, Smith RP, Leadbetter J, Laight A, McKillop D, and Wild MJ (2005) Pharmacokinetic drug interactions of gefitinib with rifampicin, itraconazole and metoprolol. *Clin Pharmacokinet* **44**:1067-1081.
- Tamaki H, Satoh H, Hori S, Ohtani H, and Sawada Y (2010) Inhibitory effects of herbal extracts on breast cancer resistance protein (BCRP) and structure-inhibitory potency relationship of isoflavonoids. *Drug Metab Pharmacokinet* **25**:170-179.
- Tamura A, Watanabe M, Saito H, Nakagawa H, Kamachi T, Okura I, and Ishikawa T (2006) Functional validation of the genetic polymorphisms of human ATP-binding cassette (ABC) transporter ABCG2: identification of alleles that are defective in porphyrin transport. *Mol Pharmacol* **70**:287-296.
- Tanaka M, Kamada I, Takahashi J, Kimura K, Matsukura H, and Tani Y (2014) Defining the Jr(a-) phenotype in the Japanese population. *Transfusion* **54**:412-417.
- Tang SC, Lagas JS, Lankheet NA, Poller B, Hillebrand MJ, Rosing H, Beijnen JH, and Schinkel AH (2012a) Brain accumulation of sunitinib is restricted by P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) and can be enhanced by oral elacridar and sunitinib coadministration. *Int J Cancer* **130**:223-233.
- Tang SC, Lankheet NA, Poller B, Wagenaar E, Beijnen JH, and Schinkel AH (2012b) P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) restrict brain accumulation of the active sunitinib metabolite N-desethyl sunitinib. *J Pharmacol Exp Ther* **341**:164-173.
- Thomas KS, Billingsley A, Amarshi N, and Nair BA (2010) Elevated international normalized ratio associated with concomitant warfarin and erlotinib. *Am J Health Syst Pharm* **67**:1426-1429.

- Tiwari AK, Sodani K, Wang SR, Kuang YH, Ashby CR, Jr., Chen X, and Chen ZS (2009)
Nilotinib (AMN107, Tasisna) reverses multidrug resistance by inhibiting the activity of
the ABCB1/Pgp and ABCG2/BCRP/MXR transporters. *Biochem Pharmacol* **78**:153-161.
- Tomaru A, Morimoto N, Morishita M, Takayama K, Fujita T, Maeda K, Kusuhara H, and
Sugiyama Y (2013) Studies on the intestinal absorption characteristics of sulfasalazine, a
breast cancer resistance protein (BCRP) substrate. *Drug Metab Pharmacokinet* **28**:71-74.
- Tran HT, Zinner RG, Blumenschein GR, Jr., Oh YW, Papadimitrakopoulou VA, Kim ES, Lu C,
Malik M, Lum BL, and Herbst RS (2011) Pharmacokinetic study of the phase III,
randomized, double-blind, multicenter trial (TRIBUTE) of paclitaxel and carboplatin
combined with erlotinib or placebo in patients with advanced Non-small Cell Lung
Cancer (NSCLC). *Invest New Drugs* **29**:499-505.
- Tsamandouras N, Dickinson G, Guo Y, Hall S, Rostami-Hodjegan A, Galetin A, and Aarons L
(2014) Identification of multiple polymorphisms effect on the pharmacokinetics of
simvastatin and simvastatin acid using a population modeling approach. *Clin Pharmacol
Ther.*
- Urquhart BL, Ware JA, Tirona RG, Ho RH, Leake BF, Schwarz UI, Zaher H, Palandra J, Gregor
JC, Dresser GK, and Kim RB (2008) Breast cancer resistance protein (ABCG2) and drug
disposition: intestinal expression, polymorphisms and sulfasalazine as an in vivo probe.
Pharmacogenet Genomics **18**:439-448.
- van de Steeg E, Greupink R, Schreurs M, Nooijen IH, Verhoeckx KC, Hanemaaijer R, Ripken D,
Monshouwer M, Vlaming ML, DeGroot J, Verwei M, Russel FG, Huisman MT, and
Wortelboer HM (2013) Drug-drug interactions between rosuvastatin and oral antidiabetic
drugs occurring at the level of OATP1B1. *Drug Metab Dispos* **41**:592-601.

- Varma MV, Scialis RJ, Lin J, Bi YA, Rotter CJ, Goosen TC, and Yang X (2014) Mechanism-based pharmacokinetic modeling to evaluate transporter-enzyme interplay in drug interactions and pharmacogenetics of glyburide. *Aaps J* **16**:736-748.
- Veeraputhiran M and Sundermeyer M (2008) Rhabdomyolysis resulting from pharmacologic interaction between erlotinib and simvastatin. *Clin Lung Cancer* **9**:232-234.
- Volk EL and Schneider E (2003) Wild-type breast cancer resistance protein (BCRP/ABCG2) is a methotrexate polyglutamate transporter. *Cancer Res* **63**:5538-5543.
- Wang L, Leggas M, Empey PE, and McNamara PJ (2012) Stereoselective interaction of pantoprazole with ABCG2. II. In vitro flux analysis. *Drug Metab Dispos* **40**:1024-1031.
- Wang Q, Strab R, Kardos P, Ferguson C, Li J, Owen A, and Hidalgo IJ (2008) Application and limitation of inhibitors in drug-transporter interactions studies. *Int J Pharm* **356**:12-18.
- Wang Y, Wang M, Qi H, Pan P, Hou T, Li J, He G, and Zhang H (2014) Pathway-dependent inhibition of paclitaxel hydroxylation by kinase inhibitors and assessment of drug-drug interaction potentials. *Drug Metab Dispos* **42**:782-795.
- Ware JA, Dalziel G, Jin JY, Pellett JD, Smelick GS, West DA, Salphati L, Ding X, Sutton R, Fridyland J, Dresser MJ, Morrisson G, and Holden SN (2013) Impact of food and the proton pump inhibitor rabeprazole on the pharmacokinetics of GDC-0941 in healthy volunteers: bench to bedside investigation of pH-dependent solubility. *Mol Pharm* **10**:4074-4081.
- Watanabe T, Kusuhara H, Maeda K, Shitara Y, and Sugiyama Y (2009) Physiologically based pharmacokinetic modeling to predict transporter-mediated clearance and distribution of pravastatin in humans. *J Pharmacol Exp Ther* **328**:652-662.

- Wei Y, Wu B, Jiang W, Yin T, Jia X, Basu S, Yang G, and Hu M (2013) Revolving door action of breast cancer resistance protein (BCRP) facilitates or controls the efflux of flavone glucuronides from UGT1A9-overexpressing HeLa cells. *Mol Pharm* **10**:1736-1750.
- Weiss J and Haefeli WE (2013) Potential of the novel antiretroviral drug rilpivirine to modulate the expression and function of drug transporters and drug-metabolising enzymes in vitro. *Int J Antimicrob Agents* **41**:484-487.
- Wilding G, Soulie P, Trump D, Das-Gupta A, and Small E (2006) Results from a pilot Phase I trial of gefitinib combined with docetaxel and estramustine in patients with hormone-refractory prostate cancer. *Cancer* **106**:1917-1924.
- Wright JA, Haslam IS, Coleman T, and Simmons NL (2011) Breast cancer resistance protein BCRP (ABCG2)-mediated transepithelial nitrofurantoin secretion and its regulation in human intestinal epithelial (Caco-2) layers. *Eur J Pharmacol* **672**:70-76.
- Xia CQ, Liu N, Miwa GT, and Gan LS (2007) Interactions of cyclosporin a with breast cancer resistance protein. *Drug Metab Dispos* **35**:576-582.
- Xiao G, Black C, Hetu G, Sands E, Wang J, Caputo R, Rohde E, and Gan LS (2012) Cerebrospinal fluid can be used as a surrogate to assess brain exposures of breast cancer resistance protein and P-glycoprotein substrates. *Drug Metab Dispos* **40**:779-787.
- Xiao Y, Davidson R, Smith A, Pereira D, Zhao S, Soglia J, Gebhard D, de Morais S, and Duignan DB (2006) A 96-well efflux assay to identify ABCG2 substrates using a stably transfected MDCK II cell line. *Mol Pharm* **3**:45-54.
- Yamamoto N, Horiike A, Fujisaka Y, Murakami H, Shimoyama T, Yamada Y, and Tamura T (2008) Phase I dose-finding and pharmacokinetic study of the oral epidermal growth

- factor receptor tyrosine kinase inhibitor Ro50-8231 (erlotinib) in Japanese patients with solid tumors. *Cancer Chemother Pharmacol* **61**:489-496.
- Yamasaki Y, Ieiri I, Kusuhara H, Sasaki T, Kimura M, Tabuchi H, Ando Y, Irie S, Ware J, Nakai Y, Higuchi S, and Sugiyama Y (2008) Pharmacogenetic characterization of sulfasalazine disposition based on NAT2 and ABCG2 (BCRP) gene polymorphisms in humans. *Clin Pharmacol Ther* **84**:95-103.
- Yanase K, Tsukahara S, Asada S, Ishikawa E, Imai Y, and Sugimoto Y (2004) Gefitinib reverses breast cancer resistance protein-mediated drug resistance. *Mol Cancer Ther* **3**:1119-1125.
- Yang JJ, Milton MN, Yu S, Liao M, Liu N, Wu JT, Gan L, Balani SK, Lee FW, Prakash S, and Xia CQ (2010) P-glycoprotein and breast cancer resistance protein affect disposition of tandutinib, a tyrosine kinase inhibitor. *Drug Metab Lett* **4**:201-212.
- Ye L, Yang X, Yang Z, Gao S, Yin T, Liu W, Wang F, Hu M, and Liu Z (2013) The role of efflux transporters on the transport of highly toxic aconitine, mesaconitine, hypaconitine, and their hydrolysates, as determined in cultured Caco-2 and transfected MDCKII cells. *Toxicol Lett* **216**:86-99.
- Yin OQ, Gallagher N, Fischer D, Zhao L, Zhou W, Leroy E, Golor G, and Schran H (2011) Effects of nilotinib on single-dose warfarin pharmacokinetics and pharmacodynamics: a randomized, single-blind, two-period crossover study in healthy subjects. *Clin Drug Investig* **31**:169-179.
- Yoshida N, Takada T, Yamamura Y, Adachi I, Suzuki H, and Kawakami J (2008) Inhibitory effects of terpenoids on multidrug resistance-associated protein 2- and breast cancer resistance protein-mediated transport. *Drug Metab Dispos* **36**:1206-1211.

- Zaher H, Khan AA, Palandra J, Brayman TG, Yu L, and Ware JA (2006) Breast cancer resistance protein (Bcrp/abcg2) is a major determinant of sulfasalazine absorption and elimination in the mouse. *Mol Pharm* **3**:55-61.
- Zamek-Gliszczyński MJ, Bedwell DW, Bao JQ, and Higgins JW (2012) Characterization of SAGE Mdr1a (P-gp), Bcrp, and Mrp2 knockout rats using loperamide, paclitaxel, sulfasalazine, and carboxydichlorofluorescein pharmacokinetics. *Drug Metab Dispos* **40**:1825-1833.
- Zelinski T, Coghlan G, Liu XQ, and Reid ME (2012) ABCG2 null alleles define the Jr(a-) blood group phenotype. *Nat Genet* **44**:131-132.
- Zhang D, He K, Herbst JJ, Kolb J, Shou W, Wang L, Balimane PV, Han YH, Gan J, Frost CE, and Humphreys WG (2013) Characterization of efflux transporters involved in distribution and disposition of apixaban. *Drug Metab Dispos* **41**:827-835.
- Zhang D, Zhu M, and Humphreys WG (2008) *Drug metabolism in drug design and development: Basic concepts and practice*. Wiley-Interscience, Hoboken, N.J.
- Zhang S, Wang X, Sagawa K, and Morris ME (2005a) Flavonoids chrysin and benzoflavone, potent breast cancer resistance protein inhibitors, have no significant effect on topotecan pharmacokinetics in rats or mdr1a/1b (-/-) mice. *Drug Metab Dispos* **33**:341-348.
- Zhang W, Yu BN, He YJ, Fan L, Li Q, Liu ZQ, Wang A, Liu YL, Tan ZR, Fen J, Huang YF, and Zhou HH (2006) Role of BCRP 421C>A polymorphism on rosuvastatin pharmacokinetics in healthy Chinese males. *Clin Chim Acta* **373**:99-103.
- Zhang Y, Gupta A, Wang H, Zhou L, Vethanayagam RR, Unadkat JD, and Mao Q (2005b) BCRP transports dipyridamole and is inhibited by calcium channel blockers. *Pharm Res* **22**:2023-2034.

- Zhou L, Schmidt K, Nelson FR, Zelesky V, Troutman MD, and Feng B (2009) The effect of breast cancer resistance protein and P-glycoprotein on the brain penetration of flavopiridol, imatinib mesylate (Gleevec), prazosin, and 2-methoxy-3-(4-(2-(5-methyl-2-phenyloxazol-4-yl)ethoxy)phenyl)propanoic acid (PF-407288) in mice. *Drug Metab Dispos* **37**:946-955.
- Zhou Q, Chen QX, Ruan ZR, Yuan H, Xu HM, and Zeng S (2013) CYP2C9*3(1075A > C), ABCB1 and SLCO1B1 genetic polymorphisms and gender are determinants of inter-subject variability in pitavastatin pharmacokinetics. *Pharmazie* **68**:187-194.
- Zouboulis-Vafiadis I, Paraskevas E, Tzourmakliotis D, Hatzikyriakou M, Mestoussi A, Vasdekis V, Katsilabros N, Arhimandritis A, and Papadokostopoulou A (2014) An open-label, non-comparative, non-interventional, multi-center, post-authorization safety study on the administration of rabeprazole to adults with gastro-esophageal reflux disease. *Ann Gastroenterol* **27**:133-138.
- Zvyaga T, Chang SY, Chen C, Yang Z, Vuppugalla R, Hurley J, Thorndike D, Wagner A, Chimalakonda A, and Rodrigues AD (2012) Evaluation of six proton pump inhibitors as inhibitors of various human cytochromes P450: focus on cytochrome P450 2C19. *Drug Metab Dispos* **40**:1698-1711.

Table 1. Summary of rosuvastatin BCRP pharmacogenomic studies and associated effect on its pharmacokinetics

Dose*	Subject Race & Sex	BCRP c.421C>A Genotype	OATP1B1 Phenotype	Effect of c.421C>A SNP	Reference
20 mg SD	Asian Chinese male	7- CC, 5 –CA, 2-AA	All subjects had OATP1B1 521TT and CYP2C9*1/*1 genotypes ((wild type)	1.78-fold AUC increase in 421CA/AA genotype (combined) ^a	Zhang 2006
20 mg SD	Caucasian (50% female)	16-CC 12-CA, 4-AA	Individuals with the OATP1B1 521CC genotype not included ^b	1.2-fold AUC increase in 421CA (NS); 2.44- and 2.3-fold AUC and C _{max} increase, respectively in 421AA ^a	Keskitalo 2009b
5, 10 and 20 mg SD (1 week washout)	Asian Chinese	15-CC, 15-CA, 6-AA	No significant effect of CYP2C9*3 and OATP1B1 c.521 T > C	1.6-fold AUC increase in 421AA ^c	Zhou 2013
10 mg/day (min 4 weeks)	Asian Chinese (52% female)	129-CC, 108- CA, 39-AA	205- OATP1B1 521TT and 62/5- TC/CC. Rosuvastatin plasma concentrations affected by both BCRP 421C>A and OATP1B1 521T>C ^e	2.2-fold increase in mean plasma concentrations in 421AA ^{a,d} BCRP 421C>A (but not OATP1B1521 T>C) strongly associated with the LDL-C response to rosuvastatin	Lee 2013

*The route of administration was oral in all cases

^a Denotes significant change, p<0.05

^b Subjects had ABCB1 1199GG reference genotype

^c Dose-normalized changes (based on 5 mg)

^d 1.99-fold increase in mean plasma concentration of N-desmethyl rosuvastatin

^e No effect of CYP2C9*1/*3 or *3/*3 on rosuvastatin PK or PD

Table 2. Summary of clinical studies using sulfasalazine as the intestinal BCRP probe.

Dose / Formulation*	Subject Race & Sex	BCRP c.421C>A Genotype	NAT2 Phenotype	Sulfasalazine AUC Variability	Effect of c.421C>A SNP or BCRP inhibitor on PK	Reference
2000 mg IR tablet	Asian men	12-CC, 16-CA, 9-AA	13-RA, 16-IA, 8-SA	1.9-fold (95% CI, genotype controlled)	1.9-fold AUC increase in 421CA, 3.5-fold AUC increase in 421AA; clear gene-dose effect	Yamasaki et al., 2008
1000 mg suspension (also tested IR tablet)	Caucasian men and women	9-CC, 5-CA, 0-AA	Not genotyped, sulfapyridine PK not studied	20.5-fold (max/min; genotype controlled)	2.4-fold AUC increase in 421CA genotype	Urquhart et al., 2008
500 mg enteric-coated tablet	Asian men	12-CC, 12-CA, 12-AA	4-RA, 20-IA, 12-SA	21- to 81-fold (max/min; genotype controlled)	None (trend for 2-fold AUC increase in AA genotype, NS); no clear gene-dose effect; no significant effect of BCRP inhibitor (40 mg pantoprazole)	Adkison et al., 2010
2000 mg / 100 µg IR tablet / suspension	Asian men	8-CC	Not genotyped, sulfapyridine PK not studied	AUC CV = 24-40%	Curcumin (2 g*) increased AUC 3.2-fold (therapeutic dose) and 2-fold (microdose)	Kushuhara et al., 2012

*The route of administration was oral in all cases

RA – rapid acetylator; IA – intermediate acetylator, SA – slow acetylator

Table 3. Compounds considered as in vivo human BCRP substrates based on available in vitro data.

Object	Object Therapeutic Class	Experimental System	Efflux Ratio ^a	K _m (μM)
(-)-pantoprazole	Proton Pump Inhibitors	MDCK-transfected cells		0.59 ¹
(+)-pantoprazole	Proton Pump Inhibitors	MDCK-transfected cells		5.32 ¹
2'-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		5.4 ²
3,2'-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		35.9 ²
3,3'-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		45.2 ²
3,4'-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		35.9 ²
3,5-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		12.8 ²
3,5-di-O-caffeoylquinic acid	Food Products	Caco-2 cells	5.1 ³	
3,6-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		31.9 ²
3,7-dihydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		3.5 ²
3'-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		5.6 ²
3-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		9.7 ²
4-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		17.5 ²
5-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		48.5 ²
6-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		26.8 ²
7-hydroxyflavone-O-glucuronide	Food Products	HeLa-transfected cells		9 ²

aconitine	Herbal Medications	MDCK–transfected cells	19.8 ⁴	
apixaban	Anticoagulants and Antiplatelets	MDCK–transfected cells	12 ⁵	
axitinib	Kinase Inhibitors	MDCK–transfected cells	3.4 ⁶	
basertib hydroxy–QPA	Kinase Inhibitors	MDCK–transfected cells	2.31 ⁷	
boceprevir	Antivirals	MDCK–transfected cells	2 ⁸	
bupropion	Dopamine–Norepinephrine Reuptake Inhibitors	Membrane vesicles		3 ⁹
canertinib	Kinase Inhibitors	MDCK–transfected cells	2.8 ¹⁰	
chlorothiazide	Diuretics	MDCK–transfected cells	36 ¹¹	
cimetidine	H–2 Receptor Antagonists	MDCK–transfected cells	3.4 ¹²	
ciprofloxacin	Antibiotics	MDCK–transfected cells	2.9 ¹³	
dantrolene	Muscle Relaxants	Caco–2 cells	5 ¹⁴	
daunorubicin	Anthracyclines	Caco–2 cells	4.9 ¹⁵	
dimenhydrinate	H–1 Receptor Antagonists	Caco–2 cells	9.8 ¹⁶	
dipyridamole	Anticoagulants and Antiplatelets	MDCK–transfected cells	2.7 ¹⁷	
dolutegravir	HIV–Integrase Strand Transfer Inhibitors	MDCK–transfected cells	3.1 ¹⁸	
edaravone sulfate	Neuroprotectors	Membrane vesicles		16.5 ¹⁹
erythromycin	Antibiotics	Caco–2 cells	8.1 ¹⁵	
estradiol–17– β –glucuronide	Estrogens	Membrane vesicles		44.2 ²⁰
estrone–3–sulfate	Estrogens	Caco–2 cells	48.5 ²¹	6.8 ^{22,b}
ethinyl estradiol sulfate	Estrogens	Membrane vesicles		2.9 ²³
etoposide	Topoisomerase Inhibitors	Caco–2 cells	12.7 ¹⁵	
flavopiridol (alvocidib)	Kinase Inhibitors	MDCK–transfected cells	6 ²⁴	

fluvastatin	HMG CoA Reductase Inhibitors (Statins)	Caco-2 cells	4.9 ¹⁵	
furosemide	Diuretics	Caco-2 cells	17.5 ¹⁵	
gemifloxacin	Antibiotics	MDCK-transfected cells		145.69 ²⁵
genistein	Food Products	LLC-PK1-transfected cells	2 ²⁶	
glyburide (glibenclamide)	Sulfonylureas	MDCK-transfected cells	9.3 ²⁷	13.07 ^{28,c}
grepafloxacin	Antibiotics	MDCK-transfected cells	9.14 ²⁹	
hematoporphyrin	Photosensitizing agent	Membrane vesicles		17.8 ³⁰
hypaconitine	Herbal Medications	MDCK-transfected cells	7.6 ⁴	
imatinib	Kinase Inhibitors	MDCK-transfected cells	2.4 ²⁴	
irinotecan	Topoisomerase Inhibitors	Caco-2 cells	3.7 ¹⁵	
-SN-38	Topoisomerase Inhibitors	MDCK-transfected cells	8.4 ¹⁰	4 ^{31,b}
-SN-38 glucuronide	Topoisomerase Inhibitors	Membrane vesicles		26 ³¹
lamivudine	Nucleoside Reverse Transcriptase Inhibitors (NRTIs)	MDCK-transfected cells	3 ³²	216.5 ³²
lapatinib	Kinase Inhibitors	MDCK-transfected cells	2.6 ³³	
mesaconitine	Herbal Medications	MDCK-transfected cells	11.1 ⁴	
methotrexate	Antimetabolites	MDCK-transfected cells	6.78 ³⁴	681 ^{35,b}
mitoxantrone	Other Antineoplastics	MDCK-transfected cells	6.1 ¹⁰	
nitrofurantoin	Anti-Infective Agents	Caco-2 cells	24.1 ¹⁵	69.4 ³⁶
norfloxacin	Antibiotics	Caco-2 cells	3 ¹⁵	
olomoucine II	Kinase Inhibitors	MDCK-transfected cells	2.27 ³⁷	
pemetrexed	Antimetabolites	Membrane vesicles		390 ³⁸

pitavastatin	HMG CoA Reductase Inhibitors (Statins)	Membrane vesicles		5.73 ³⁹
prazosin	Alpha/Beta Adrenergic Antagonists	MDCK–transfected cells	11.1 ⁴⁰	
rivaroxaban	Anticoagulants and Antiplatelets	Caco–2 cells	2.82 ⁴¹	
rosuvastatin	HMG CoA Reductase Inhibitors (Statins)	Caco–2 cells	83.2 ⁴⁰	2.02 ^{42,b}
sildenafil	Erectile Dysfunction Treatments	MDCK–transfected cells	3.59 ⁴³	22.4 ⁴³
sorafenib	Kinase Inhibitors	MDCK–transfected cells	2.7 ⁴⁴	
sulfasalazine	NSAIDS	Caco–2 cells	87 ⁴⁵	0.7 ^{46,b}
sunitinib	Kinase Inhibitors	MDCK–transfected cells	1.9 ^{47,d}	
–N–desethyl sunitinib (SU12662)	Kinase Inhibitors	MDCK–transfected cells	13.5 ⁴⁸	
tandutinib	Kinase Inhibitors	Caco–2 cells	5 ⁴⁹	4 ⁴⁹
temocapril	Angiotensin Converting Enzyme (ACE) Inhibitors	Caco–2 cells	5.56 ⁵⁰	
teniposide	Topoisomerase Inhibitors	Caco–2 cells	10.8 ¹⁵	
teriflunomide	Other Immunomodulators	Caco–2 cells	9.37 ⁵¹	
topotecan	Topoisomerase Inhibitors	Caco–2 cells	6.9 ¹⁵	213.28 ^{52,e}
troglitazone sulfate	Thiazolidinediones	MDCK–transfected cells	6.42 ⁵³	
ulifloxacin	Antibiotics	MDCK–transfected cells	2.95 ²⁹	
varденаfil	Erectile Dysfunction Treatments	MDCK–transfected cells	4.12 ⁴³	14.3 ⁴³

Metabolites are listed below the respective parent compound. K_m and efflux ratio values are presented as reported in the respective references.

^a compounds with efflux ratios >2 were selected for further consideration

^b kinetic experiments were performed in membrane vesicles

^c kinetic experiments were performed in HEK–transfected cells

^d included in the list because metabolite has an efflux ratio >2

^e kinetic experiments were performed in MDCK–transfected cells

¹ (Wang et al., 2012); ² (Wei et al., 2013); ³ (Farrell et al., 2012); ⁴ (Ye et al., 2013); ⁵ (Zhang et al., 2013); ⁶ FDA (2012) Drug approval package: INLYTA[®] (axitinib) [FDA application no, (NDA) 202324].; ⁷ (Marchetti et al., 2013); ⁸ (Chu et al., 2013a); ⁹ (Hemauer et al., 2010); ¹⁰ (Xiao et al., 2006); ¹¹ (Beery et al., 2012); ¹² (Pavek et al., 2005); ¹³ (Merino et al., 2006); ¹⁴ (Xiao et al., 2012); ¹⁵ (Lin et al., 2011); ¹⁶ (Crowe and Wright, 2012); ¹⁷ (Zhang et al., 2005b); ¹⁸ (Reese et al., 2013); ¹⁹ (Mizuno et al., 2007); ²⁰ (Chen et al., 2003); ²¹ (Li et al., 2011a); ²² (Imai et al., 2003); ²³ (Han et al., 2010); ²⁴ (Zhou et al., 2009); ²⁵ (Jin et al., 2013); ²⁶ (Imai et al., 2004); ²⁷ (Varma et al., 2014); ²⁸ (Pollex et al., 2010); ²⁹ (Ando et al., 2007); ³⁰ (Tamura et al., 2006); ³¹ (Nakatomi et al., 2001); ³² (Kim et al., 2007); ³³ (Polli et al., 2008); ³⁴ (Xia et al., 2007); ³⁵ (Volk and Schneider, 2003); ³⁶ (Wright et al., 2011); ³⁷ (Hofman et al., 2013); ³⁸ (Li et al., 2011b); ³⁹ (Hirano et al., 2005); ⁴⁰ (Li et al., 2012); ⁴¹ (Gong et al., 2013); ⁴² (Kitamura et al., 2008); ⁴³ (Choi and Song, 2012); ⁴⁴ (Lagas et al., 2010); ⁴⁵ (Wang et al., 2008); ⁴⁶ (Jani et al., 2009); ⁴⁷ (Tang et al., 2012a); ⁴⁸ (Tang et al., 2012b); ⁴⁹ (Yang et al., 2010); ⁵⁰ (Ohura et al., 2011); ⁵¹ FDA (2012) Drug approval package: AUBAGIO[®] (teriflunomide) [FDA application no, (NDA) 202992].; ⁵² (Li et al., 2008); ⁵³ (Enokizono et al., 2007a)

Table 4. *In vivo* drug interaction studies involving rosuvastatin and sulfasalazine indicating at least partial contribution by BCRP.

Object	Object Dose*	Precipitant	Precipitant Therapeutic Class	Precipitant Dose*	% Change in AUC ^a
sulfasalazine	2 g SD	curcumin	Food Products	2 g SD	220 ¹
	500 mg SD	pantoprazole	Proton Pump Inhibitors	40 mg SD	83 ^{2,b}
	100 µg SD	curcumin	Food Products	2 g SD	83 ¹
rosuvastatin	10 mg QD x 10D	cyclosporine	Immunosuppressants	75 to 200 mg BID (stable dose for at least 2 months)	610 ^{3,c}
	10 mg SD	atazanavir + ritonavir	Protease Inhibitors	300 mg atazanavir + 100 mg ritonavir QD x 6D	210 ⁴
	20 mg QD x 7D	lopinavir + ritonavir	Protease Inhibitors	400 mg lopinavir + 100 mg ritonavir BID x 7D	110 ⁵
	10 mg SD	eltrombopag	Platelet Stimulating Agents	75 mg QD x 5D	55 ⁶
	10 mg SD	elvitegravir + cobicistat	AIDS Treatments	150 mg elvitegravir + 150 mg cobicistat QD x 10D	40 ⁷
10 mg SD	tipranavir + ritonavir	Protease Inhibitors	500 mg tipranavir + 200 mg ritonavir BID x 11D	37 ⁸	

*The route of administration was oral in all cases

^a - Values were extracted from the DIDB and calculated as the change in mean AUC values.

^b - AUC change observed with 421A/C genotype and was not statistically different from control

^c - change in AUC was calculated using healthy historical controls

¹ (Kusuhara et al., 2012); ² (Adkison et al., 2010); ³ (Simonson et al., 2004); ⁴ (Busti et al., 2008); ⁵ (Kiser et al., 2008); ⁶ (Allred et al., 2011); ⁷ (Custodio et al., 2014); ⁸ (Pham et al., 2009)

Table 5. Compounds considered as in vivo BCRP inhibitors, based on in vitro IC₅₀ or K_i values < 10 μM

Precipitant	Therapeutic Class	K _i (μM)	IC ₅₀ (μM)
afatinib	Kinase Inhibitors	–	0.75 ¹
aripiprazole	Antipsychotics	–	3.5 ²
axitinib	Kinase Inhibitors	–	4.4 ³
curcumin	Food Products	–	1.6 ⁴
cyclosporine	Immunosuppressants	6.7 ⁵	13.4 ^a
erlotinib	Kinase Inhibitors	0.15 ⁶	0.13 ⁶
elacridar	drug distribution modulator	–	0.31 ⁷
fluvastatin	HMG CoA Reductase Inhibitors	5.43 ⁸	10.86 ^a
fumitremorgin C (FTC)	None	0.55 ⁶	0.25 ⁶
gefitinib	Kinase Inhibitors	–	1.01 ⁹
ivermectin	Antiparasitics	1.4 ¹⁰	2.8 ^a
ko143	Transporter Modulators	–	0.013 ⁵
lapatinib	Kinase Inhibitors	–	0.025 ¹¹
nilotinib	Kinase Inhibitors	0.69 ¹²	1.38 ^a
novobiocin	Antibiotics	–	0.063 ¹³
pantoprazole	Proton Pump Inhibitors	–	5.5 ¹⁴
pitavastatin	HMG CoA Reductase Inhibitors	2.92 ⁸	5.84 ^a
ponatinib	Kinase Inhibitors	–	0.013 ¹⁵
quercetin	Food Products	–	0.6 ¹⁶
quizartinib (AC220)	Kinase Inhibitors	–	0.5 ¹⁷
rabeprazole	Proton Pump Inhibitors	–	8.5 ¹⁴
regorafenib	Kinase Inhibitors	–	0.0447 ¹⁸
rilpivirine	NonNucleoside Reverse Transcriptase Inhibitors	–	1.5 ¹⁹
sulfasalazine	NSAIDS	–	0.46 ¹³
sunitinib	Kinase Inhibitors	0.32 ²⁰	0.64 ^a
tacrolimus	Immunosuppressants	–	6 ¹⁶
teriflunomide	Other Immunomodulators	–	0.146 ²¹

trametinib	Kinase Inhibitors	–	1.1 ²²
trifluoperazine	Antipsychotics	–	7.56 ²³
vismodegib	Cancer Treatments	–	2.4 ²⁴

IC₅₀ and K_i values are presented as reported in the respective references.

^a IC₅₀ value was calculated as 2xK_i, assuming linearity

¹ FDA (2013) Drug approval package: GILOTRIF® (afatinib dimaleate) [FDA application no, (NDA) 201292]; ² (Nagasaka et al., 2012); ³ (Reyner et al., 2013); ⁴ (Kusuhara et al., 2012); ⁵ (Xia et al., 2007); ⁶ (Noguchi et al., 2009); ⁷ (Si et al., 2013); ⁸ (Hirano et al., 2005); ⁹ (Yanase et al., 2004); ¹⁰ (Jani et al., 2011); ¹¹ (Polli et al., 2008); ¹² (Tiwari et al., 2009); ¹³ (Elsby et al., 2011); ¹⁴ (Suzuki et al., 2009); ¹⁵ FDA (2012) Drug approval package: ICLUSIG® (ponatinib hydrochloride) [FDA application no, (NDA) 203469]; ¹⁶ (Saito et al., 2006); ¹⁷ (Bhullar et al., 2013); ¹⁸ FDA (2012) Drug approval package: STIVARGA® (regorafenib) [FDA application no, (NDA) 203085]; ¹⁹ (Weiss and Haefeli, 2013); ²⁰ (Kawahara et al., 2010); ²¹ FDA (2012) Drug approval package: AUBAGIO® (teriflunomide) [FDA application no, (NDA) 202992]; ²² FDA (2013) Drug approval package: MEKINIST® (trametinib dimethyl sulfoxide) [FDA application no, (NDA) 204114]; ²³ (Pan et al., 2013); ²⁴ FDA (2012) Drug approval package: ERIVEDGE® (vismodegib) [FDA application no, (NDA) 203388]

Table 6. In vitro and in vivo data and considerations for potential systemic BCRP inhibitors.

Precipitant	I ₁ (μ M, C _{max})	Plasma Binding (%)	Dose	BCRP parameters			Other possible systemic interactions				
				K _i (μ M)	IC ₅₀ (μ M)	[I ₁]/IC ₅₀	In vivo PK-based DDI data	Transporter / Enzyme	IC ₅₀ (μ M)	[I ₁]/IC ₅₀	In vivo PK-based DDI data
regorafenib	8.1 ¹	99.5 ¹	160 mg (SS) ¹	–	0.0447 TOP ²	180	No published inhibition data ³	P-gp	0.8 ²	10	
								CYP3A4	22.2 ^{a,2}	0.36	
								CYP2C8	1.2 ^{a,2}	6.8	
								CYP2C9	9.4 ^{a,2}	0.86	
								CYP2B6	10.4 ^{a,2}	0.78	
								CYP2C19	32.8 ^{a,2}	0.25	
								UGT1A1	6 ²	1.4	SN-38 glucuronide AUC +44% (when co-administered with irinotecan) ²
UGT1A9	4.2 ²	1.9									
sulfasalazine	79 ⁴	>99.7 ⁴	3–4 g SD ⁴	–	0.46 E3S ⁵	170	No effect with methotrexate ⁶ , however in vivo substrate ⁷	OATP1B1	0.56 ⁸	140	
								OATP1B3	12 ⁸	6.6	
								OATP2B1	3 ⁸	26	

lapatinib	4.2 ^{b,9}	>99 ⁹	1250 mg QD (SS) ⁹	-	0.025 CIM ¹⁰	170 ^c	Decreased topotecan clearance, -18% (renal -22%) ¹¹ , SN-38 +45% AUC ¹²	P-gp	1.5 ¹³	2.8	Digoxin AUC +2.8-fold ⁹
								OATP1B1	4 ¹⁰	1.0	SN-38 AUC +45% ¹²
								CYP3A	39 ¹⁴	0.11	Midazolam AUC +45% (oral) +22% (IV) ⁹
								CYP2D6	13.8 ¹⁴	0.30	
								CYP2C8	1.2 ^{a,14}	3.5	Paclitaxel AUC +23% ⁸
								CYP2C9	11.8 ¹⁴	0.36	
								CYP2C19	27.2 ¹⁴	0.15	
erlotinib	6.06 ¹⁵	90 ¹⁵	150 mg QD x 21D ¹⁵	0.15 E3S ¹⁶	0.13 E3S ¹⁶	47	No effect on topotecan ¹⁷	P-gp	2 ¹⁶	3.0	
								OATP1B1	21 ⁸	0.29	
								OATP1B3	44 ⁸	0.14	
								OATP2B1	0.55 ⁸	11	
								CYP3A4	2.56 ^{a,18}	2.4	2-fold increase in exposure of OSI-930 ²³ , everolimus AUC +21% ²⁴ , case report of toxicity with simvastatin ²⁵
								CYP3A5	80 ^{a,19}	0.08	2-fold increase in exposure of OSI-930 ²³ , everolimus AUC +21% ²⁴ , case report of toxicity with simvastatin ²⁵

DMD Fast Forward. Published on January 13, 2015 as DOI: 10.1124/dmd.114.062174
This article has not been copyedited and formatted. The final version may differ from this version.

									CYP2D6	37% ²⁰	–	
									CYP2C8	8.04 ^{a,18}	0.75	No effect on paclitaxel ²⁶
									CYP2C9	49% ²⁰	–	Case report of increased INR with warfarin ²⁷
									CYP1A2	30% ²⁰	–	
									CYP2C19	42.9 ²¹	0.14	
									UGT1A1	1.28 ^{a,22}	4.7	No effect on irinotecan/SN-38 ²⁸
									P-gp	1.7 ²⁹	2.5	
									OCT1	2.92 ³¹	0.47	
									OCT2	>30 ³¹		
									OCT3	0.345 ³¹	4.0	
nilotinib	4.3 ²⁹	98 ²⁹	400 mg BID x 15D ²⁹	0.69 MTX ³⁰	1.38	3.1	No inhibition data		CYP3A4	0.4 ³²	11	Midazolam AUC +30% ²⁹
									CYP2C8	0.4 ²¹	11	
									CYP2C9	0.264 ^{a,29}	16	No effect on S-warfarin ³⁴
									CYP2D6	2.92 ^{a,29}	1.5	
									CYP2C19	7.64 ^{a,29}	0.56	

								UGT1A1	0.28 ^{a,33}	15	
								UGT1A7	7.8 ^{a,33}	0.55	
								P-gp	17.9 ³⁹	0.36	No effect on digoxin ^{44,45}
								OATP1B1	54% ⁴⁰	–	
								CYP3A4	1 ⁴¹	6.5	No effect on carbamazepine ⁴⁶ , nifedipine ^{47,48} , tacrolimus ⁴⁹
pantoprazole	6.5 ³⁵	98 ³⁵	40 mg SD ³⁵	–	5.5 MTX ³⁶	1.2	Sulfasalazine AUC +83.3% (not significant) with 421A/C genotype ³⁷ , Case report of toxicity with methotrexate ³⁸	CYP2D6	15 ⁴¹	0.43	No effect on metoprolol ⁵⁰
								CYP2C9	13.0 ^{a,42}	0.50	No effect on S-warfarin ⁵¹ , diclofenac ^{52,53}
								CYP2C19	4.1 ⁴³	0.46	No effect on phenytoin ^{54,55} , diazepam ⁵⁶
								CYP2B6	200 ⁴¹	0.03	
								P-gp	0.0156 ⁶⁰	21	Doxorubicin AUC +46% ⁵⁷ , topotecan (IV) AUC +143% ⁵⁹ , paclitaxel AUC increase ⁶³
								OATP1B1	<<20 ⁶¹	–	
elacridar	0.327 ⁵⁷	Not available	400 mg BID x 3D ⁵⁷	–	0.31 MIT ⁵⁸	1.1	Topotecan (IV) AUC +143% ⁵⁹	OATP1B3	>20 ⁸	–	
								OATP2B1	<20 ⁸	–	
								CYP3A4	4.9 ⁶²	0.07	

									CYP2C9	19.1 ⁶²	0.02	
									CYP1A2	16 ⁶²	0.02	
									CYP2C19	2.05 ⁶²	0.16	
									CYP3A4	9.6 ^{a,18}	0.08	No effect on docetaxel ⁶⁶ everolimus ⁶⁷
gefitinib	0.8 ⁶⁴	90 ⁶⁴	225 mg QD x 14D ⁶⁴	–	1.01 E3S ⁶⁵	0.79	No inhibition data		CYP2D6	13.8 ²¹	0.06	Metoprolol AUC +49.5% ⁶⁸
									CYP2C8	17.4 ^{a,18}	0.05	
									CYP2C19	36 ⁶²	0.02	

IC₅₀ and K_i values are presented as reported in the respective references, along with the substrate used: CIM – cimetidine, E3S – estrone-3-sulfate, MTX – methotrexate, MIT – mitoxantrone, TOP – topotecan.

^a – IC₅₀ was calculated as 2 x K_i, assuming linearity

^b – C_{max} for proposed 250 mg dose = 0.54 μM (Bence et al., 2005)

^c – [I₁]/IC₅₀ value for proposed 250 mg dose = 22

¹ STIVARGA[®] product label (05/29/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/203085s001lbl.pdf); ² FDA (2012) Drug approval package: STIVARGA[®] (regorafenib) [FDA application no, (NDA) 203085]; ³ ongoing clinical trial to investigate effect as BCRP inhibitor: <http://clinicaltrials.gov/show/NCT02106845>; ⁴ (Goodman et al., 2001); ⁵ (Elsby et al., 2011); ⁶ (Haagsma et al., 1996); ⁷ (Kusuhara et al., 2012); ⁸ (Karlgrén et al., 2012b); ⁹ TYKERB[®] product label (10/18/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/022059s016s017lbl.pdf); ¹⁰ (Polli et al., 2008); ¹¹ (Molina et al., 2008); ¹² (Midgley et al., 2007); ¹³ (Sugimoto et al., 2011); ¹⁴ FDA (2012) Drug approval package: TYKERB[®] (lapatinib ditosylate) [FDA application no, (NDA) 022059]; ¹⁵ (Yamamoto et al., 2008); ¹⁶ (Noguchi et al., 2009); ¹⁷ (Stewart et al., 2014); ¹⁸ (Wang et al., 2014); ¹⁹ (Li et al., 2010); ²⁰ (Dong et al., 2011); ²¹ (Kim et al., 2013); ²² (Liu et al., 2010); ²³ (Macpherson et al., 2013); ²⁴ (Bullock et al., 2011); ²⁵ (Veeraputhiran and Sundermeyer, 2008); ²⁶ (Tran et al., 2011); ²⁷ (Thomas et al., 2010); ²⁸ (Messersmith et al., 2004); ²⁹ FDA (2012) Drug approval package: TASIGNA[®] (nilotinib hydrochloride monohydrate) [FDA application no, (NDA) 022068]; ³⁰ (Tiwari et al., 2009); ³¹ (Minematsu and Giacomini, 2011); ³² (Kenny et al., 2012); ³³ (Ai et al., 2014); ³⁴ (Yin et al., 2011); ³⁵ PROTONIX[®] product label (12/10/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/020987s048,022020s010lbl.pdf); ³⁶ (Suzuki et al., 2009); ³⁷ (Adkison et al., 2010); ³⁸ (Ranchon et al., 2011); ³⁹ (Pauli-Magnus et al., 2001); ⁴⁰ (Oostendorp et al., 2009); ⁴¹ (Moody et al., 2013); ⁴² (Li et al., 2004); ⁴³ (Zvyaga et al., 2012); ⁴⁴ (Hartmann et al., 1995); ⁴⁵ (Hartmann et al., 1996); ⁴⁶ (Huber et al., 1998); ⁴⁷ (Bliesath et al., 1996b); ⁴⁸ (Bliesath et al., 1996a); ⁴⁹ (Lorf et al., 2000); ⁵⁰ (Koch et al., 1996); ⁵¹ (Duursema et al., 1995); ⁵² (Bliesath et al., 1996d); ⁵³ (Bliesath et al., 1996c); ⁵⁴ (Middle et al., 1995); ⁵⁵ (Middle et al., 1996); ⁵⁶ (Gugler et al., 1996); ⁵⁷ (Planting et al., 2005); ⁵⁸ (Ahmed-Belkacem et al., 2005); ⁵⁹ (Kruijtz et al., 2002); ⁶⁰ (Dickens et al., 2013); ⁶¹ (Karlgrén et al., 2012a); ⁶² (Englund et al., 2014); ⁶³ (Malingre et al., 2001); ⁶⁴ (Goodman et al., 2005); ⁶⁵ (Yanase et al., 2004); ⁶⁶ (Wilding et al., 2006); ⁶⁷ (Milton et al., 2007); ⁶⁸ (Swaisland et al., 2005)

Table 7. In vitro and in vivo data and considerations for potential intestinal BCRP inhibitors.

Precipitant	I ₂ (μM)	Dose	BCRP parameters			Other possible intestinal Interactions				
			K _i (μM)	IC ₅₀ (μM)	I ₂ /IC ₅₀	In vivo PK-based DDI data	Transporter/Enzyme	IC ₅₀ (μM)	I ₂ /IC ₅₀	In vivo PK-based DDI data
lapatinib ^a	5300 ^b	1250 mg QD (SS) ¹	–	0.025 CIM ²	210000 ^c	Decreased topotecan clearance, -18% (renal -22%) ² , SN-38 +45% AUC ⁴	P-gp	1.5 ⁵	3500	Digoxin AUC +2.8-fold ¹ Midazolam AUC +45% (oral) +22% (IV) ¹
							CYP3A	39 ⁶	140	
sulfasalazine ^a	40000	3–4 g SD ⁷	–	0.46 E3S ⁸	87000	No effect with methotrexate ⁹ , however in vivo substrate ¹⁰	OATP2B1	3 ¹¹	13000	
							P-gp	0.8 ¹³	1600	
regorafenib ^a	1300	160 mg (SS) ¹²	–	0.0447 TOP ¹³	28000	No inhibition data ¹⁴	CYP3A4	22.2 ^{d,13}	58	
							UGT1A1	6 ¹³	210	SN-38 glucuronide AUC +44% (when co-administered with irinotecan) ¹³
curcumin ^e	22000	2 g SD ¹⁰	0.7 SSZ ¹⁰	1.6 SSZ ¹⁰	14000	Sulfasalazine exposure increased significantly ¹⁰	P-gp	~20 ¹⁵	1100	Talinolol AUC +33–54% ^{17,18}
							CYP3A4	13.9 ¹⁶	1600	
erlotinib ^a	1500	150 mg QD x 21D ¹⁹	0.15 ²⁰	0.13 E3S ²⁰	12000	No effect on topotecan ²¹	P-gp	2 ²⁰	760	
							OATP2B1	0.55 ¹¹	2800	
							CYP3A4	2.56 ^{d,22}	600	2-fold increase in exposure of OSI-930 ²⁵ , everolimus AUC +21% ²⁶ , case

report of toxicity
with simvastatin ²⁷2-fold increase in
exposure of OSI-
930 ²⁵, everolimus
AUC +21% ²⁶, case
report of toxicity
with simvastatin ²⁷No effect on
irinotecan/SN-38
²⁸Doxorubicin AUC
+46% ²⁹, topotecan
AUC +143% ³¹,
paclitaxel AUC
increase ³⁴

							CYP3A5	80 ^{c,23}	19	
							UGT1A1	1.28 ^{d,24}	1200	
elacridar ^a	2800	400 mg BID x 3D ²⁹	–	0.31 MIT ³⁰	9200	Topotecan AUC +143% ³¹	P-gp	0.0156 ³²	180000	
							OATP2B1	<20 ¹¹	–	
							CYP3A4	4.9 ³³	580	
nilotinib ^a	2800	400 mg BID x 15D ³⁵	0.69 MTX ³⁶	1.38	2000	No inhibition data	CYP3A4	0.4 ³⁷	7100	Midazolam AUC +30% ³⁵
							UGT1A1	0.28 ^{d,38}	10000	
gefitinib ^a	2000	225 mg QD x 14D ³⁹	–	1.01 E3S ⁴⁰	2000	No inhibition data	CYP3A4	9.6 ^{d,22}	210	No effect on docetaxel ⁴¹ everolimus ⁴²
sunitinib	500	50 mg QD x 28D	0.32 E3S ⁴⁴	0.64	780	No inhibition data, in vivo substrate ⁴⁵	P-gp	15.2 ^{d,44}	33	Docetaxel AUC +44.7% ⁴⁷ , Case report of toxicity with colchicine ⁴⁸

43						CYP3A4	19.0 ^{d,46}	26	Docetaxel AUC +44.7% ⁴⁷	
pantoprazole ^a	370	40 mg SD ⁴⁹	–	5.5 MTX ⁵⁰	67	Sulfasalazine AUC +83.3% with 421A/C genotype ⁵¹ ,	P-gp	17.9 ⁵³	21	No effect on digoxin ^{55,56}
						Case report of toxicity with methotrexate ⁵²	CYP3A4	1 ⁵⁴	370	No effect on carbamazepine ⁵⁷ , nifedipine ^{58,59} , tacrolimus ⁶⁰
rabeprazole	440	40 mg AD x 8D ⁶¹	–	8.5 MTX ⁵⁰	52	Delayed methotrexate elimination ⁵⁰	CYP3A4	4.9 ⁵⁴	91	No effect on tacrolimus ⁶²

IC₅₀ and K_i values are presented as reported in the respective references, along with the substrate used: CIM – cimetidine, E3S – estrone-3-sulfate, MTX – methotrexate, MIT – mitoxantrone, SSZ – sulfasalazine, TOP – topotecan.

^a – Also present in [I₁]/IC₅₀ list

^b – I₂ value for proposed 250 mg dose = 1000 μM

^c – [I₂]/IC₅₀ value for proposed 250 mg dose = 40,000

^d – IC₅₀ value was calculated as 2 x K_i, assuming linearity

^e – Curcumin also caused a small, but significant, decrease in the clearance of the CYP1A2 substrate, caffeine (–29%) (Juan et al., 2007)

¹ TYKERB[®] product label (10/18/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/022059s016s017lbl.pdf); ² (Polli et al., 2008); ³ (Molina et al., 2008); ⁴ (Midgley et al., 2007); ⁵ (Sugimoto et al., 2011); ⁶ FDA (2012) Drug approval package: TYKERB[®] (lapatinib ditosylate) [FDA application no, (NDA) 022059]; ⁷ (Goodman et al., 2001); ⁸ (Elsby et al., 2011); ⁹ (Haagsma et al., 1996); ¹⁰ (Kusuhara et al., 2012); ¹¹ (Karlgrén et al., 2012b); ¹² STIVARGA[®] product label (05/29/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/203085s001lbl.pdf); ¹³ FDA (2012) Drug approval package: STIVARGA[®] (regorafenib) [FDA application no, (NDA) 203085]; ¹⁴ ongoing clinical trial to investigate effect as BCRP inhibitor: <http://clinicaltrials.gov/show/NCT02106845>; ¹⁵ (Si et al., 2013); ¹⁶ (Appiah-Opong et al., 2007); ¹⁷ (Juan et al., 2007); ¹⁸ (He et al., 2012); ¹⁹ (Yamamoto et al., 2008); ²⁰ (Noguchi et al., 2009); ²¹ (Stewart et al., 2014); ²² (Wang et al., 2014); ²³ (Li et al., 2010); ²⁴ (Liu et al., 2010); ²⁵ (Macpherson et al., 2013); ²⁶ (Bullock

et al., 2011);²⁷ (Veeraputhiran and Sundermeyer, 2008);²⁸ (Messersmith et al., 2004);²⁹ (Planting et al., 2005);³⁰ (Ahmed-Belkacem et al., 2005);³¹ (Kruijtzer et al., 2002);³² (Dickens et al., 2013);³³ (Englund et al., 2014);³⁴ (Malingre et al., 2001);³⁵ FDA (2012) Drug approval package: TASIGNA[®] (nilotinib hydrochloride monohydrate) [FDA application no, (NDA) 022068];³⁶ (Tiwari et al., 2009);³⁷ (Kenny et al., 2012);³⁸ (Ai et al., 2014);³⁹ (Goodman et al., 2005);⁴⁰ (Yanase et al., 2004);⁴¹ (Wilding et al., 2006);⁴² (Milton et al., 2007);⁴³ SUTENT[®] product label (08/30/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/021938s024s025lbl.pdf);⁴⁴ (Kawahara et al., 2010);⁴⁵ (Oberoi et al., 2013);⁴⁶ FDA (2006) Drug approval package: SUTENT[®] (sunitinib malate) [FDA application no, (NDA) 021938].;⁴⁷ (Robert et al., 2010);⁴⁸ (Abodunde et al., 2013);⁴⁹ PROTONIX[®] product label (12/10/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/020987s048,022020s010lbl.pdf);⁵⁰ (Suzuki et al., 2009);⁵¹ (Adkison et al., 2010);⁵² (Ranchon et al., 2011);⁵³ (Pauli-Magnus et al., 2001);⁵⁴ (Moody et al., 2013);⁵⁵ (Hartmann et al., 1995);⁵⁶ (Hartmann et al., 1996);⁵⁷ (Huber et al., 1998);⁵⁸ (Bliesath et al., 1996b);⁵⁹ (Bliesath et al., 1996a);⁶⁰ (Lorf et al., 2000);⁶¹ ACIPHEX[®] product label (04/19/2013, http://www.accessdata.fda.gov/drugsatfda_docs/label/2013/020973s032lbl.pdf);⁶² (Itagaki et al., 2004)

Figure 1

