

DMD-AR-2020-000316R1

The Long Non-coding RNA HNF4A-AS1 Negatively Regulates Cytochrome P450 Enzymes in Huh7 Cells via Histone Modifications

Pei Wang¹, Shitong Chen¹, Yiting Wang, Xiaofei Wang, Liang Yan, Kun Yang, Xiao-bo Zhong, Shengna Han, and Lirong Zhang

Department of Pharmacology, School of Basic Medical Sciences, Zhengzhou University, Zhengzhou, Henan, China (P.W., S.C., Y.W., X.W., K.Y., S.N., L.Z.); Department of Pharmacy, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan, China (L.Y.); Department of Pharmaceutical Sciences, School of Pharmacy, University of Connecticut, Storrs, Connecticut, USA (X.-b.Z.).

¹P.W. and S.C. contributed equally to this work.

DMD-AR-2020-000316R1

Running Title: HNF4A-AS1 regulates CYPs via histone modifications

Corresponding author: Dr. Lirong Zhang, Department of Pharmacology, School of Basic Medical Sciences, Zhengzhou University, Zhengzhou, Henan, China.

Phone: +86 371-66658807; E-mail: zhanglirongzzu@126.com.

Number of text pages: 38

Number of figures: 6

Number of tables: 2

Number of references: 47

Number of words in Abstract: 213

Number of words in Introduction: 570

Number of words in Discussion: 1329

ABBREVIATIONS:

AHR, aryl hydrocarbon receptor; CAR, constitutive androstane receptor; ChIP, chromatin immunoprecipitation; CYPs, cytochrome P450s; DMEM, dulbecco's modified Eagle's medium; DMSO, dimethyl sulfoxide; FBS, fetal bovine serum; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; HNF1A, hepatocyte nuclear factor 1 alpha; HNF1A-AS1, HNF1A antisense RNA 1; HNF4A,

DMD-AR-2020-000316R1

hepatocyte nuclear factor 4 alpha; HNF4A-AS1, HNF4A antisense RNA 1;
HREs, HNF4A response elements; H3K4me3, histone 3 lysine 4 trimethylation;
IgG, immunoglobulin G; lncRNA, long non-coding RNA; OE-HNF4A-AS1,
overexpression of HNF4A-AS1; PXR, pregnane X receptor; qRT-PCR,
quantitative real-time polymerase chain reaction; RIF, rifampicin; S.D, standard
deviation; siRNAs, small interfering RNAs.

DMD-AR-2020-000316R1

ABSTRACT

The maintenance of homeostasis of cytochrome P450s (CYPs) under both physiological and xenobiotic exposure conditions is ensured by the action of positive and negative regulators. In the current study, the hepatocyte nuclear factor 4 alpha (HNF4A) antisense RNA 1 (HNF4A-AS1), an antisense lncRNA of *HNF4A*, was found to be a negative regulator of the basal and rifampicin (RIF)-induced expression of nuclear receptors and downstream CYPs. In Huh7 cells, knockdown of HNF4A-AS1 resulted in elevated expression of HNF4A, pregnane X receptor (PXR), and CYPs (including CYP3A4) under both basal and RIF-induced conditions. Conversely, overexpression of HNF4A-AS1 led to decreased basal expression of constitutive androstane receptor (CAR), aryl hydrocarbon receptor (AHR), PXR, and all studied CYPs. Of note, significantly-diminished induction levels of PXR, CYP1A2, 2C8, 2C19, and 3A4 by RIF were also observed in HNF4A-AS1 plasmid transfected Huh7 cells. Moreover, the negative feedback of HNF4A on HNF4A-AS1 mediated gene expression was validated using a loss-of-function experiment in this study. Strikingly, our data showed that increased enrichment levels of histone 3 lysine 4 trimethylation (H3K4me3) and HNF4A in the *CYP3A4* promoter contribute to the elevated CYP3A4 expression after HNF4A-AS1 knockdown. Overall, the current study reveals that histone modifications contribute to the negative regulation of nuclear receptors and CYPs by HNF4A-AS1 in basal

DMD-AR-2020-000316R1

and drug-induced levels.

DMD-AR-2020-000316R1

Significance statement

Utilizing loss-of-function and gain-of-function experiments, the current study systematically investigated the negative regulation of HNF4A-AS1 on the expression of nuclear receptors (including HNF4A, CAR, AHR, and PXR) and CYPs (including CYP1A2, 2E1, 2B6, 2D6, 2C8, 2C9, 2C19, and 3A4) in both basal and RIF-induced levels in Huh7 cells. Notably, this study is the first to reveal the contribution of histone modification to the HNF4A-AS1 mediated expression of CYP3A4 in Huh7 cells.

DMD-AR-2020-000316R1

Introduction

Cytochrome P450 (CYP) enzymes, the predominant phase I enzymes in human liver, are responsible for the metabolism of approximately 75% of the clinically used drugs; additionally, CYPs are also of paramount importance for the detoxification of xenobiotics (Pikuleva, 2006; Fisher et al., 2009). Considerable inter-individual variabilities in the expression and induction of CYPs have been observed, which markedly affect the effectiveness and safety of therapeutic agents (Tang et al., 2005; Zanger and Schwab, 2013). Thus, elucidating the precise mechanism underlying the basal and drug-induced expression of CYPs is essential for precision medicine.

It is well studied that nuclear receptors, such as hepatocyte nuclear factor 4 alpha (HNF4A) and 1 alpha (HNF1A), aryl hydrocarbon receptor (AHR), pregnane X receptor (PXR), and constitutive androstane receptor (CAR) are crucial transcriptional regulators of CYPs (Cheung et al., 2003; Kamiyama et al., 2007; Aleksunes and Klaassen, 2012; Hakkola et al., 2018). Notably, HNF4A is considered to be at the top hierarchy of the transcription factor network (Kuo et al., 1992; Kamiya et al., 2003). HNF4A was reported to be involved in HNF1A-mediated transcriptional activation (Eeckhoutte et al., 2004). Additionally, the indispensable role of HNF4A in the PXR and CAR mediated induction of CYPs has also been reported (Kamiya et al., 2003; Tirona et al., 2003; Li and Chiang, 2006; Takezawa et al., 2012). Whereas, these studies

DMD-AR-2020-000316R1

are all focused on positive regulatory mechanisms in CYP expression. Far less information is available on the negative regulation of CYPs, thereby limiting our knowledge on the precise mechanism underlying the maintenance of the homeostasis of CYP expression in normal and xenobiotic (including drug) exposure conditions.

Long non-coding RNAs (lncRNAs) are non-coding RNAs with a length of more than 200 nucleotides (Atkinson et al., 2012; Yang et al., 2014; Ning et al., 2019). Recent studies have reported that lncRNAs participate in the regulation of neighboring genes (Halley et al., 2014; Villegas et al., 2014; Villegas and Zaphiropoulos, 2015). Our previous study has unveiled that HNF4A antisense RNA 1 (HNF4A-AS1), a neighboring antisense lncRNA of *HNF4A*, negatively regulated the mRNA expression of nuclear receptors, including HNF4A and PXR, and the downstream CYPs at the basal level (Chen et al., 2018). However, the underlying mechanism and the role of HNF4A-AS1 in the drug-induced expression of CYPs have not been elucidated thus far.

lncRNAs can act as cofactors for transcriptional factors and jointly regulate gene expression (Zhao et al., 2014; Mi et al., 2017). Additionally, lncRNAs can also regulate gene expression by recruiting complexes to the promoters of target genes and causing alterations of histone modifications of genes (Wang et al., 2011). Of note, studies conducted by us and other groups have shown that transcription factors can affect gene transcription by regulating histone

DMD-AR-2020-000316R1

modifications in gene promoter regions (Xie et al., 2009; Yan et al., 2017).

Therefore, we speculate that HNF4A-AS1 can regulate the expression of CYPs via altering the histone modification status of *CYP* genes, thus affecting the binding of HNF4A.

In the current study, we performed a systematic analysis to investigate the role of HNF4A-AS1 in the expression of transcription factor-mediated CYPs-in Huh7 cells. Importantly, the underlying mechanism was elucidated in this study. Our findings reveal that HNF4A-AS1 negatively regulates the basal and rifampicin (RIF)-induced expression of CYPs. Moreover, the elevated expression of CYP3A4 after HNF4A-AS1 knockdown is related to the increased enrichment levels of histone 3 lysine 4 trimethylation (H3K4me3) and of HNF4A in the promoter of *CYP3A4*.

DMD-AR-2020-000316R1

Materials and Methods

Chemicals and Reagents. Huh7 cells were purchased from the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). RIF was obtained from Sigma-Aldrich (St Louis, MO). Small interfering RNAs (siRNAs), including si-HNF4A, si-HNF4A-AS1, and a negative control (si-NC), were provided by GenePharma Co., Ltd. (Shanghai, China). HNF4A-AS1 expression plasmids were obtained from GeneChem Co., Ltd. (Shanghai, China). Lipofectamine 3000, opti-MEM, and SYBR Select Master Mix were purchased from Thermo Fisher Scientific (Carlsbad, CA). Nanofector transfection reagent was obtained from Procell Co., Ltd. (Wuhan, China). The TriPure isolation reagent and protease inhibitor cocktail were purchased from Roche (Basel, Switzerland). Primers were synthesized by SunYa Co. (Zhejiang, China). Antibodies against HNF4A (ab174653), PXR (bs-2334R), CYP3A4 (ab124921), 1A2 (ab151728), 2C9 (ab150364), and 2C19 (ab185213) were obtained from Abcam (Cambridge, MA). Horseradish peroxidase-linked secondary antibodies (SA00001-1 and SA00001-2) and the antibody against glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (60004-1-Ig) were purchased from Proteintech (Wuhan, China). The antibody recognizing H3K4me3 (17-678) and polyvinylidene fluoride (PVDF) membranes were obtained from MD Millipore (Billerica, MA). Dulbecco's modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were purchased from Corning

DMD-AR-2020-000316R1

Inc. (Armonk, NY). RIPA reagent, penicillin and streptomycin mixture, dimethyl sulfoxide (DMSO), and other chemical reagents were provided by Solarbio Science & Technology Co. (Beijing, China).

Cell Culture and Transfection. Huh7 cells were cultured in DMEM medium supplemented with 10% FBS and 1% penicillin and streptomycin mixture at 37°C with 5% CO₂. Before transfection, Huh7 cells were allowed to grow for up to 24 hours on six-well plates reaching 80%–90% confluence. For gene knockdown experiments, 40 pmol/well of siRNAs targeting HNF4A, HNF4A-AS1, or control siRNAs were transfected into Huh7 cells with the Lipofectamine 3000 reagent according to the manufacturer's instructions. For gene overexpression experiments, Nanofecter transfection reagent was used to transfect 2.5 µg/well of plasmids into Huh7 cells following the manufacturer's protocol. Cells were harvested 48 hours after transfection.

Drug Treatment. Twenty-four hours after transfection, cells were incubated with 10 µM RIF or 0.1% DMSO (vehicle) for 24 hours before harvested.

RNA Isolation and Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR). Total RNAs were isolated from Huh7 cells using a TriPure isolation reagent according to the manufacturer's instruction. The quality and concentrations of RNAs were measured by a Nanodrop 2000c Spectrophotometer (Thermo Fisher Scientific). One microgram total RNAs were reverse transcribed using a PrimeScript RT reagent kit. qRT-PCR was

DMD-AR-2020-000316R1

performed using a SYBR Select Master Mix with specific primers (Table 1) in an Applied Biosystems 7500 Fast Real-Time PCR System (Thermo Fisher Scientific). Relative RNA expression levels were determined by normalizing focused gene expression against GAPDH expression using the $2^{-\Delta\Delta C_t}$ method.

Protein Isolation and Western Blot Analysis. Whole protein homogenates of Huh7 cells were prepared using a RIPA reagent supplemented with a protease inhibitor cocktail. Protein concentrations were determined by a bicinchoninic acid method according to the manufacturer's instructions (Beyotime Institute of Biotechnology, Hangzhou, China). Western blot analysis was performed using 80 µg total protein samples, as described previously (Yan et al., 2017; Wang et al., 2019a) with minor modifications. The dilutions of primary antibodies against HNF4A, PXR, and CYP3A4 were 1:1,000. The dilutions of primary antibodies against CYP1A2, 2C9, and 2C19 were 1:2,000. The dilutions of the primary antibody against GAPDH was 1:5,000. Horseradish peroxidase-linked secondary antibodies were diluted in 1:10,000. Protein bands were visualized using an enhanced chemiluminescence method with a FluorChem E system (ProteinSample, San Jose, California). The protein expression levels of focused genes were normalized against GAPDH.

Chromatin Immunoprecipitation (ChIP) Analysis. ChIP was performed as described previously (Yan et al., 2017; Wang et al., 2019a) with minor

DMD-AR-2020-000316R1

modifications. Briefly, siRNA-transfected cells were incubated with 1% formaldehyde for 10 minutes; fixation was halted by glycine (125 mM) for 5 minutes at room temperature. Cells were then washed twice with cold phosphate-buffered saline and lysed using an SDS lysis buffer (0.1% SDS, 50 mM Tris-HCl, pH 8.0, 1 mM EDTA, and 1% Triton-X) with a proteinase inhibitor cocktail on ice for 15 minutes. Cell lysates were sonicated with a Bioruptor Pico sonication system (Diagenode, Seraing, Belgium) to obtain DNA fragments ranging between ~200 and 1,000 bp. DNA fragments were immunoprecipitated using antibodies against H3K4me3 and HNF4A. Mouse immunoglobulin G (IgG) was used as a negative control. The purification of DNA was performed with a PCR purification kit (QIAGEN GmbH, Hilden, Germany) followed by qPCR analysis with specific primers (Table 2) using a SYBR green method. Enrichment was calculated as the percentage of chromatin input and normalized against IgG.

Statistical Analysis. All studies with Huh7 cells described here were performed as three independent experiments. Data are shown as mean \pm standard deviation (S.D). Statistical significances between two groups were analyzed by two-tailed unpaired Student's *t* test using SPSS 21.0 software (IBM Corp., Armonk, NY).

DMD-AR-2020-000316R1

Results

Negative Regulation of HNF4A-AS1 on the Basal Expression of Nuclear Receptors and CYPs in Huh7 Cells.

To uncover the role of HNF4A-AS1 in the regulation of nuclear receptors and CYPs, loss-of-function and gain-of-function studies were performed in Huh7 cells. The expression of HNF4A-AS1, five nuclear receptors (HNF4A, HNF1A, CAR, AHR, and PXR), and eight CYPs (CYP1A2, 2E1, 2B6, 2D6, 2C8, 2C9, 2C19, and 3A4) was measured.

The knockdown efficiency of HNF4A-AS1 after siRNA transfection into Huh7 cells was confirmed by qRT-PCR. As seen in Fig.1A, the expression of HNF4A-AS1 decreased approximately 57% after si-HNF4A-AS1 transfection. Knockdown of HNF4A-AS1 led to increased expression of HNF4A at both mRNA and protein levels (Fig. 1A and B). For the focused nuclear receptors, 1.7-fold increased expression of PXR was observed by HNF4A-AS1 knockdown, while the basal expression of HNF1A, CAR, and AHR was unchanged at the mRNA level (Fig. 1C). Meanwhile, the basal mRNA expression of CYP1A2, 2C8, 2C9, 2C19, and 3A4 was also significantly higher in the HNF4A-AS1 knockdown group than that in the control group (Fig. 1D). There was no significant difference in the mRNA expression of CYP2E1, 2B6, and 2D6 after HNF4A-AS1 knockdown (Fig. 1D). Importantly, consistent with the increased expression of PXR and CYP3A4 at the mRNA level, the protein

DMD-AR-2020-000316R1

expression of PXR and CYP3A4 was also significantly elevated after HNF4A-AS1 knockdown (Fig. 1E and F). Moreover, higher protein expression levels of CYP1A2 and 2C9 were also observed in the HNF4A-AS1 group ($P<0.05$), while the protein expression of CYP2C19 remained unchanged ($P>0.05$) (supplemental Fig. S1).

The expression of HNF4A-AS1 in the overexpression of HNF4A-AS1 (OE-HNF4A-AS1) group was approximately 673 times that in the control group (Fig. 2A), which confirmed the overexpression efficiency of HNF4A-AS1. Compared with the control group, the mRNA expression of CAR, AHR, and PXR was approximately 50% lower in the HNF4A-AS1 overexpression group (Fig. 2B). However, the mRNA expression of HNF4A and HNF1A was not affected by HNF4A-AS1 overexpression ($P>0.05$) (Fig. 2A and B). Additionally, as shown in Fig. 2C, HNF4A-AS1 overexpression also resulted in significantly decreased expression of studied CYPs except for CYP2D6 at the mRNA level. These findings indicate that HNF4A-AS1 acts as a negative regulator in the basal expression of CYPs, probably, via the negative regulation of nuclear receptors.

HNF4A-AS1 Attenuates the RIF-induced Expression of CYPs in Huh7 Cells.

To further explore the role of HNF4A-AS1 in the drug-induced expression of CYPs, siRNA or plasmid transfected Huh7 cells were treated with RIF. As seen

DMD-AR-2020-000316R1

in Fig. 3A, knockdown of HNF4A-AS1 abolished the induction of HNF4A-AS1 expression by RIF, whereas elevated that of HNF4A. Significantly higher induction levels of PXR were also observed in the HNF4A-AS1 knockdown group ($P < 0.05$), whereas the RIF-induced expression of other nuclear receptors (including HNF1A, CAR, and PXR) was unaffected by the HNF4A-AS1 knockdown ($P > 0.05$) (Fig. 3B). Additionally, after HNF4A-AS1 knockdown, the induction of CYP1A2, 2C8, 2C9, 2C19, and 3A4 expression by RIF was significantly increased ($P < 0.05$), while that of CYP2E1, 2B6, and 2D6 was not affected ($P > 0.05$) (Fig. 3C).

Conversely, overexpression of HNF4A-AS1 led to higher induction levels of HNF4A-AS1 and lower that of PXR by RIF (Fig. 3D and E). However, the RIF-induced expression of HNF4A, HNF1A, CAR, and AHR remained unchanged after HNF4A-AS1 overexpression ($P > 0.05$) (Fig. 3D and E). For CYPs, the induction levels of CYP1A2, 2C8, 2C19, and 3A4 decreased after HNF4A-AS1 overexpression (Fig. 3F). Altogether, these results suggest that HNF4A-AS1 attenuates the drug-induced expression of CYPs.

Negative feedback of HNF4A on HNF4A-AS1 mediated genes expression.

To verify the role of HNF4A in the transcriptional regulation of HNF4A-AS1 and the related CYPs, knockdown of HNF4A was performed in Huh7 cells. After effective knockdown of HNF4A, the expression of HNF4A-AS1 decreased (Fig. 4A and B). Lower expression levels of all studied nuclear

DMD-AR-2020-000316R1

receptors, including HNF1A, CAR, AHR, and PXR, were also observed at the mRNA level after HNF4A knockdown (Fig. 4C). Additionally, the knockdown of HNF4A also led to reduced mRNA expression of CYPs (Fig. 4D). Consistent with the decreased expression at the mRNA level, the protein expression of PXR and CYP3A4 was lower in the HNF4A knockdown group than that in the control group (Fig. 4E and F). Moreover, as seen in Fig. 5, the RIF-induced expression of most focused genes, including HNF1A and CYPs, was significantly diminished after HNF4A knockdown at the RNA level. There was also a trend of lower induced expression levels of CAR, AHR, and PXR by RIF in HNF4A knockdown *versus* control cells ($P>0.05$) (Fig. 5B). These findings demonstrate that HNF4A participates in the transcriptional regulation of HNF4A-AS1, nuclear receptors, and CYPs under both basal and RIF-induced expression-levels conditions.

HNF4A-AS1 Knockdown leads to Elevated Enrichment Levels of Histone Modifications and HNF4A in the Promoter of *CYP3A4*.

To explore the underlying mechanism by which HNF4A-AS1 regulated the expression of CYP3A4, ChIP-qPCR experiments were conducted in si-NC or si-HNF4A-AS1 transfected Huh7 cells. As shown in Fig. 6A, two pairs of primers around the HNF4A response elements (HREs) in the promoter of *CYP3A4* were synthesized. Knockdown of HNF4A-AS1 resulted in elevated enrichment levels of H3K4me3, a gene activate mark, in the HREs of the

DMD-AR-2020-000316R1

CYP3A4 promotor, which was consistent with the higher expression of *CYP3A4* in the HNF4A-AS1 knockdown group (Fig. 6B). Moreover, the enrichment level of HNF4A also increased after HNF4A-AS1 knockdown (Fig. 6C). Altogether, these findings indicate that HNF4A-AS1 might regulate the expression of *CYP3A4* by a scaffold mechanism.

DMD-AR-2020-000316R1

Discussion

It is well known that the expression of CYPs is transcriptionally regulated by nuclear receptors (Honkakoski and Negishi, 2000; Jover et al., 2009; Nie et al., 2017). As a master regulator, HNF4A plays a curial role in the complex regulatory network of basal and drug-induced CYP expression (Tirona et al., 2003; Kamiyama et al., 2007). Recently, an increasing number of studies have elucidated the epigenetic regulation of CYPs (Klaassen et al., 2011; Tang and Chen, 2015; Li et al., 2019). Notably, as promising epigenetic regulators, lncRNAs were reported to be important for multiple biological or physiological processes (Fatica and Bozzoni, 2014; Schmitt and Chang, 2016; Ghafouri-Fard et al., 2020; Zhou et al., 2020). In the current study, utilizing loss-of-function and gain-of function experiments, we systemically verified that HNF4A-AS1 was involved in the negative regulation of HNF4A-mediated gene expression (including that of CYPs) at the basal level and under drug-induced conditions in Huh7 cells. Importantly, the alteration of histone modification status was proved as the underlying mechanism for the HNF4A-AS1-mediated negative-regulation of CYPs in this study.

Mounting evidence suggests that lncRNAs can be involved in the regulation of their neighboring genes (Villegas and Zaphiropoulos, 2015; Gibbons et al., 2018; Qi et al., 2019). Particularly, lncRNAs are known to interact on their antisense strands with transcription factors (Villegas and Zaphiropoulos, 2015;

DMD-AR-2020-000316R1

Gil and Ulitsky, 2020). Our previous studies proved that *HNF1A* antisense lncRNA, HNF1A-AS1, played as a positive regulator of HNF1A-mediated gene expression in Huh7 and HepaRG cells (Chen et al., 2018; Wang et al., 2019b). In the present study, using loss-of-function and gain-of-function studies, HNF4A-AS1 was validated to have opposite regulatory effects to HNF1A-AS1, on the basal expression of most studied genes in Huh7 cells (Fig. 1 and 2), which confirmed our previous findings in HNF4A-AS1-knockdown HepaRG cells (Chen et al., 2018).

Though higher mRNA expression of most CYPs (including CYP1A2, 2C9, 2C19, and 3A4) after HNF4A-AS1 knockdown was observed in both Huh7 and HepaRG cells (Fig. 1)(Chen et al., 2018), differential impacts on genes were noticed. For instance, after efficient knockdown of HNF4A-AS1, the basal mRNA expression of HNF4A and PXR was significantly elevated in Huh7 cells (Fig. 1A and C) but remained unchanged in HepaRG cells (Chen et al., 2018). Increased protein expression of HNF4A and PXR by HNF4A-AS1 knockdown was also observed in Huh7 cells (Fig. 1B and E). However, overexpression of HNF4A-AS1 in Huh7 cells did not affect the mRNA expression of HNF4A (Fig. 2A). This may be attributed to the low intrinsic expression of HNF4A in Huh7 cells. Another possible explanation for the above-mentioned findings is that HNF4A-AS1 may serve as a central platform for the binding of effector molecules repressing the expression of HNF4A in Huh7 cells. That is, the

DMD-AR-2020-000316R1

knockdown of HNF4A-AS1 may lead to dismantling of the lncRNA-effector scaffold, thus increasing the expression of HNF4A, whereas overexpression of HNF4A-AS1 may have no impact on the scaffolding complexes, and consequently on the expression of HNF4A. However, these are aspects that warrant further investigation. Additionally, transfection of siRNA against HNF4A-AS1 into HepaRG cells resulted in the altered mRNA expression of CAR and AHR (Chen et al., 2018); whereas, in Huh7 cells, the expression of CAR and PXR was not affected after HNF4A-AS1 knockdown, although it decreased after overexpression of HNF4A-AS1 (Fig. 1C and 2B). The differential impacts of HNF4A-AS1 on the basal expression of genes in Huh7 and HepaRG cells may be ascribed to differences in the intrinsic expression of HNF4A-AS1, HNF4A-related nuclear receptors, and CYPs. Overall, both studies confirmed the negative regulation of HNF4A-AS1 in the basal and drug-induced expression of HNF4A-mediated genes.

It is well documented that PXR and CAR are xenobiotic receptors that play curial roles in the induction or inhibition of CYPs (Timsit and Negishi, 2007; Hernandez et al., 2009; Chai et al., 2013). Additionally, several studies have reported that HNF4A is indispensable for the PXR- and CAR-mediated induction of CYPs (Tirona et al., 2003; Li and Chiang, 2006; Takezawa et al., 2012). Based on our previous findings, we speculated that HNF4A-AS1 could be involved in the drug-induced expression of CYPs. Importantly, consistent

DMD-AR-2020-000316R1

with the elevated basal expression of HNF4A, PXR, and CYPs (such as CYP1A2 and 3A4) in HNF4A-AS1 knockdown Huh7 cells, the induction of these genes by RIF was improved after HNF4A-AS1 knockdown (Fig. 3A-C). Conversely, overexpression of HNF4A-AS1 resulted in diminished induction levels of PXR, CYP1A2, 2C8, 2C19, and 3A4 in Huh7 cells (Fig. 3D-F). These results indicate that HNF4A-AS1 is involved in the negative regulation of drug-induced CYP expression. Under both normal physical and xenobiotic exposure conditions, HNF4A-AS1 may act as a “break” to sustain constant intracellular levels of CYPs. Importantly, to better understand the human implications of these findings, studies using primary human hepatocytes, human induced pluripotent stem cell-derived hepatocytes, or HepaRG cells should be conducted in the future.

In a recent study, using dual-luciferase assay, Guo S and Lu H found that the endogenous expression of HNF4A-AS1 in HepG2 and HEK293 cells was transcriptionally regulated by P1-HNF4A, the predominant subtype of HNF4A in adult liver (Guo and Lu, 2019). Thus, in this study, we also assessed the impacts of HNF4A on the expression of HNF4A-AS1, other nuclear receptors, and CYPs under basal and RIF-induced conditions. As shown in Fig. 4, knockdown of HNF4A led to decreased expression of HNF4A-AS1, nuclear receptors (HNF1A, CAR, and PXR), and CYPs (including CYP2E1, 2B6, 2C8, 2C19, and 3A4) in Huh7 cells, which was consistent with our previous findings

DMD-AR-2020-000316R1

in HepaRG cells. Similarly, the RIF-induced expression of these CYPs was also diminished after HNF4A knockdown in both Huh7 (Fig. 5) and HepaRG cells (Chen et al., 2018). These findings verified the previously published results and further indicated that there was a negative feedback loop in the HNF4A-AS1/HNF4A-CYPs axis. Whereas, the precise mechanisms need to be elucidated in further researches.

The involvement of histone modifications in the regulation of CYPs has been proved by more and more studies (Yan et al., 2017; Wang et al., 2019a; Pande et al., 2020). For instance, H3K27ace was reported to be involved in HNF4A-mediated CYP2C9 expression. Englert N and colleagues found that HNF4A could recruit the co-regulator MED25 to the *CYP2C9* promoter, thus promoting the transcription of CYP2C9 (Englert et al., 2015). However, alterations in histone modification status are usually non-spontaneous, and their activation requires the action of upstream regulatory factors. Therefore, we speculate that HNF4A-AS1, playing as a cofactor, would interact with HNF4A by recruiting histone modification enzymes to the HREs in the *CYP3A4* promoter, thereby regulating the expression of CYPs. Given the highest expression level among the entire CYPs in liver, CYP3A4 was selected as a representative gene to uncover the underlying mechanisms that how HNF4A-AS1 regulated the expression of CYPs in this study. After the efficient knockdown of HNF4A-AS1, the HREs (-7,836 to -6,093 bp and -362 to +53 bp)

DMD-AR-2020-000316R1

in the *CYP3A4* promoter were found to have higher enrichment levels of H3K4me3, a gene-active factor. In addition, interference with HNF4A-AS1 resulted in elevated enrichment levels of HNF4A in the promoter of *CYP3A4*. Altogether, these findings validate our hypothesis and indicate that HNF4A-AS1 may regulate the expression of *CYP3A4* by recruiting histone methylation enzymes to the promoter of *CYP3A4*, altering the histone modification status, thereby affecting the binding of HNF4A to *CYP3A4* promoter. However, whether HNF4A-AS1 interacts with HNF4A and histone modification enzymes was not elucidated in this study. This said, these findings provide potential future directions for the investigation of the molecular mechanisms of HNF4A-AS1 in HNF4A-mediated regulatory functions in the human liver.

In summary, this study observed a negative regulation of HNF4A-AS1 on the basal and drug-induced expression of HNF4A, CYPs, and other nuclear receptors. Our study also revealed that elevated enrichment levels of H3K4me3 and HNF4A in the *CYP3A4* promotor was responsible for the induced *CYP3A4* expression after HNF4A-AS1 knockdown. Therefore, the HNF4A-AS1/HNF4A-CYPs axis is an epigenetic-dependent novel negative feedback regulatory loop that should be considered for maintenance of the homeostasis of CYPs in the liver.

DMD-AR-2020-000316R1

Authorship Contributions

Participated in research design: P. Wang, Yan, Zhong, Han, Zhang

Conducted experiments: P. Wang, Chen, Y. Wang, X. Wang, Yan, Yang

Performed data analysis: P. Wang, Chen, Zhang

Wrote or contributed to the writing of the manuscript: P. Wang, Chen, Zhong,

Zhang

DMD-AR-2020-000316R1

References

- Aleksunes LM and Klaassen CD (2012) Coordinated regulation of hepatic phase I and II drug-metabolizing genes and transporters using AhR-, CAR-, PXR-, PPARalpha-, and Nrf2-null mice. *Drug metabolism and disposition: the biological fate of chemicals* **40**:1366-1379.
- Atkinson SR, Marguerat S, and Bähler J (2012) Exploring long non-coding RNAs through sequencing. *Seminars in cell & developmental biology* **23**:200-205.
- Chai X, Zeng S, and Xie W (2013) Nuclear receptors PXR and CAR: implications for drug metabolism regulation, pharmacogenomics and beyond. *Expert Opinion on Drug Metabolism & Toxicology* **9**:253-266.
- Chen L, Bao Y, Piekos SC, Zhu K, Zhang L, and Zhong XB (2018) A Transcriptional Regulatory Network Containing Nuclear Receptors and Long Noncoding RNAs Controls Basal and Drug-Induced Expression of Cytochrome P450s in HepaRG Cells. *Mol Pharmacol* **94**:749-759.
- Cheung C, Akiyama TE, Kudo G, and Gonzalez FJ (2003) Hepatic expression of cytochrome P450s in hepatocyte nuclear factor 1-alpha (HNF1alpha)-deficient mice. *Biochemical pharmacology* **66**:2011-2020.
- Eeckhoutte J, Formstecher P, and Laine B (2004) Hepatocyte nuclear factor 4alpha enhances the hepatocyte nuclear factor 1alpha-mediated activation of transcription. *Nucleic acids research* **32**:2586-2593.

DMD-AR-2020-000316R1

- Englert NA, Luo G, Goldstein JA, and Surapureddi S (2015) Epigenetic Modification of Histone 3 Lysine 27 MEDIATOR SUBUNIT MED25 IS REQUIRED FOR THE DISSOCIATION OF POLYCOMB REPRESSIVE COMPLEX 2 FROM THE PROMOTER OF CYTOCHROME P450 2C9. *J Biol Chem* **290**:2264-2278.
- Fatica A and Bozzoni I (2014) Long non-coding RNAs: new players in cell differentiation and development. *Nat Rev Genet* **15**:7-21.
- Fisher CD, Lickteig AJ, Augustine LM, Ranger-Moore J, Jackson JP, Ferguson SS, and Cherrington NJ (2009) Hepatic cytochrome P450 enzyme alterations in humans with progressive stages of nonalcoholic fatty liver disease. *Drug Metab Dispos* **37**:2087-2094.
- Ghafouri-Fard S, Shoorei H, Mohaqiq M, and Taheri M (2020) Non-coding RNAs regulate angiogenic processes. *Vascular Pharmacology* **133**.
- Gibbons HR, Shaginurova G, Kim LC, Chapman N, Spurlock CF, III, and Aune TM (2018) Divergent lncRNA GATA3-AS1 Regulates GATA3 Transcription in T-Helper 2 Cells. *Frontiers in Immunology* **9**.
- Gil N and Ulitsky I (2020) Regulation of gene expression by cis-acting long non-coding RNAs. *Nat Rev Genet* **21**:102-117.
- Guo S and Lu H (2019) Novel mechanisms of regulation of the expression and transcriptional activity of hepatocyte nuclear factor 4alpha. *J Cell Biochem* **120**:519-532.

DMD-AR-2020-000316R1

- Hakkola J, Bernasconi C, Coecke S, Richert L, Andersson TB, and Pelkonen O (2018) Cytochrome P450 Induction and Xeno-Sensing Receptors Pregnane X Receptor, Constitutive Androstane Receptor, Aryl Hydrocarbon Receptor and Peroxisome Proliferator-Activated Receptor α at the Crossroads of Toxicokinetics and Toxicodynamics. *Basic & clinical pharmacology & toxicology* **123 Suppl 5**:42-50.
- Halley P, Kadakkuzha BM, Faghihi MA, Magistri M, Zeier Z, Khorkova O, Coito C, Hsiao J, Lawrence M, and Wahlestedt C (2014) Regulation of the apolipoprotein gene cluster by a long noncoding RNA. *Cell reports* **6**:222-230.
- Hernandez JP, Mota LC, and Baldwin WS (2009) Activation of CAR and PXR by Dietary, Environmental and Occupational Chemicals Alters Drug Metabolism, Intermediary Metabolism, and Cell Proliferation. *Current pharmacogenomics and personalized medicine* **7**:81-105.
- Honkakoski P and Negishi M (2000) Regulation of cytochrome P450 (CYP) genes by nuclear receptors. *The Biochemical journal* **347**:321-337.
- Jover R, Moya M, and Jose Gomez-Lechon M (2009) Transcriptional Regulation of Cytochrome P450 Genes by the Nuclear Receptor Hepatocyte Nuclear Factor 4-Alpha. *Current Drug Metabolism* **10**:508-519.
- Kamiya A, Inoue Y, and Gonzalez FJ (2003) Role of the hepatocyte nuclear factor 4alpha in control of the pregnane X receptor during fetal liver

DMD-AR-2020-000316R1

development. *Hepatology* **37**:1375-1384.

Kamiyama Y, Matsubara T, Yoshinari K, Nagata K, Kamimura H, and Yamazoe Y (2007) Role of human hepatocyte nuclear factor 4 alpha in the expression of drug-metabolizing enzymes and transporters in human Hepatocytes assessed by use of small interfering RNA. *Drug Metabolism and Pharmacokinetics* **22**:287-298.

Klaassen CD, Lu H, and Cui JY (2011) Epigenetic regulation of drug processing genes. *Toxicology Mechanisms and Methods* **21**:312-324.

Kuo CJ, Conley PB, Chen L, Sladek FM, Darnell JE, Jr., and Crabtree GR (1992) A transcriptional hierarchy involved in mammalian cell-type specification. *Nature* **355**:457-461.

Li DY, Tolleson WH, Yu DK, Chen S, Guo L, Xiao WM, Tong WD, and Ning BT (2019) Regulation of cytochrome P450 expression by microRNAs and long noncoding RNAs: Epigenetic mechanisms in environmental toxicology and carcinogenesis. *J Environ Sci Health Pt C-Environ Carcinog Ecotoxicol Rev* **37**:180-214.

Li T and Chiang JY (2006) Rifampicin induction of CYP3A4 requires pregnane X receptor cross talk with hepatocyte nuclear factor 4alpha and coactivators, and suppression of small heterodimer partner gene expression. *Drug Metab Dispos* **34**:756-764.

Mi L, Zhao XY, Li S, Yang G, and Lin JD (2017) Conserved function of the long

DMD-AR-2020-000316R1

noncoding RNA Blnc1 in brown adipocyte differentiation. *Molecular metabolism* **6**:101-110.

Nie YL, He H, Li JF, Meng XG, Yan L, Wang P, Wang SJ, Bi HZ, Zhang LR, and Kan QC (2017) Hepatic expression of transcription factors affecting developmental regulation of UGT1A1 in the Han Chinese population. *Eur J Clin Pharmacol* **73**:29-37.

Ning B, Yu D, and Yu AM (2019) Advances and challenges in studying noncoding RNA regulation of drug metabolism and development of RNA therapeutics. *Biochem Pharmacol* **169**:113638.

Pande P, Zhong X-B, and Ku WW (2020) Histone Methyltransferase G9a Regulates Expression of Nuclear Receptors and Cytochrome P450 Enzymes in HepaRG Cells at Basal Level and in Fatty Acid Induced Steatosis. *Drug metabolism and disposition: the biological fate of chemicals*.

Pikuleva IA (2006) Cholesterol-metabolizing cytochromes P450. *Drug Metab Dispos* **34**:513-520.

Qi W, Li Z, Xia L, Dai J, Zhang Q, Wu C, and Xu S (2019) LncRNA GABPB1-AS1 and GABPB1 regulate oxidative stress during erastin-induced ferroptosis in HepG2 hepatocellular carcinoma cells. *Scientific Reports* **9**.

Schmitt AM and Chang HY (2016) Long Noncoding RNAs in Cancer Pathways. *Cancer Cell* **29**:452-463.

Takezawa T, Matsunaga T, Aikawa K, Nakamura K, and Ohmori S (2012)

DMD-AR-2020-000316R1

Lower expression of HNF4alpha and PGC1alpha might impair rifampicin-mediated CYP3A4 induction under conditions where PXR is overexpressed in human fetal liver cells. *Drug Metab Pharmacokinet* **27**:430-438.

Tang C, Lin JH, and Lu AYH (2005) Metabolism-based drug-drug interactions: What determines individual variability in cytochrome P450 induction? *Drug Metabolism and Disposition* **33**:603-613.

Tang X and Chen S (2015) Epigenetic Regulation of Cytochrome P450 Enzymes and Clinical Implication. *Current Drug Metabolism* **16**:86-96.

Timsit YE and Negishi M (2007) CAR and PXR: The xenobiotic-sensing receptors. *Steroids* **72**:231-246.

Tirona RG, Lee W, Leake BF, Lan LB, Cline CB, Lamba V, Parviz F, Duncan SA, Inoue Y, Gonzalez FJ, Schuetz EG, and Kim RB (2003) The orphan nuclear receptor HNF4alpha determines PXR- and CAR-mediated xenobiotic induction of CYP3A4. *Nat Med* **9**:220-224.

Villegas VE, Rahman MF, Fernandez-Barrena MG, Diao Y, Liapi E, Sonkoly E, Ståhle M, Pivarcsi A, Annaratone L, Sapino A, Ramírez Clavijo S, Bürglin TR, Shimokawa T, Ramachandran S, Kapranov P, Fernandez-Zapico ME, and Zaphiropoulos PG (2014) Identification of novel non-coding RNA-based negative feedback regulating the expression of the oncogenic transcription factor GLI1. *Molecular oncology* **8**:912-926.

DMD-AR-2020-000316R1

- Villegas VE and Zaphiropoulos PG (2015) Neighboring gene regulation by antisense long non-coding RNAs. *International journal of molecular sciences* **16**:3251-3266.
- Wang KC, Yang YW, Liu B, Sanyal A, Corces-Zimmerman R, Chen Y, Lajoie BR, Protacio A, Flynn RA, Gupta RA, Wysocka J, Lei M, Dekker J, Helms JA, and Chang HY (2011) A long noncoding RNA maintains active chromatin to coordinate homeotic gene expression. *Nature* **472**:120-124.
- Wang P, Liu G, Nie Y, Han S, Li J, Zhong XB, and Zhang L (2019a) Epigenetic Memory Is Involved in the Persistent Alterations of Drug-Processing Genes in Adult Mice due to PCN-Activated PXR during Early Life. *Toxicol Sci*.
- Wang Y, Yan L, Liu J, Chen S, Liu G, Nie Y, Wang P, Yang W, Chen L, Zhong X, Han S, and Zhang L (2019b) The HNF1alpha-Regulated LncRNA HNF1alpha-AS1 Is Involved in the Regulation of Cytochrome P450 Expression in Human Liver Tissues and Huh7 Cells. *J Pharmacol Exp Ther* **368**:353-362.
- Xie Y, Ke S, Ouyang N, He J, Xie W, Bedford MT, and Tian Y (2009) Epigenetic regulation of transcriptional activity of pregnane X receptor by protein arginine methyltransferase 1. *The Journal of biological chemistry* **284**:9199-9205.
- Yan L, Wang YT, Liu JY, Nie YL, Zhong XB, Kan QC, and Zhang LR (2017) Alterations of Histone Modifications Contribute to Pregnane X

DMD-AR-2020-000316R1

Receptor-Mediated Induction of CYP3A4 by Rifampicin. *Mol Pharmacol* **92**:113-123.

Yang L, Froberg JE, and Lee JT (2014) Long noncoding RNAs: fresh perspectives into the RNA world. *Trends in biochemical sciences* **39**:35-43.

Zanger UM and Schwab M (2013) Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation. *Pharmacology & therapeutics* **138**:103-141.

Zhao XY, Li S, Wang GX, Yu Q, and Lin JD (2014) A long noncoding RNA transcriptional regulatory circuit drives thermogenic adipocyte differentiation. *Molecular cell* **55**:372-382.

Zhou Y, Sun W, Qin Z, Guo S, Kang Y, Zeng S, and Yu L (2020) LncRNA regulation: New frontiers in epigenetic solutions to drug chemoresistance. *Biochemical pharmacology*:114228-114228.

DMD-AR-2020-000316R1

Footnotes

This work was supported by the National Natural Science Foundation of China [Grants 81773815, 82073931, and U1604163]; and the National Key Research and Development Project [Grant 2018YFA0107303]. The authors declare that they have no conflict of interest.

DMD-AR-2020-000316R1

Figure legends

Fig. 1. Impact of HNF4A-AS1 knockdown on the basal expression of genes in Huh7 cells. si-HNF4A-AS1 or negative control siRNA was transfected into Huh7 cells for 24 hours. The RNA and protein expression levels of genes after HNF4A-AS1 knockdown were measured by qRT-PCR and Western Blot, respectively. (A) Expression of HNF4A and HNF4A-AS1 at the RNA level. (B) The protein expression of HNF4A. (C) The relative mRNA expression of selected nuclear receptors. (D) The relative mRNA expression of CYPs. (E) The protein expression of PXR. (F) The protein expression of CYP3A4. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. **P* < 0.05; ***P* < 0.01 vs. the si-NC group.

Fig. 2. Impact of HNF4A-AS1 overexpression on the basal expression of genes in Huh7 cells. HNF4A-AS1 expression plasmid or negative control plasmid was transfected into Huh7 cells for 24 hours. The RNA expression of genes after HNF4A-AS1 overexpression was measured by qRT-PCR. (A) Expression of HNF4A and HNF4A-AS1 at the RNA level. (B) The relative mRNA expression of selected nuclear receptors. (C) The relative mRNA expression of CYPs. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001 vs. the OE-NC group. OE-HNF4A-AS1,

DMD-AR-2020-000316R1

overexpression of HNF4A-AS1.

Fig. 3. HNF4A-AS1 repressed the RIF-induced expression of genes in Huh7 cells. si-HNF4A-AS1, si-NC, OE-HNF4A-AS1, or OE-NC transfected Huh7 cells were treated with RIF for 24 hours. The RIF-induced expression of genes after HNF4A-AS1 knockdown or overexpression at the RNA level was detected by qRT-PCR method. (A-C) Knockdown of HNF4A-AS1 elevated the RIF-induced expression of genes. (D-F) Overexpression of HNF4A-AS1 diminished the RIF-induced expression of genes. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. **P* < 0.05; ***P* < 0.01; ****P* < 0.001 vs. the control group.

Fig. 4. Impact of HNF4A knockdown on the basal expression of genes in Huh7 cells. si-HNF4A or negative control siRNA was transfected into Huh7 cells for 24 hours. The RNA and protein expression of selected genes after HNF4A knockdown were measured by qRT-PCR and Western Blot method, respectively. (A) Expression of HNF4A and HNF4A-AS1 at the RNA level. (B) The protein expression of HNF4A. (C) The relative mRNA expression of selected nuclear receptors. (D) The relative mRNA expression of CYPs. (E) The protein expression of PXR. (F) The protein expression of CYP3A4. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. **P* < 0.05; ***P* < 0.01;

DMD-AR-2020-000316R1

*** $P < 0.001$ vs. the si-NC group.

Fig. 5. Effect of HNF4A knockdown on the RIF-induced expression of genes in Huh7 cells. si-HNF4A or si-NC transfected Huh7 cells were treated with RIF for 24 hours. The RIF-induced expression of genes after HNF4A knockdown in the RNA level was determined by qRT-PCR method. (A) Expression of HNF4A and HNF4A-AS1 at the RNA level. (B) The relative mRNA expression of selected nuclear receptors. (C) The relative mRNA expression of CYPs. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ vs. the control group.

Fig. 6. Suppression of HNF4A-AS1 elevated the enrichment levels of histone modifications and HNF4A in the promoter of *CYP3A4* in Huh7 cells. (A) Schematic locations of HNF4A response elements (HREs) and ChIP-qPCR primers in the *CYP3A4* promoter. (B) Enrichment levels of H3K4me3 around the HREs in *CYP3A4* promoter. (C) Enrichment levels of HNF4A around the HREs in *CYP3A4* promoter. ChIP-qPCR analysis was conducted to measure the enrichment levels of H3K4me3 and HNF4A in the promoter of *CYP3A4* and data were presented as fold enrichment over IgG. Data are shown as mean \pm S.D. of three independent experiments. Statistical analysis was performed using unpaired Student's *t* test. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ vs. the si-NC group.

Table 1. Primer sequences for qRT-PCR analysis

Gene	Sequence (5'-3')	Orientation
<i>GAPDH</i>	GCACCGTCAAGGCTGAGAAC	Sense
	TGGTGAAGACGCCAGTGGA	Antisense
<i>HNF4A-AS1</i>	TGGAGCTGGGATCTGACACT	Sense
	ATGACCGGTGTGCAGTCAAG	Antisense
<i>HNF4A</i>	CGTGCTGCTCCTAGGCAA	Sense
	GTCAAGGATGCGTATGGACAC	Antisense
<i>HNF1A</i>	TGGGTCCTACGTTACCAAC	Sense
	TCTGCACAGGTGGCATGAGC	Antisense
<i>CAR</i>	CAGAAGCGGTGCATACCTCA	Sense
	AGATGGCACTGACTCTGCAA	Antisense
<i>AhR</i>	ACATCACCTACGCCAGTCGC	Sense
	TCTATGCCGCTTGGAAGGAT	Antisense
<i>PXR</i>	CAACCTACATGTTCAAAGGCATC	Sense
	ACACTCCCAGGTTCCAGTCTC	Antisense
<i>CYP1A2</i>	CCAACGTCATTGGTGCCATG	Sense
	GTGATGTCCCGGACACTGTTC	Antisense
<i>CYP2E1</i>	TCCTAGGGCACAGTCGTAGT	Sense
	AGGGAAGGTACTGCCTCTGAT	Antisense
<i>CYP2B6</i>	AGACGCCTTCAATCCTGACC	Sense
	CCTTCACCAAGACAAATCCGC	Antisense
<i>CYP2D6</i>	CCTACGCTTCCAAAAGGCTTT	Sense
	AGAGAACAGGTCAGCCACCACT	Antisense
<i>CYP2C8</i>	GGACTTTATGGATTGCTTCCTG	Sense
	CCATATCTCAGAGTGGTGCTTG	Antisense
<i>CYP2C9</i>	GACATGAACAACCCTCAGGACTTT	Sense
	TGCTTGTCTGTCTGTCCCA	Antisense
<i>CYP2C19</i>	GAACACCAAGAATCGATGGACA	Sense
	TCAGCAGGAGAAGGAGAGCATA	Antisense
<i>CYP3A4</i>	CCCTTTGGAAGTGGACCCAG	Sense
	ACGGTGCCATCCCTTGACTC	Antisense

Table 2. Primer sequences for ChIP-qPCR

Primer	Position	Sequence (5'-3')	Orientation
Primer 1	-362 ~ +53	GCACCGTCAAGGCTGAGAAC	Sense
		GCCTGAGAACACCAGAGACC	Antisense
Primer 2	-6,093 ~ -6,574	CAGCAGACAGAGAACCCAGA	Sense
		GCCTGAGAACACCAGAGACC	Antisense




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6