

DMD-MR-2021-000418

Adverse Drug Reactions and Toxicity of the FDA- approved Antisense Oligonucleotide Drugs

Feryal Alhamadani, Kristy Zhang, Rajvi Parikh, Hangyu Wu, Theodore P.

Rasmussen, Raman Bahal, Xiao-bo Zhong, and José E. Manautou

*Department of Pharmaceutical Sciences, School of Pharmacy, University
of Connecticut, Storrs, Connecticut (F.A., K.Z., H.W., T.P.R., R.B., X-b.Z.,*

J.E.M.); Department of Physiology and Neurobiology, University of

Connecticut, Storrs, Connecticut (R.P)

DMD-MR-2021-000418

Running Title: ADRs and toxicity of the FDA-approved ASO drugs

Corresponding author: Dr. Jose E. Manautou, Department of Pharmaceutical Sciences, School of Pharmacy, University of Connecticut, 69 N Eagleville Road, Storrs, Connecticut 06269, USA. Telephone: 860-486-3852; Fax: 860-486-5792; E-mail: jose.manautou@uconn.edu

Dr. Xiao-bo Zhong, Department of Pharmaceutical Sciences, School of Pharmacy, University of Connecticut, 69 N Eagleville Road, Storrs, Connecticut 06269, USA. Telephone: 860-486-3697; Fax: 860-486-5792; E-mail: xiaobo.zhong@uconn.edu

Number of text pages: 42

Number of figures: 1

Number of tables: 3

Number of references: 63

Number of words in Abstract: 250

Number of words in Introduction: 351

Number of words in the main text: 6466

DMD-MR-2021-000418

ABBREVIATIONS:

ADR, adverse drug reaction; ALT, alanine aminotransferase; AMD, age-related macular degradation; ASO, antisense oligonucleotide; AST, aspartate aminotransferase; CMV, cytomegalovirus; DMD, Duchenne muscular atrophy; FDA, Food and Drug Administration; MRI, magnetic resonance imaging; pre-mRNA, pre-messenger RNA; REMS, risk evaluation and mitigation strategy; SMA, spinal muscular atrophy; SMN2, survival motor neuron-2; TTR: transthyretin; ULN, upper limit of normal; UPCR, urine protein to creatinine ratio; VEGF, vascular endothelial growth factor; VLDL, very low-density lipoprotein.

DMD-MR-2021-000418

ABSTRACT

The market for large molecule biological drugs has grown rapidly, including antisense oligonucleotide (ASO) drugs. ASO drugs work as single-stranded synthetic oligonucleotides that reduce production or alter functions of disease-causing proteins through various mechanisms, such as mRNA degradation, exon skipping, and ASO-protein interactions. After the first ASO drug Fomivirsen was approved in 1998, the US Food and Drug Administration has approved 10 ASO drugs to date. Although ASO drugs are efficacious in treating some diseases that are untargetable by small-molecule chemical drugs, concerns on adverse drug reactions (ADRs) and toxicity cannot be ignored. Illustrative of this, Mipomersen was recently taken off the market due to its hepatotoxicity risk. This paper reviews ADRs and toxicity from FDA drug labeling, preclinical studies, clinical trials, and post-marketing real-world studies on the ten FDA-approved ASO drugs, including Fomivirsen and Pegaptanib Mipomersen, Nusinersen, Inotersen, Defibrotide, Eteplirsen, Golodirsen, Viltolarsen, and Casimersen. Unique and common ADRs and toxicity for each ASO drug are summarized here. The risk of developing hepatotoxicity, kidney toxicity, and hypersensitivity reactions co-exists for multiple ASO drugs. Special precautions need to be in place when certain ASO drugs are administered. Further discussion is extended on studying the mechanisms of ADRs and toxicity of these drugs, evaluating the existing physiological and pathological states of patients, optimizing the dose and route of administration, and formulating personalized treatment plans to improve the clinical utility of FDA-approved ASO drugs and discovery and development of new ASO drugs with reduced ADRs.

DMD-MR-2021-000418

Significance Statement

The current review provides a comprehensive analysis of unique and common ADRs and the toxicity of FDA-approved ASO drugs. The information can help better manage the risk of severe hepatotoxicity, kidney toxicity, and hypersensitivity reactions in the usage of currently approved ASO drugs and the discovery and development of new and safer ASO drugs.

Introduction

If a drug at an administered dose results in a plasma drug concentration over the minimal toxic concentration, it may cause adverse drug reactions (ADRs) and toxicity. This is also the case for next-generation therapeutics, such as antisense oligonucleotide (ASO) drugs. ASOs are a class of biological entities with a fast-growing pace of development in recent years, with an innovative and promising approach for treating both rare and common human diseases. ASO drugs are therapeutic agents capable of reducing the abundance of proteins that cause human diseases (Roberts et al., 2020). ASO drugs employ single-stranded chains of synthetic oligonucleotides that block the ability of mRNA molecules to be translated into proteins through a variety of mechanisms (Crooke, 2017). Compared to small molecule-based chemical drugs, ASO drugs have shown distinct advantages in targeting diseases at the genetic level and may become highly useful in routine clinical practice to treat a diverse number of human diseases and conditions.

Although ASO drugs are highly promising pharmacotherapies, ADRs and toxicity are equally concerning in the development of ASO drugs as with more conventional therapeutic agents. The safety assessment of ASOs has already revealed that this class of drugs has some shared ADRs with varying levels of severity. Some ASO drugs have shown certain toxicity levels during clinical trials, such as the case of Mipomersen and hepatotoxicity (Wong and Goldberg, 2014). The U.S. Food and Drug Administration (FDA) withdrew Mipomersen in 2019 from the market after only five years since use approval due to its hepatotoxic liability. Therefore, it is critical to understand what types

DMD-MR-2021-000418

of ADRs and toxicities can be caused by ASO or ASO drug candidates and understand the underlying molecular mechanisms. The purpose of this review is to provide critical analysis of both frequently encountered and idiosyncratic ADRs and toxicities among the ten FDA-approved ASO drugs. The article will cover a brief historical perspective on ASO drugs, recent fundamental discoveries, current challenges and knowledge gaps, and future perspective directions in consideration of ADRs and toxicity of the ten FDA-approved ASO drugs.

Brief Historical Perspective

The development of ASOs as therapeutic agents spans more than 30 years since ASOs were first demonstrated to influence RNA processing and modulate the translation of proteins from mRNAs back in 1978 (Stephenson and Zamecnik, 1978). It took a subsequent 20 years for the first ASO drug, Fomivirsen (Table 1), to be approved by the FDA in 1998 (Marwick, 1998). After a slow pace during the following 18 years, the FDA approved only two new ASO drugs (Pegaptanib in 2004 and Mipomersen in 2013). Since then, the pace has markedly quickened, and the FDA has approved a total of seven new drugs (Defibrotide, Eteplirsen, Nusinersen, Inotersen, Golodirsen, Viltolarsen, and Casimersen) up to the present time. Currently, over 100 ASOs are in the clinical drug development pipelines, and many more are in various pre-clinical development stages (Dhuri et al., 2020). In the coming years, ASO drugs may become

DMD-MR-2021-000418

frequently used for routine clinical treatment of human diseases, and perhaps they can become as prominent in their clinical utility as small-molecule chemical drugs.

ADRs and toxicity are always of concern during ASO drug discovery and development. For example, when Mipomersen was approved by the FDA, a black box warning existed regarding the risk of liver enzymes (transaminases) elevation and hepatic steatosis. The drug was only prescribed to patients who participated in a Risk Evaluation and Mitigation Strategy (Dixon et al., 2014). After further use for several years under restrictive guidelines, the FDA withdrew Mipomersen from the market in 2019 due to an unacceptable risk of hepatotoxicity. Numerous class-wide and idiosyncratic ADRs and toxicities are known for other FDA-approved ASO drugs. Therefore, understanding why ADRs and toxicity occur at mechanistic and physiological levels is prudent for developing future efficacious ASO drugs while avoiding such safety liabilities.

Key Recent Advances

A comprehensive analysis of ADRs and associated toxicity is provided for each of the following FDA-approved ASO drugs: Fomivirsen, Pegaptanib, Mipomersen, Defibrotide, Eteplirsen, Nusinersen, Inotersen, Golodirsen, Viltolarsen, and Casimersen.

Fomivirsen (Vitravene):

DMD-MR-2021-000418

Fomivirsen is the first FDA-approved ASO drug. Known commercially as Vitravene, it was manufactured by Isis Pharmaceuticals (now Ionis Pharma) and received FDA approval on August 26, 1998 for the treatment of cytomegalovirus (CMV) retinitis (Roehr, 1998). CMV retinitis is an infection that targets the light-sensing cells in the retina (Port et al., 2017). Fomivirsen is a phosphorothioate ASO drug that inhibits human CMV replication through an antisense mechanism (de Smet et al., 1999).

The commonly observed ADRs of Fomivirsen in clinical trials were increased intraocular pressure and ocular inflammation. Approximately 25% of patients experienced these ADRs during loading dosing (Perry and Balfour, 1999; Vitravene Study, 2002). Other ADRs of Fomivirsen reported during the clinical trials included ophthalmic dysfunctions, such as anterior chamber inflammation, abnormal vision, cataract, decreased visual acuity, eye pain, and floaters. Fomivirsen was observed to also accumulate in the plasma at detectable concentrations, and also in the liver, and kidneys. Some of the more frequent systemic side effects included anemia, diarrhea, abnormal pain, asthenia, fever, infection, nausea, headache, pneumonia, sepsis, rash, vomiting, sinusitis, and systemic CMV infection. Additional ADRs that occurred less frequently in patients included application site reactions, decreased peripheral vision, hypotony, corneal edema, eye irritation, retinal vascular disease, optic neuritis, vitreous opacity, and visual field defects.

Limited open label clinical studies have been done on Fomivirsen to evaluate its safety. However, systemic accumulation of Fomivirsen was noted in animal studies following single or repeated intravitreal injections in monkeys at levels below 70 ng/mL in plasma and 350 ng/g in tissues. Metabolites of Fomivirsen were detected in the liver,

DMD-MR-2021-000418

kidney, and plasma at a concentration of 14 ng/mL in plasma and 70 ng/g in tissues in monkeys treated with Fomivirsen every other week for up to 3 months.

Fomivirsen was withdrawn from the European market in 2002 and the U.S. in 2006 due to the development of highly effective antiretroviral therapies for the same indication. In addition, the number of CMV cases dramatically decreased and Ionis Pharmaceuticals stopped marketing the drug.

Pegaptanib Sodium (Macugen):

Pegaptanib sodium, sold under the name of Macugen, is manufactured by Eyetech Pharmaceuticals and was approved by the FDA in December 17, 2004 for the treatment of neovascular (wet) age-related macular degeneration (AMD) (Tobin, 2006). AMD can present itself in two forms, as blurred or no vision in the center of the visual field (Nowak, 2006). The number of AMD patients was estimated to be 196 million in 2020 and the projections are for 288 million cases by 2040 globally (Jonas et al., 2017). Macugen is an aptamer oligonucleotide with a three-dimensional conformation, enabling it to bind to extracellular vascular endothelial growth factor (VEGF) as an antagonist (Katz and Goldbaum, 2006). VEGF induces angiogenesis and increases vascular permeability and inflammation, which contribute to AMD. Pegaptanib inhibits VEGF, suppressing pathological neovascularization.

ADRs were observed in two clinical trials with controlled, double masked, randomized studies in approximately 1,200 patients with neovascular AMD (Group et al., 2006). The patients were randomized and administered either 0.3 mg of Pegaptanib sodium or placebo once every six weeks for 8-9 treatments, intravitreally. The group

DMD-MR-2021-000418

treated with 0.3 mg Pegaptanib sodium exhibited a statistically significant reduction in the rate of vision decline as compared to the placebo group. The most frequently reported ADRs in the clinical studies included anterior chamber inflammation, blurred vision, cataract, conjunctival hemorrhage, corneal edema, eye discharge, eye irritation, eye pain, hypertension, increased intraocular pressure (IOP), ocular discomfort, punctate keratitis, reduced visual acuity, visual disturbance, vitreous floaters, and vitreous opacities (observed in 10-40% of patients). Approximately 6-10% of patients treated with Pegaptanib sodium demonstrated ADRs that included ocular (blepharitis, conjunctivitis, photopsia, and vitreous disorder) and non-ocular (bronchitis, diarrhea, dizziness, headache, nausea, and urinary tract infection) reactions. Approximately 1-5% of patients treated exhibited ocular side effects, such as allergic conjunctivitis, conjunctival edema, corneal abrasion, corneal deposits, corneal epithelium disorder, endophthalmitis, eye inflammation, eye swelling, eyelid irritation, meibomianitis, mydriasis, periorbital hematoma, retinal edema, and vitreous hemorrhage. Approximately 1-5% of patients reported some non-ocular events, including arthritis, bone spurs, carotid artery occlusion, cerebrovascular accident, chest pain, contact dermatitis, contusion, diabetes mellitus, dyspepsia, hearing loss, pleural effusion, transient ischemic attack, urinary retention, vertigo, and vomiting. Serious ADRs related to the treatment with Pegaptanib sodium occurred in <1% of those treated, and these included endophthalmitis, retinal detachment, and iatrogenic traumatic cataract.

Warnings and precautions in the FDA-approved drug label for Macugen include endophthalmitis, increases in intraocular pressure, and anaphylaxis. Monitoring for

DMD-MR-2021-000418

infection, intraocular pressure, and the perfusion of the optic nerve head is recommended by the FDA.

ADRs of Pegaptanib in post-marketing settings were further compared with the pre-marketing clinical trials findings (Penedones et al., 2014). Both in pre- and post-marketing settings, ocular ADRs were more frequent than non-ocular ADRs. In addition to the observed ADRs in the pre-marketing settings, the post-marketing findings have been typified by an increased number of events, such as retinal pigmented epithelium tears, thromboembolic events, and mortality.

Mipomersen (Kynamro):

Mipomersen, was manufactured by Genzyme Corporation and approved by the FDA in January 29, 2013 (Hair et al., 2013) to reduce low density lipoprotein-cholesterol (LDL-C), apolipoprotein B (apo B), total cholesterol (TC), and non-high density lipoprotein-cholesterol (non-HDL-C) in patients with homozygous familial hypercholesterolemia (HoFH). Mipomersen is an ASO drug that inhibits APO B-100 synthesis used in combination with diet modification and lipid lowering medications (Bell et al., 2011).

ADRs were described in four phase 3 clinical trials (ClinicalTrials.gov NCT00607373, NCT00706849, NCT00770146, and NCT00794664). These clinical trials were randomized, double-blinded, placebo-controlled studies with a total of 390 patients at a ratio of 2:1 for weekly subcutaneous injections of 200 mg of mipomersen or placebo for a median treatment of 25 weeks (Santos et al., 2015). The most common ADR, injection site reactions, occurred in 84% of patients and typically consisted of pain, local swelling,

DMD-MR-2021-000418

erythema, pruritus, and tenderness. Furthermore, flu-like symptoms, such as fatigue, influenza-like illness, chills, pyrexia, arthralgia, myalgia, or malaise were reported in approximately 30% of the patients within 2 days of treatment initiation (Reeskamp et al., 2019). Other less common ADRs (3-15%) included vascular disorder (hypertension), cardiac disorders (angina pectoris and palpitations), musculoskeletal and connective tissue disorders (pain in extremity and musculoskeletal pain), gastrointestinal disorders (nausea, vomiting, and abdominal pain), and nervous system and psychiatric disorders (headache, insomnia).

A black box warning for the risk of hepatotoxicity, including elevation of transaminases and hepatic steatosis, was placed in the FDA-approved drug labeling of Mipomersen. In clinical trials, 12% of the patients with HoFH treated with Mipomersen compared to 0% of subjects treated with placebo had elevations in alanine transaminase (ALT) ≥ 3 x the upper limits of normal (ULN), and 9% in the Mipomersen group compared to 0% of placebo patients had at least one elevation in ALT ≥ 5 x ULN (Thomas et al., 2013). The FDA suggested that before initiating treatment with Mipomersen, a full panel of liver function tests that include ALT, aspartate transaminase (AST), total bilirubin, and alkaline phosphatase should be conducted. Patients with moderate or severe hepatic impairment, or active liver diseases, including unexplained persistent elevations in serum transaminases are contraindicated in the use of Mipomersen.

Another FDA warning is hepatic steatosis. During clinical trials in patients with HoFH and hyperlipidemia, the median absolute increase in hepatic fat was 10% after 26 weeks of treatment compared to baseline (0%). The increase in hepatic fat (hepatic

DMD-MR-2021-000418

steatosis) is a risk factor for advanced liver diseases, including steatohepatitis and cirrhosis (Nassir et al., 2015). Because of the risk of hepatotoxicity, Mipomersen is only available through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS), called the Kynamro REMS (Gupta, 2015). Mipomersen is no longer available after its distribution was terminated in 2019 due to the risk of hepatotoxicity (Lui et al., 2021).

Defibrotide Sodium (Defitelio):

Defibrotide sodium, manufactured by Jazz Pharmaceuticals and sold under the name Defitelio, was approved by the FDA in March 30, 2016 for the treatment of adult or pediatric patients with hepatic veno-occlusive disease (VOD) and renal or pulmonary dysfunction following hematopoietic stem-cell transplantation (HSCT) (Choy, 2016). Defibrotide is a mixture of single- and double-stranded phosphodiester oligonucleotides (9-80, average 50 nucleotides) extracted through controlled depolymerization of porcine gut mucosa (Kornblum et al., 2006). Although the mechanism of action has not been fully elucidated, *in vitro* studies showed that Defibrotide sodium reduces hepatic sinusoidal endothelial cell activation and protects them from damage. Also *in vitro*, it promotes hydrolysis of fibrin clots by enhancing the enzymatic activity of factors such as plasmin and tissue plasminogen activator (Richardson et al., 2017). The clinically recommended dosage of Defibrotide is 6.25 mg/kg given every 6 hours as an intravenous infusion over 2 hours, which should be dosed based on the baseline body weight of the patient.

DMD-MR-2021-000418

ADRs of Defibrotide were defined based on interventional and expanded clinical trial studies, including NCT00272948, NCT00358501, and NCT00628498 (Kessler-Icekson et al., 1988; Corbacioglu et al., 2012; Richardson et al., 2016). Patients with VOD received either Defibrotide at 6.25 mg/kg intravenously every 6 hours or placebo (given as a 2-hour infusion) for at least 21 days up to 60 days. The most common ADRs included hypotension, diarrhea, vomiting, nausea, and epistaxis (incidence >10%). The most common ADRs with an incidence of 5% or greater included hypotension (11%) and pulmonary alveolar hemorrhage (7%).

The warnings and precautions in the FDA-approved drug label of Defitelio include hemorrhage and hypersensitivity reactions. As previously stated, since Defibrotide increases the *in vitro* activity of fibrinolytic enzymes, it may increase the risk of bleeding in patients with VOD after HSCT. Hypersensitivity reactions occurred in less than 2% of patients and included rashes, urticaria, and angioedema. Patients on Defibrotide treatment should be monitored for both bleeding and hypersensitivity reactions.

Eteplirsen (Exondys 51)

Eteplirsen, is manufactured by Sarepta Therapeutics and received an accelerated FDA-approval on September 19, 2016 (Syed, 2016) to treat Duchenne muscular dystrophy (DMD). DMD is the most common type of muscular dystrophy, affecting 7.1 per 100,000 males and 2.8 per 100,000 in the general global population (Crisafulli et al., 2020). Muscle weakness usually begins around the age of four, worsens quickly, and most are unable to walk by the age of twelve (Bushby et al., 2010). DMD is caused by mutations of the dystrophin gene, encoding dystrophin protein responsible for

DMD-MR-2021-000418

connecting the actin cytoskeleton of each muscle fiber to the underlying extracellular matrix (Elangkovan and Dickson, 2021). Although there is no known cure, some medications may help to improve symptoms, including steroids, anticonvulsants, and immunosuppressants (Beytia Mde et al., 2012). Recently, ASO-mediated exon-skipping therapies have been developed as precision medicine for the treatment of DMD (Matsuo, 2021). Eteplirsen is an ASO drug indicated for the treatment of DMD in patients who have a confirmed mutation of the dystrophin gene that is circumvented by ASO-induced skipping of exon 51 during splicing of the dystrophin pre-mRNA. Exondys 51, as Eteplirsen is commercially known, binds to exon 51 of dystrophin pre-mRNA, resulting in exclusion of this exon during mRNA processing (Lim et al., 2017). The clinical dose recommended for Exondys 51 is 30 mg/kg once weekly given as 35 to 60 min intravenous infusion.

Eteplirsen's ADRs were reported in a phase 2 clinical study with 12 participants (NCT01396239). This was a double-blind placebo-controlled randomized trial (Mendell et al., 2013). The participating patients were randomized to weekly IV infusions of 30 and 50 mg/kg Eteplirsen, or placebo for 24 weeks. The most common ADRs of Eteplirsen were balance disorder, vomiting, and contact dermatitis. These ADRs had an incidence at least 25% higher than placebo in all 12 DMD patients treated with either dose of Eteplirsen (Mendell et al., 2013). Because of the small numbers of patients in this study, these ADR frequencies may not reflect the frequencies observed in practice. In a follow-up phase 3 clinical trial (NCT02255552) with 109 patients who received at least 30 mg/kg/week of Eteplirsen for up to 144 weeks, at least 10% of patients reported vomiting, contusion, excoriation, arthralgia, rash, catheter site pain, and respiratory tract

DMD-MR-2021-000418

infection. Less than 1% of patients had bronchospasm, cough, dyspnea, fever, flushing, hypersensitivity reaction, hypotension and urticaria from post-marketing surveillance and case reports. Transient erythema, facial flushing, and elevated temperature had been reported on days of Eteplirsen infusion. Since only three clinical studies had evaluated Eteplirsen, the information for monitoring parameters is lacking. There is no animal data available to assess the use of Eteplirsen during pregnancy and the effect of the drug on milk production.

A warning and precaution for hypersensitivity reaction has been placed in the FDA-approved drug label of Eteplirsen. In the clinical trials, some patients treated with Eteplirsen presented with hypersensitivity reactions, including rash, pyrexia, flushing, urticaria, bronchospasm, dyspnea, and cough. The FDA recommends following the institute appropriate medical treatment for these ADRs. Another recommendation is to consider slowing drug infusion or interrupting Eteplirsen therapy altogether if hypersensitivity reactions arise.

Nusinersen (Spinraza)

Nusinersen, is manufactured by Biogen and was approved by the FDA in December 23, 2016 for the treatment of spinal muscular atrophy (SMA) in pediatric and adult patients (Hoy, 2017). SMA is a rare neuromuscular disorder that results in the loss of motor neurons and progressive muscle wasting. Reported prevalence of SMA is approximately 1-3 per 100,000 among different populations (Bertolin et al., 2019; Ito et al., 2021). Nusinersen is the first FDA-approved drug to treat SMA (Son and Yokota, 2018). This is a survival motor neuron-2 (SMN2)-directed ASO drug designed to treat

DMD-MR-2021-000418

SMA caused by mutations in chromosome 5q that lead to SMN protein deficiency (Claborn et al., 2019). Nusinersen is initiated as 4 loading doses and a maintenance dose. The first three doses are administered at 14-day intervals and the fourth one is administered 30 days after the third dose. Maintenance doses are given every 4 months after the fourth loading dose.

ADRs were determined in numerous clinical trials, including NCT01494701, NCT01703988, NCT01839656, NCT02193074, NCT02292537, NCT01780246, and NCT02052791 (Darras et al., 2019). The ADRs were analyzed from 323 infants and children that either received Nusinersen at 12 mg per administration at 14-day intervals or underwent sham procedures (Darras et al., 2019). The most common ADRs reported in approximately 20% of treated patients were vomiting, constipation, back pain, headache, pyrexia, and lower respiratory tract infections. ADRs of lower frequency included decreased weight, seasonal allergy symptoms, falls, epistaxis, flatulence, upper respiratory tract congestion, urinary tract infection, and ear infection.

After administration of Nusinersen therapy is associated with thrombocytopenia and coagulation abnormalities. In the clinical studies (Mercuri et al., 2018; Darras et al., 2019), 16% Nusinersen-treated patients with normal or elevated platelet levels at baseline developed a platelet level below the lower limits of normal. Because of the risk of thrombocytopenia and coagulation abnormalities, patients treated with Nusinersen may be at increased risk of bleeding complications. Nusinersen is present in and excreted by the kidney. Renal toxicity, including potentially fatal glomerulonephritis, was observed after administration of Nusinersen. In the same clinical studies, 58% of Nusinersen-treated patients had elevated urine protein, compared to 34% sham-control

DMD-MR-2021-000418

group. The mechanism of renal and liver injury is most likely due to drug accumulation with repeated doses, because subcutaneously or intravenously administered ASOs bind to plasma proteins that are filtered at the glomerulus and then reabsorbed in the proximal tubule (Frazier, 2015). Given these safety findings, warnings and precautions for thrombocytopenia, coagulation abnormalities, and renal toxicity have been placed in the FDA-approved drug label of Nusinersen. Platelet count, coagulation tests, and quantitative spot urine protein testing at baseline and prior to each dose of Nusinersen are highly recommended by the FDA.

The ADRs identified in the clinical trials have been further confirmed from post-marketing and case reports, including angioedema, aseptic meningitis, hydrocephalus, hypersensitivity reaction, maculopapular rash, meningitis, serious infection, and skin rash (Mendonca et al., 2021). Long term animal studies to evaluate carcinogenic potential of Nusinersen were not performed. There are not sufficient data to evaluate Nusinersen-associated developmental risks in pregnant women and no adverse events were observed in animal studies of fetal development. In intrathecal toxicity studies in juvenile primates, administration of Nusinersen resulted in brain histopathology at the moderate and high doses and acute, transient deficits in lower spinal reflexes at the high dose in each study.

An observational clinical trial (NCT04419233) is expected to recruit 50 participants to evaluate the safety of Nusinersen in the post-marketing setting in China and is estimated to be completed in January 2023. Another observational clinical trial (NCT04317794) is also in recruitment phases with an expected 145 participants to evaluate the safety of Nusinersen in the post-marketing setting in Korea. The estimated

DMD-MR-2021-000418

completion date of the study is July 2027. The data released from these studies will provide information for ADRs and toxicity in diverse populations.

Inotersen (Tegsedi)

Inotersen, is manufactured by Ionis Pharmaceuticals and was approved by the FDA in October 5, 2018 to treat the polyneuropathy of hereditary transthyretin (TTR)-mediated amyloidosis (hATTR-PN) in adults (Keam, 2018). hATTR-PN is a rare disease with an estimated 10,000 afflicted patients worldwide (Schmidt et al., 2018), but without treatment. This is an inherited, progressive, and fatal disease. Current options for the treatment of the disease include liver transplantation and TTR stabilizers of tafamidis and diflunisal (Luigetti et al., 2020). Recently, more promising gene-silencing therapies have been developed, including Patisiran, the first FDA-approved siRNA drug (Zhang et al., 2021) and the ASO drug Inotersen. Inotersen works by degrading mutant and wild type TTR mRNAs by binding to the TTR mRNAs therefore reducing serum TTr and TTR protein deposits in tissues (Gales, 2019). The clinically recommended dose for Inotersen is 284 mg once weekly subcutaneously. The injection should be given on the same day every week.

ADRs were observed in a phase 2/3 clinical trial (NCT01737398). This clinical trial was an international, randomized, double-blinded, placebo-controlled study to assess the efficacy and safety of Inotersen (Benson et al., 2018). During the clinical trial, 173 adult patients with polyneuropathy caused by hATTR-PN were randomly assigned in a 2:1 ratio to receive weekly subcutaneous injections of Inotersen (300 mg) or placebo for 64 weeks. The most common ADRs occurring in approximately 20% of Inotersen-

DMD-MR-2021-000418

treated patients and more frequently than with placebo, were nausea, fever, fatigue, thrombocytopenia, injection site reactions, and headaches. The less common ADRs (<20%) included peripheral edema, chills, anemia, vomiting, myalgia, decreased renal function, arrhythmia, arthralgia, pre-syncope, decreased appetite, paresthesia, dyspnea, elevated liver function test, orthostasis, influenza like illness, contusion, bacterial infection, eosinophilia, and dry mouth.

The FDA mandated the inclusion of a black box warning in the drug's packing insert describing the safety concerns with the use of Inotersen. The most serious concerns are thrombocytopenia and glomerulonephritis. In the clinical trial of NCT01737398, Inotersen caused low platelet count, which can lead to sudden and unpredictable life-threatening thrombocytopenia (Benson et al., 2018). Inotersen-associated thrombocytopenia was also reported in the clinical trials of the NEURO-TTR (NCT04136184) and NEURO-TTR open-label extension (Narayanan et al., 2020). The key strategies to prevent thrombocytopenia include monitoring the platelet count and withholding therapy if the patient develops signs or symptoms of thrombocytopenia (unusual or prolonged bleeding, neck stiffness or atypical headache). Caution should be used when using antiplatelet drugs or anticoagulants concomitantly with Inotersen because of the risk of bleeding. Inotersen may also cause glomerulonephritis that may result in dialysis-dependent renal failure. In the same clinical trial, glomerulonephritis occurred in 3% Inotersen-treated patients, compared with no patient on placebo (Benson et al., 2018). The mechanism of action for glomerulonephritis can be explained by accumulation of ASOs in proximal tubule cells, which leads to increased tubular proteinuria. Urine protein to creatinine ratio (UPCR) greater than 5 times the ULN

DMD-MR-2021-000418

occurred in 15% of Inotersen-treated patients, compared to 8% of patients on placebo. Therefore, Inotersen should generally not be initiated in patients with a UPCR of 1,000 mg/g or greater (Benson et al., 2018). Because of the risks of thrombocytopenia and glomerulonephritis, Inotersen is available only through a restricted distribution program called the Tegsedi REMS.

Other warnings and precautions have been indicated in the FDA-approved drug label of Inotersen, including stroke and cervicocephalic arterial dissection, inflammatory and immune effects, liver injury, hypersensitive reactions, and reduced serum vitamin A levels. In the clinical trial, stroke and cervicocephalic arterial dissection occurred within 2 days of the first Inotersen dose. Patients also experienced symptoms of cytokine release (e.g. nausea, vomiting, muscular pain, and weakness) and a high sensitivity C-reactive protein level greater than 100 mg/L (Benson et al., 2018). The liver is a site where ASOs accumulates. In the clinical studies, 8% of Inotersen-treated patients had increased ALT levels at least 3 times the ULN, compared to 3% of patients on placebo. Furthermore, 3% of Inotersen-treated patients had ALT values at least 8 times the ULN, compared to no patients on placebo. Monitoring ALT, AST, and total bilirubin at baseline and every four months during treatment with Inotersen is recommended.

From post-marketing and case reports, < 1% of patients reported autoimmune hepatitis, cerebrovascular accident, coronary artery dissection, hepatobiliary disease, hypersensitivity reaction, immune thrombocytopenia, lower back pain, paraplegia, speech disturbance, vasculitis, and weight loss.

A worldwide clinical trial for pregnancy outcome in women treated with Inotersen is underway (NCT 04270058) and is expected to be completed in 2030. A data collection

DMD-MR-2021-000418

clinical trial (NCT04850105) with an estimated 240 participants currently under recruitment is aimed at further characterizing the long-term safety of Inotersen in patients with hATTR-PN under real-world conditions. Aforementioned studies will provide more evidence on how to manage safety concerns of Inotersen.

Golodirsen (Vyondys 53):

Golodirsen, is manufactured by Sarepta Therapeutics and was approved by the FDA on December 12, 2019 (Heo, 2020). Golodirsen is indicated for the treatment of DMD in patients with a mutation in the dystrophin gene, which can be treated by ASO-induced skipping of exon 53 in the dystrophin pre-mRNA. Golodirsen is designed to bind to exon 53 of dystrophin pre-mRNA, thus excluding the exon during processing. The clinically recommended dose is 30 mg/kg once weekly in 35 to 60 min infusion intravenously.

ADRs were reported in a phase 1/2 clinical trial (NCT02310906). This was a randomized, placebo-controlled dose-titration study (part 1) followed by open-label evaluation (part 2) with 39 participants to assess safety and tolerability of Golodirsen (Frank et al., 2020). The common ADRs observed in at least 20% of patients included headaches, pyrexia, fall, abdominal pain, nasopharyngitis, cough, vomiting, and nausea. Less common ADRs reported in ~ 5% of treated patients were administration site pain, diarrhea, dizziness, ligament sprain, contusion, influenza, oropharyngeal pain, rhinitis, skin abrasions, ear infections, seasonal allergies, tachycardia, site related reaction, constipation, and bone fractures (Frank et al., 2020). An extension study (NCT03532542) to evaluate Golodirsen in patients with DMD is under enrolling stages

DMD-MR-2021-000418

by invitation in a phase 3 study with an estimated 260 participants and is expected to be completed in August 2026.

Warnings and precautions for hypersensitivity reactions and renal toxicity have been placed in the FDA-approved drug label of Golodirsén. Hypersensitivity reactions occurred in Golodirsén-treated patients in the clinical trials with phenotypes including rash, pruritus, pyrexia, dermatitis, urticaria, and skin exfoliation. If a hypersensitivity reaction occurs, patient care should be managed following the institute appropriate medical treatment and slower infusion rates or interruption of Golodirsén therapy should be considered. Renal toxicity was not observed in the clinical studies; however, it was reported in preclinical studies with different species animals receiving Golodirsén. Mice administered either IV or subcutaneous injections of Golodirsén developed renal tubular degeneration, as well as degeneration of the transitional epithelium of the urinary bladder at all tested doses. Tubular degeneration by IV administration was accompanied by increases in serum urea nitrogen. Primates after IV administration of Golodirsén also developed kidney dysfunction with tubular basophilia, dilatation, and mononuclear cell infiltration. The FDA recommends monthly proteinuria monitoring by dipstick urinalysis and monitoring of serum cystatin C every three months.

Sarepta Therapeutics is currently conducting a placebo controlled, post-marketing confirmation trial named ESSENCE (NCT02500381) to support the accelerated approval. The trial is expected to be concluded in April 2024.

Viltolarsén (Viltepso):

DMD-MR-2021-000418

Viltolarsen, sold under the name of Viltepso and manufactured by Nippon Shinyaku Co, was granted accelerated approval by the FDA on July 12, 2020 based on its capacity to increase dystrophin levels in skeletal muscle in DMD patients (Dhillon, 2020). Continued approval is dependent upon verification of the clinical benefits in confirmatory trials. Viltolarsen is an ASO drug also indicated for the treatment of DMD in patients who have a confirmed mutation of the dystrophin gene that is treatable by ASO-induced skipping of exon 53. Viltolarsen is designed to bind to exon 53 of dystrophin pre-mRNA, resulting in exclusion of this exon during mRNA processing. This exclusion leads to the increase in dystrophin production, which helps keep muscle cells intact. The clinically recommended dose is 80 mg/kg once weekly in 60 min infusion intravenously.

ADRs were reported in a phase II clinical trial (NCT02740972). This is a multi-center, 2-period, randomized, placebo-controlled study with 16 participants (Clemens et al., 2020). The initial period of the clinical study (4 weeks) consisted of patients being randomized (double blind), with administration via an injection with Viltolarsen or placebo. All patients then received 20 weeks of open label Viltolarsen (40 mg/kg or 80 mg/kg) once a week. The most common observed ADRs included injection site reaction, upper respiratory tract infection, cough, and pyrexia (observed in >15% of patients treated). Other less common ADRs included contusion, arthralgia, diarrhea, vomiting, and abdominal pain. Several clinical studies to assess the safety and tolerability of Viltolarsen are under recruiting status, including NCT 04956289 (phase 2, 20 participants, to be completed in May 2024), NCT04060199 (phase 3, 74 participants, to be completed in November 2024), NCT04768062 (phase 3, 74 participants, to be

DMD-MR-2021-000418

completed in June 2026), and NCT04687020 (phase 4, 16 participants, to be completed in November 2031).

A warning and precaution for renal toxicity has been placed in the FDA drug label of Viltolarsen. Although kidney toxicity was not directly observed in the clinical studies, preclinical studies reported potential risk of renal toxicity. Viltolarsen was administered to juvenile male mice via subcutaneous injection on postnatal day 7 and intravenous weekly injections from postnatal day 14 to 70. It was administered at doses of 0, 15, 60, 240 and 1,200 mg/kg. Renal toxicity was observed at the highest dose. In mice at the 240 and 1,200 mg/kg doses, there was a dose-dependent increase in deaths caused by renal toxicity, as well as a dose-dependent incidence and severity of renal tubular effects, such as degeneration. Reduced body weight and delayed sexual maturation were also observed at the highest dose. At 60 mg/kg, no renal toxicity was observed, and plasma exposure was similar to that in humans at 80 mg/kg. Long-term animal studies to assess the carcinogenic potential of Viltolarsen were not conducted. There were also no adequate data on the developmental risk associated with the use of Viltolarsen in pregnant women. In addition, no adverse effect on embryofetal development was observed in animal studies. Although renal toxicity was not observed in clinical studies, renal function should be monitored in patients taking Viltolarsen since renal toxicity and fatal glomerulonephritis were reported in preclinical studies.

The post-market data are not available for viltolarsen due to the short length of time since its approval for use.

Casimersen (Amondys 45)

DMD-MR-2021-000418

Developed by Sarepta Therapeutic Inc., Casimersen, is the newest FDA-approved ASO drug, which belongs to the phosphorodiamidate morpholino oligomer subclass. The FDA approved the drug in February 25, 2021 (Shirley, 2021) and its primary function is helping people who are suffering from DMD with a confirmed mutation on exon 45 of the *DMD* gene. Casimersen is designed to bind pre-mRNA of exon 45 of the DMD pre-mRNA, leading to skipping of the exon during mRNA processing. Therefore, patients can produce shorter functional dystrophin. Casimersen is approved under the provisions of accelerated approval regulations based on its capacity to increase dystrophin production in skeletal muscle of patients with DMD. The clinically recommended dose of Casimersen 45 is 30 mg/kg once weekly in 35 to 60 min infusion intravenously with 0.2-micron filter.

ADRs were reported in an initiated phase 1 clinical trial with 12 participants at therapeutic escalating doses (NCT02530905). The clinical trial was a multicenter, randomized, double-blind, placebo-controlled, dose-titration study (Wagner et al., 2021). The recruited patients were all males, age 7-21 years, who were diagnosed with DMD and confirmed genetic mutation amenable for treatment involving ASO-induced exon 45 skipping. The patients were randomized 2:1 to receive ascending doses of Casimersen at 4, 10, 20, and 30 mg/kg or placebo once a week through IV infusion for 12 weeks. After that, all patients received Casimersen at 30 mg/kg in an open-label extension period for up to 132 weeks. The common ADRs observed (>20% patients) included upper respiratory tract infections, cough, pyrexia, headache, arthralgia, and oropharyngeal pain. In addition to the observed common ADRs, other less common ADRs (<10% patients) included ear pain, nausea, ear infection, post-traumatic pain,

DMD-MR-2021-000418

dizziness, and light-headedness. An extension study to evaluate Casimersen in patients with DMD is underway with an estimated enrollment of 260 participants (NCT03532542), which is expected to be completed in 2026. Another double-blind, placebo-controlled, multi-center study to evaluate the efficacy and safety of Casimersen is also ongoing (so called ESSENCE, NCT02500381), which has 222 estimated participants and is expected to be completed in 2024.

A warning and precaution for kidney toxicity has been placed in the FDA-approved drug label of Casimersen. Although kidney toxicity was not shown in the clinical trial, kidney toxicity was reported in preclinical studies, including potentially fatal glomerulonephritis. In the preclinical studies, IV administration to juvenile male rats led to renal tubular degeneration and necrosis at the tested doses. Therefore, kidney function should be monitored when patients are treated with Casimersen. Meanwhile, serum cystatin C, glomerular filtration rate, urine dipstick, and urine protein-to-creatinine ratio are recommended to be measured before starting Amondys 45 in the drug label. Further, patients with renal dysfunction are recommended to be closely monitored during treatment with Casimersen.

Since the drug received FDA approval very recently, the post-market safety data is not available. To evaluate ADRs and toxicity of Casimersen, more data and information must be collected and/or generated.

Drug-specific and Class-wide ADRs and Toxicities among All FDA-approved ASO Drugs

DMD-MR-2021-000418

Due to different properties of absorption, distribution, metabolism, and excretion of each ASO drug involving different target organs and tissues, some specific ADRs are unique to particular ASO drugs (Table 2). Just to name a few, ocular ADRs are mainly found in Fomivirsen and Pegaptanib, which are directly administered into the eyes by intravitreal injection. On the other hand, although each FDA-approved ASO drug has a unique oligonucleotide sequence, they are also administered by different routes of delivery to target different organs and tissues. Some ADRs are found to occur across the spectrum of ASO drugs in general, which are also summarized in Table 2. A common ADR are application site reactions, which are likely caused by injection. Headache, pyrexia, and fever may reflect ADRs that are mediated by immune responses to the ASO drugs. Cough may be caused by respiratory infection, while vomiting and nausea are common gastrointestinal ADRs to the ASO drugs. Some common ADRs, such as headache, fever, flu-like symptoms, abdominal pain, nausea, and fatigue can be treated easily with non-prescriptions drugs.

The major safety concerns for the FDA-approved ASO drugs are severe hepatotoxicity, renal toxicity, and hypersensitivity reactions, which are commonly manifested with several ASO drugs (Figure 1 and Table 3). In particular, kidney toxicity has been indicated in Nusinersen, Inotersen, Golodirsen, Viltolarsen, and Casimersen, which are the ASO drugs developed for treatment of muscular dystrophy. Monitoring kidney function is critical while under treatment with these ASO drugs. Hypersensitivity reactions are listed as a precaution in Defibrotide, Inotersen, Eteplirsen, and Golodirsen. If a hypersensitivity reaction occurs, patients should carefully follow the institute appropriate medical treatment, and as a last resource or in cases of severe

DMD-MR-2021-000418

manifestation of these adverse reactions, discontinuing treatment is recommended. Risk of hepatotoxicity with elevation of ALT and AST and hepatic steatosis has been indicated in Mipomersen and Inotersen, which are the ASO drugs targeting hepatic gene expression. Because of severity of hepatotoxicity, Mipomersen has been discontinued from the market in 2019. Inotersen has numerous warnings and precautions on its black box label. Because of the risks of thrombocytopenia and glomerulonephritis, Inotersen is only available through a restricted distribution program.

Potential Mechanisms of the Common ADRs and Toxicity of the ASO Drugs

Although each ASO drug has a specific mechanism and a unique panoply of ADRs and toxicity, they also share some common mechanisms of toxicity as a class of drugs. ASO can cause toxicity via two possible mechanisms, via either hybridization dependent off-target effects or RNase-H1 dependent reduction in unintended off-target RNAs (Burel et al., 2016). The hybridization dependent off-target effects can occur due to the binding of ASOs to complementary sites of non-selective RNAs that have a homologous sequence to the intended target RNA (Kamola et al., 2017). Due to their various chemical functional groups, ASOs may produce non-specific interactions with plasma proteins, undesired organ-targeting, and possible binding with other small-molecule-based drugs that can result in toxicity. Studies have shown that hepatotoxicity of ASOs is mediated by the RNase-H1-dependent reduction in unintended off-target RNAs in mice (Burel et al., 2016).

The mechanism of liver toxicity of Mipomersen has been linked to the potential of the drug to induced endoplasmic reticulum stress (Hashemi et al., 2014). The induced

DMD-MR-2021-000418

stress impairs very low-density lipoprotein (VLDL) assembly via proteasomal and non-proteasomal pathways, which might induce a reduction of apo B mRNA levels, therefore causing drug-induced reduction of VLDL, which increases the risk of hepatic steatosis (Bell et al., 2011; Rinaldi and Wood, 2018). Preclinical studies established that hepatic Kupffer cells and renal proximal tubular epithelium receive the highest distribution of Mipomersen. The liver showed increased incidence of minimal cytoplasmic vacuolation at high exposure levels of Mipomersen in addition to proinflammatory signals. In the kidneys, minimal slight degeneration of the tubular epithelium was observed in monkeys.

The possible mechanism of kidney toxicity of the ASO drugs is the accumulation of the oligonucleotide within the lysosomes of the proximal tubules, resulting in physiological perturbation of tubular absorptive capacity and increased tubular proteinuria {Frazier, 2015 #96}. This was mainly associated to ASO ionic interactions with extracellular, cell-surface, or intracellular proteins. Repeated dosing results in ASOs accumulation in these tissues, which reaches steady state levels and the degree of toxicity then depends on the concentrations and inherent potency of the accumulated ASO. At high doses, the prominence of granules in epithelial cells correlates with an increased incidence or severity of degradation in the kidney; this is mostly a result of lysosomal breakdown.

Current Challenges, Knowledge Gaps, and Perspective on Future Directions

DMD-MR-2021-000418

Several FDA-approved ASO drugs target rare diseases. A limitation of the safety data is the very limited number of patients included in clinical trials, which is understandable given the low frequency of these diseases. However, common and less common ADRs and severe toxicity have emerged for this class of drugs. The listed ADRs, warnings, and precautions in the FDA-approved drug labels are supported, in part, by preclinical studies. Although some of the ADRs have been observed in clinical studies, they may not be entirely reflective of real-world conditions when ASO drugs are used to target diseases at a worldwide scale. Hence, it is critical to further extend the observations from clinical trials to examine patient samples in the post-marketing real-world settings to validate the findings on ADRs and toxicity from the pre-clinical and clinical trials in pre-marketing settings.

Another current challenge and knowledge gap is to understand inclusive mechanisms causing hepatotoxicity, renal toxicity, and hypersensitivity reactions by the ASO drugs. Although some speculations on mechanisms for hepatotoxicity and renal toxicity have been proposed, the precise molecular and biochemical events at the genetic level involving development of toxicity are unclear. Numerous questions need to be urgently addressed considering the rapid pace of discovery and development of ASO drugs. Going forward, the following questions should be addressed: What are the distinct features of ASO drugs-induced toxicity when compared to other small molecule-based drugs? What are the underlying molecular events occurring during ASO drug-induced toxicity? What is the capacity for liver and kidney to repair damage and/or develop tolerance to ASO drug-induced toxicity? What pharmacological interventions can be applied to mitigate the risk of ASO drug-induced toxicity? Does ASO drug-

DMD-MR-2021-000418

induced toxicity impairs function of the liver and kidneys as it relates to absorption, distribution, metabolism, and excretion of other small molecule-based drugs, altering their therapeutic efficacy and ADRs (drug-drug interactions)? These questions can lead to numerous perspective future directions to be addressed pertinent to the success of ASO based drug discovery and development for treating myriads of devastating diseases.

DMD-MR-2021-000418

Author contributions

Participated in research design: *F.A., K.Z., R.P., H.W., T.P.R., R.B., X-b.Z., J.E.M.*

Conducted experiments:

Performed data analysis: *F.A., K.Z., R.P., H.W.*

Wrote or contributed to the writing of the manuscript: *F.A., K.Z., R.P., H.W., T.P.R., R.B., X-b.Z., J.E.M.*

DMD-MR-2021-000418

Footnotes.

This study was partly supported by the National Institutes of Health (NIH) National Institute of General Medical Sciences [Grant R35GM140862 to X.Z.] and the University of Connecticut Research Excellence Program [to J.E.M.].

The authors declare there are no actual or perceived conflicts of interest with the contents of this article.

DMD-MR-2021-000418

Figure legend

Fig. 1. Shared warnings of toxicity in multiple FDA-approved ASO drugs. Hypersensitive reactions, such as rash and urticaria, are shown in Defibrotide, Inotersen, Eteplirsen, and Golodirsen. Hepatotoxicity is indicated after treatment with Mipomersen and Inotersen. Kidney toxicity is shown with the treatment of Nusinersen, Inotersen, Golodirsen, Viltolarsen, and Casimersen.

DMD-MR-2021-000418

Table 1. FDA-approved ASO drugs between 1998 and 2021

ASO Drug	Brand name	Company for manufacture	Disease to be treated	Therapeutic target	FDA approval date
A. ASO drugs targeting on mRNAs and pre-mRNAs					
Fomivirsen	Vitravene	Isis Pharmaceuticals	Cytomegalovirus retinitis	Cytomegalovirus IE2 mRNA	NDA* 20-961 08/26/1998 [#]
Mipomersen	Kynamro	Genzyme	Homozygous familial hypercholesterolemia	Apolipoprotein B mRNA	NDA 203568 01/29/2013 [^]
Eteplirsen	Exondys 51	Sarepta Therapeutics	Duchenne muscular dystrophy	Exon 51 dystrophin pre-mRNA	NDA 206488 09/19/2016
Nusinersen	Spinraza	Biogen	Spinal muscular atrophy	Survival motor neuron-2 (SMN-2) mRNA	NDA 209531 12/23/2016
Inotersen	Tegsedi	Ionis Pharmaceuticals	Polyneuropathy of amyloidosis	Transthyretin (TTR) mRNA	NDA 211172 10/05/2018
Golodirsen	Vyondys 53	Sarepta Therapeutics	Duchenne muscular dystrophy	Exon 53 dystrophin pre-mRNA	NDA 211970 12/12/2019
Viltolarsen	Viltepso	Nippon Shinyaku Co	Duchenne muscular dystrophy	Exon 53 dystrophin pre-mRNA	NDA 212154 08/12/2020
Casimersen	Amondys 45	Sarepta Therapeutics	Duchenne muscular dystrophy	Exon 45 dystrophin pre-mRNA	NDA 213026 02/25/2021
B. ASO drugs targeting on proteins					

DMD-MR-2021-000418

Pegaptanib	Macugen	Eyetech Pharmaceuticals	Neovascular macular degeneration	Vascular endothelial growth factor	NDA 21- 756 12/17/2004
Defibrotide	Defitelio	Jazz Pharmaceuticals	Veno-occlusive disease	Multiple mechanisms of action	NDA 208114 03/30/2016

*NDA: new drug application number.

#: withdrew in the EU market in 2002 and in the US in 2006

^: withdrew in the US market in 2019.

DMD-MR-2021-000418

Table 2. Common ADRs of the FDA-approved ASO drugs

ASO Drug	Common ADRs	Shared common ADRs by multiple ASO drugs
Fomivirsen	Increased intraocular pressure and ocular inflammation	Injection site reactions Headache Pyrexia Fever Respiratory infection Cough Vomiting Nausea
Pegaptanib	Anterior chamber inflammation, blurred vision, cataract, conjunctival hemorrhage, corneal edema, eye discharge, eye irritation, eye pain, hypertension, increased intraocular pressure (IOP), ocular discomfort, punctate keratitis, reduced visual acuity, visual disturbance, vitreous floaters, and vitreous opacities	
Mipomersen	Injection site reactions and flu-like symptoms.	
Defibrotide	Hypotension, diarrhea, vomiting, nausea, and epistaxis.	
Eteplirsen	Lower respiratory infection, pyrexia, constipation, headache, vomiting, and back pain.	
Nusinersen	Injection site reactions, nausea, headache, fatigue, thrombocytopenia, and fever.	
Inotersen	Balance disorder, vomiting, and contact dermatitis.	
Golodirsen	Headaches, pyrexia, fall, abdominal pain, nasopharyngitis, cough, vomiting, and nausea.	
Viltolarsen	injection site reaction, upper respiratory tract infection, cough, and pyrexia.	
Casimersen	upper respiratory tract infections, cough, pyrexia, headache, arthralgia, and oropharyngeal pain.	

DMD-MR-2021-000418

Table 3. Warnings and precautions for each ASO drug in the FDA-approved drug labeling

ASO drug	FDA warnings and precautions	Shared warnings and precautions by multiple ASO drugs
Fomivirsen	Ocular inflammation Increased intraocular pressure	Hepatotoxicity Kidney toxicity Hypersensitive reactions
Pegaptanib	Endophthalmitis Increased intraocular pressure	
Mipomersen	Black box: Risk of hepatotoxicity <ul style="list-style-type: none"> • Elevation of ALT and AST • Hepatic steatosis 	
Defibrotide	Hemorrhage Hypersensitive reactions	
Nusinersen	Thrombocytopenia Coagulation abnormalities Renal toxicity	
Inotersen	Black box: <ul style="list-style-type: none"> • Thrombocytopenia • Glomerulonephritis and renal toxicity Elevation of ALT and AST Stroke and cervicocephalic arterial dissection Inflammatory and immune effects Hypersensitive reactions Reduced serum vitamin A	
Eteplirsen	Hypersensitive reactions	
Golodirsen	Renal toxicity Hypersensitive reactions	
Viltolarsen	Kidney toxicity	
Casimersen	Kidney toxicity	

DMD-MR-2021-000418

References

- Bell DA, Hooper AJ, and Burnett JR (2011) Mipomersen, an antisense apolipoprotein B synthesis inhibitor. *Expert Opin Investig Drugs* **20**:265-272.
- Benson MD, Waddington-Cruz M, Berk JL, Polydefkis M, Dyck PJ, Wang AK, Plante-Bordeneuve V, Barroso FA, Merlini G, Obici L, Scheinberg M, Brannagan TH, 3rd, Litchy WJ, Whelan C, Drachman BM, Adams D, Heitner SB, Conceicao I, Schmidt HH, Vita G, Campistol JM, Gamez J, Gorevic PD, Gane E, Shah AM, Solomon SD, Monia BP, Hughes SG, Kwoh TJ, McEvoy BW, Jung SW, Baker BF, Ackermann EJ, Gertz MA, and Coelho T (2018) Inotersen Treatment for Patients with Hereditary Transthyretin Amyloidosis. *N Engl J Med* **379**:22-31.
- Bertolin C, Querin G, Martinelli I, Pennuto M, Pegoraro E, and Soraru G (2019) Insights into the genetic epidemiology of spinal and bulbar muscular atrophy: prevalence estimation and multiple founder haplotypes in the Veneto Italian region. *Eur J Neurol* **26**:519-524.
- Beytia Mde L, Vry J, and Kirschner J (2012) Drug treatment of Duchenne muscular dystrophy: available evidence and perspectives. *Acta Myol* **31**:4-8.
- Bushby K, Finkel R, Birnkrant DJ, Case LE, Clemens PR, Cripe L, Kaul A, Kinnett K, McDonald C, Pandya S, Poysky J, Shapiro F, Tomezsko J, Constantin C, and Group DMDCW (2010) Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and psychosocial management. *Lancet Neurol* **9**:77-93.
- Choy M (2016) Pharmaceutical Approval Update. *P T* **41**:416-441.
- Claborn MK, Stevens DL, Walker CK, and Gildon BL (2019) Nusinersen: A Treatment for Spinal Muscular Atrophy. *Ann Pharmacother* **53**:61-69.
- Clemens PR, Rao VK, Connolly AM, Harper AD, Mah JK, Smith EC, McDonald CM, Zaidman CM, Morgenroth LP, Osaki H, Satou Y, Yamashita T, Hoffman EP, and Investigators CD (2020) Safety, Tolerability, and Efficacy of Viltolarsen in Boys With Duchenne Muscular Dystrophy Amenable to Exon 53 Skipping: A Phase 2 Randomized Clinical Trial. *JAMA Neurol* **77**:982-991.
- Corbacioglu S, Cesaro S, Faraci M, Valteau-Couanet D, Gruhn B, Rovelli A, Boelens JJ, Hewitt A, Schrum J, Schulz AS, Muller I, Stein J, Wynn R, Greil J, Sykora KW, Matthes-Martin S, Fuhrer M, O'Meara A, Toporski J, Sedlacek P, Schlegel PG, Ehlert K, Fasth A, Winiarski J, Arvidson J, Mauz-Korholz C, Ozsahin H, Schrauder A, Bader P, Massaro J, D'Agostino R, Hoyle M, Iacobelli M, Debatin KM, Peters C, and Dini G (2012) Defibrotide for prophylaxis of hepatic veno-occlusive disease in paediatric haemopoietic stem-cell transplantation: an open-label, phase 3, randomised controlled trial. *Lancet* **379**:1301-1309.
- Crisafulli S, Sultana J, Fontana A, Salvo F, Messina S, and Trifiro G (2020) Global epidemiology of Duchenne muscular dystrophy: an updated systematic review and meta-analysis. *Orphanet J Rare Dis* **15**:141.
- Crooke ST (2017) Molecular Mechanisms of Antisense Oligonucleotides. *Nucleic Acid Ther* **27**:70-77.
- Darras BT, Farrar MA, Mercuri E, Finkel RS, Foster R, Hughes SG, Bhan I, Farwell W, and Gheuens S (2019) An Integrated Safety Analysis of Infants and Children with Symptomatic Spinal Muscular Atrophy (SMA) Treated with Nusinersen in Seven Clinical Trials. *CNS Drugs* **33**:919-932.

DMD-MR-2021-000418

- de Smet MD, Meenken CJ, and van den Horn GJ (1999) Fomivirsen - a phosphorothioate oligonucleotide for the treatment of CMV retinitis. *Ocul Immunol Inflamm* **7**:189-198.
- Dhillon S (2020) Viltolarsen: First Approval. *Drugs* **80**:1027-1031.
- Dhuri K, Bechtold C, Quijano E, Pham H, Gupta A, Vikram A, and Bahal R (2020) Antisense Oligonucleotides: An Emerging Area in Drug Discovery and Development. *J Clin Med* **9**.
- Dixon DL, Sisson EM, Butler M, Higbea A, Muoio B, and Turner B (2014) Lomitapide and mipomersen: novel lipid-lowering agents for the management of familial hypercholesterolemia. *J Cardiovasc Nurs* **29**:E7-E12.
- Elangkovan N and Dickson G (2021) Gene Therapy for Duchenne Muscular Dystrophy. *J Neuromuscul Dis*.
- Frank DE, Schnell FJ, Akana C, El-Husayni SH, Desjardins CA, Morgan J, Charleston JS, Sardone V, Domingos J, Dickson G, Straub V, Guglieri M, Mercuri E, Servais L, Muntoni F, and Group S-NS (2020) Increased dystrophin production with golodirsen in patients with Duchenne muscular dystrophy. *Neurology* **94**:e2270-e2282.
- Frazier KS (2015) Antisense oligonucleotide therapies: the promise and the challenges from a toxicologic pathologist's perspective. *Toxicol Pathol* **43**:78-89.
- Gales L (2019) Tegsedi (Inotersen): An Antisense Oligonucleotide Approved for the Treatment of Adult Patients with Hereditary Transthyretin Amyloidosis. *Pharmaceuticals (Basel)* **12**.
- Group VISiONCT, Chakravarthy U, Adamis AP, Cunningham ET, Jr., Goldbaum M, Guyer DR, Katz B, and Patel M (2006) Year 2 efficacy results of 2 randomized controlled clinical trials of pegaptanib for neovascular age-related macular degeneration. *Ophthalmology* **113**:1508 e1501-1525.
- Gupta S (2015) LDL cholesterol, statins and PCSK 9 inhibitors. *Indian Heart J* **67**:419-424.
- Hair P, Cameron F, and McKeage K (2013) Mipomersen sodium: first global approval. *Drugs* **73**:487-493.
- Hashemi N, Odze RD, McGowan MP, Santos RD, Stroes ESG, and Cohen DE (2014) Liver histology during Mipomersen therapy for severe hypercholesterolemia. *J Clin Lipidol* **8**:606-611.
- Heo YA (2020) Golodirsen: First Approval. *Drugs* **80**:329-333.
- Hoy SM (2017) Nusinersen: First Global Approval. *Drugs* **77**:473-479.
- Ito M, Yamauchi A, Urano M, Kato T, Matsuo M, Nakashima K, and Saito K (2021) Epidemiological investigation of spinal muscular atrophy in Japan. *Brain Dev*.
- Jonas JB, Cheung CMG, and Panda-Jonas S (2017) Updates on the Epidemiology of Age-Related Macular Degeneration. *Asia Pac J Ophthalmol (Phila)* **6**:493-497.
- Katz B and Goldbaum M (2006) Macugen (pegaptanib sodium), a novel ocular therapeutic that targets vascular endothelial growth factor (VEGF). *Int Ophthalmol Clin* **46**:141-154.
- Keam SJ (2018) Inotersen: First Global Approval. *Drugs* **78**:1371-1376.
- Kessler-Icekson G, Schlesinger H, Leger JJ, Leger J, Braverman Y, and Binah O (1988) Ventricular myosin of the shrew *Crocidura russula*, correlation with contractile properties. *J Mol Cell Cardiol* **20**:1069-1073.
- Kornblum N, Ayyanar K, Benimetskaya L, Richardson P, Iacobelli M, and Stein CA (2006) Defibrotide, a polydisperse mixture of single-stranded phosphodiester oligonucleotides

DMD-MR-2021-000418

- with lifesaving activity in severe hepatic veno-occlusive disease: clinical outcomes and potential mechanisms of action. *Oligonucleotides* **16**:105-114.
- Lim KR, Maruyama R, and Yokota T (2017) Eteplirsen in the treatment of Duchenne muscular dystrophy. *Drug Des Devel Ther* **11**:533-545.
- Lui DTW, Lee ACH, and Tan KCB (2021) Management of Familial Hypercholesterolemia: Current Status and Future Perspectives. *J Endocr Soc* **5**:bvaa122.
- Luigetti M, Romano A, Di Paolantonio A, Bisogni G, and Sabatelli M (2020) Diagnosis and Treatment of Hereditary Transthyretin Amyloidosis (hATTR) Polyneuropathy: Current Perspectives on Improving Patient Care. *Ther Clin Risk Manag* **16**:109-123.
- Marwick C (1998) First "antisense" drug will treat CMV retinitis. *JAMA* **280**:871.
- Matsuo M (2021) Antisense Oligonucleotide-Mediated Exon-skipping Therapies: Precision Medicine Spreading from Duchenne Muscular Dystrophy. *JMA J* **4**:232-240.
- Mendell JR, Rodino-Klapac LR, Sahenk Z, Roush K, Bird L, Lowes LP, Alfano L, Gomez AM, Lewis S, Kota J, Malik V, Shontz K, Walker CM, Flanigan KM, Corridore M, Kean JR, Allen HD, Shilling C, Melia KR, Sazani P, Saoud JB, Kaye EM, and Eteplirsen Study G (2013) Eteplirsen for the treatment of Duchenne muscular dystrophy. *Ann Neurol* **74**:637-647.
- Mercuri E, Darras BT, Chiriboga CA, Day JW, Campbell C, Connolly AM, Iannaccone ST, Kirschner J, Kuntz NL, Saito K, Shieh PB, Tulinius M, Mazzone ES, Montes J, Bishop KM, Yang Q, Foster R, Gheuens S, Bennett CF, Farwell W, Schneider E, De Vivo DC, Finkel RS, and Group CS (2018) Nusinersen versus Sham Control in Later-Onset Spinal Muscular Atrophy. *N Engl J Med* **378**:625-635.
- Narayanan P, Curtis BR, Shen L, Schneider E, Tami JA, Paz S, Burel SA, Tai LJ, Machemer T, Kwoh TJ, Xia S, Shattil SJ, Witztum JL, Engelhardt JA, Henry SP, Monia BP, and Hughes SG (2020) Underlying Immune Disorder May Predispose Some Transthyretin Amyloidosis Subjects to Inotersen-Mediated Thrombocytopenia. *Nucleic Acid Ther* **30**:94-103.
- Nassir F, Rector RS, Hammoud GM, and Ibdah JA (2015) Pathogenesis and Prevention of Hepatic Steatosis. *Gastroenterol Hepatol (N Y)* **11**:167-175.
- Nowak JZ (2006) Age-related macular degeneration (AMD): pathogenesis and therapy. *Pharmacol Rep* **58**:353-363.
- Penedones A, Mendes D, Alves C, and Batel Marques F (2014) Safety monitoring of ophthalmic biologics: a systematic review of pre- and postmarketing safety data. *J Ocul Pharmacol Ther* **30**:729-751.
- Perry CM and Balfour JA (1999) Fomivirsen. *Drugs* **57**:375-380; discussion 381.
- Port AD, Orlin A, Kiss S, Patel S, D'Amico DJ, and Gupta MP (2017) Cytomegalovirus Retinitis: A Review. *J Ocul Pharmacol Ther* **33**:224-234.
- Reeskamp LF, Kastelein JJP, Moriarty PM, Duell PB, Catapano AL, Santos RD, and Ballantyne CM (2019) Safety and efficacy of mipomersen in patients with heterozygous familial hypercholesterolemia. *Atherosclerosis* **280**:109-117.
- Richardson PG, Grupp SA, Pagliuca A, Krishnan A, Ho VT, and Corbacioglu S (2017) Defibrotide for the treatment of hepatic veno-occlusive disease/sinusoidal obstruction syndrome with multiorgan failure. *Int J Hematol Oncol* **6**:75-93.
- Richardson PG, Riches ML, Kernan NA, Brochstein JA, Mineishi S, Termuhlen AM, Arai S, Grupp SA, Guinan EC, Martin PL, Steinbach G, Krishnan A, Nemecek ER, Giralt S,

DMD-MR-2021-000418

- Rodriguez T, Duerst R, Doyle J, Antin JH, Smith A, Lehmann L, Champlin R, Gillio A, Bajwa R, D'Agostino RB, Sr., Massaro J, Warren D, Miloslavsky M, Hume RL, Iacobelli M, Nejadnik B, Hannah AL, and Soiffer RJ (2016) Phase 3 trial of defibrotide for the treatment of severe veno-occlusive disease and multi-organ failure. *Blood* **127**:1656-1665.
- Rinaldi C and Wood MJA (2018) Antisense oligonucleotides: the next frontier for treatment of neurological disorders. *Nat Rev Neurol* **14**:9-21.
- Roberts TC, Langer R, and Wood MJA (2020) Advances in oligonucleotide drug delivery. *Nat Rev Drug Discov* **19**:673-694.
- Roehr B (1998) Fomivirsen approved for CMV retinitis. *J Int Assoc Physicians AIDS Care* **4**:14-16.
- Santos RD, Raal FJ, Catapano AL, Witztum JL, Steinhagen-Thiessen E, and Tsimikas S (2015) Mipomersen, an antisense oligonucleotide to apolipoprotein B-100, reduces lipoprotein(a) in various populations with hypercholesterolemia: results of 4 phase III trials. *Arterioscler Thromb Vasc Biol* **35**:689-699.
- Schmidt HH, Waddington-Cruz M, Botteman MF, Carter JA, Chopra AS, Hopps M, Stewart M, Fallet S, and Amass L (2018) Estimating the global prevalence of transthyretin familial amyloid polyneuropathy. *Muscle Nerve* **57**:829-837.
- Shirley M (2021) Casimersen: First Approval. *Drugs* **81**:875-879.
- Son HW and Yokota T (2018) Recent Advances and Clinical Applications of Exon Inclusion for Spinal Muscular Atrophy. *Methods Mol Biol* **1828**:57-68.
- Stephenson ML and Zamecnik PC (1978) Inhibition of Rous sarcoma viral RNA translation by a specific oligodeoxyribonucleotide. *Proc Natl Acad Sci U S A* **75**:285-288.
- Syed YY (2016) Eteplirsen: First Global Approval. *Drugs* **76**:1699-1704.
- Thomas GS, Cromwell WC, Ali S, Chin W, Flaim JD, and Davidson M (2013) Mipomersen, an apolipoprotein B synthesis inhibitor, reduces atherogenic lipoproteins in patients with severe hypercholesterolemia at high cardiovascular risk: a randomized, double-blind, placebo-controlled trial. *J Am Coll Cardiol* **62**:2178-2184.
- Tobin KA (2006) Macugen treatment for wet age-related macular degeneration. *Insight* **31**:11-14.
- Vitravene Study G (2002) Safety of intravitreal fomivirsen for treatment of cytomegalovirus retinitis in patients with AIDS. *Am J Ophthalmol* **133**:484-498.
- Wagner KR, Kuntz NL, Koenig E, East L, Upadhyay S, Han B, and Shieh PB (2021) Safety, tolerability, and pharmacokinetics of casimersen in patients with Duchenne muscular dystrophy amenable to exon 45 skipping: A randomized, double-blind, placebo-controlled, dose-titration trial. *Muscle Nerve* **64**:285-292.
- Wong E and Goldberg T (2014) Mipomersen (kynamro): a novel antisense oligonucleotide inhibitor for the management of homozygous familial hypercholesterolemia. *P T* **39**:119-122.
- Zhang MM, Bahal R, Rasmussen TP, Manautou JE, and Zhong XB (2021) The growth of siRNA-based therapeutics: Updated clinical studies. *Biochem Pharmacol* **189**:114432.

Fig. 1

