

SUPPLEMENTAL FIGURES

Evaluation of Normothermic Machine Perfusion of Porcine Livers as a Novel Preclinical Model to Predict Biliary Clearance and Transporter-mediated Drug-Drug Interactions using Statins

L.J. Stevens^{1,2}, A.Z.X. Zhu³, P.P. Chothe⁴, S.K. Chowdhury⁴, J. M. Donkers², W.H.J. Vaes², C.A.J. Knibbe⁵, I.P.J. Alwayn¹ and E. van de Steeg²

¹ Department of Surgery, Leiden University Medical Centre (LUMC) Transplant Center, Leiden, the Netherlands

² The Netherlands Organization for Applied Scientific Research (TNO), Zeist, the Netherlands

³ Quantitative Solutions, Takeda Pharmaceutical International, Cambridge, MA, USA

⁴ Department of Drug Metabolism & Pharmacokinetics, Takeda Pharmaceuticals International, Cambridge, MA, USA

⁵ Division of Systems Biomedicine and Pharmacology, Leiden Academic Centre for Drug Research (LACDR), Leiden, the Netherlands & Department of Clinical Pharmacy, St. Antonius Hospital Nieuwegein and Utrecht, the Netherlands

Drug Metabolism and disposition

DMD-AR-2021-000521

Table S3. The Multiple Reaction Monitoring Transition (MRM) of Compounds

Compound	MRM Transition (m/z)
Atorvastatin	559.30→466.00
Atorvastatin lactone	541.30→448.10
2-hydroxy atorvastatin	575.40→440.00
2-hydroxy atorvastatin lactone	557.30→448.10
4-hydroxy atorvastatin	575.30→440.20
4-hydroxy atorvastatin lactone	557.40→448.10
Pitavastatin	422.20→318.10
Rosuvastatin	482.10→258.20
Rifampicin	823.50→399.20
Glyburide (internal standard)	494.20→369.10
Carbamazepine (internal standard)	237.10→194.10
Rifampicin_d3 (internal standard)	826.50→794.70
Chrysin (internal standard)	255.10→153.00