

DMD #50278

Which Metabolites Circulate?

Cho-Ming Loi, Dennis A. Smith and Deepak Dalvie

Pharmacokinetics, Dynamics, and Metabolism, Pfizer Worldwide Research and Development,
San Diego, CA 92121 (*CML, DD*); Honary Professor, Department of Chemistry, University of
Capetown, South Africa, Visiting Professor, Institute of Translational Medicine, University of
Liverpool, UK (*DAS*)

DMD #50278

Running Title: Which Metabolites Circulate?

Corresponding Author: Cho-Ming Loi

Address: Pfizer, Inc., 10646 Science Center Drive, San Diego, CA 92121

Telephone: (858)622-7361

E-Mail address: Cho-Ming.Loi@Pfizer.com

Number of Text Pages: 38

Number of Tables: 7

Number of Figures: 7

Number of word in the Abstract: 247

DMD #50278

Non-standard Abbreviations:

ADME: Absorption, distribution, metabolism, excretion

AUC: Area-under-the concentration time curve

AUC_m: Area-under-the concentration time curve for a metabolite

BSEP: Bile salt export pump

CL_f: Formation clearance of metabolite

CL_m: Elimination clearance of metabolite

CL_{parent}: Clearance of parent drug

CL_s = Total systemic clearance

CL_{R, U} = Unbound renal clearance

CL_{NR, U} = Unbound non-renal (hepato-biliary) clearance

CYP: Cytochrome P450

DDI: Drug-drug interactions

EM: Extensive metabolizer

EMA: European Medicines Agency

F_a: Fraction of dose absorbed after oral administration

FDA: Food and Drug Administration

F_{h,m}: Systemic availability of the metabolite

f_m: Fraction of dose converted to a given metabolite from the parent

f_p: Lipoidal permeability

MIST: Drug metabolites in safety testing

M/P ratio: Relative abundance of metabolite to parent based on the ratio of plasma/serum AUC values

MRP: Multidrug resistance protein

MRP2: Multidrug resistance protein 2

DMD #50278

MRP3: Multidrug resistance protein 3

PAPS: 3'-phosphoadenosine 5'-phosphosulfate

PM: Poor metabolizer

tPSA: Topological polar surface area

UDPGA: Uridine 5'-diphospho-glucuronic acid

UGT: Uridine diphosphate glucuronosyltransferase

UM: Ultra rapid metabolizer

DMD #50278

Abstract

Characterization of the circulating metabolites for a new chemical entity in humans is essential for safety assessment, an understanding of their contributions to pharmacologic activities, and their potential involvement in drug-drug interactions (DDI). This review examines the abundance of metabolites relative to the total parent drug (M/P ratio) from 125 drugs in relation to their structural and physicochemical characteristics, lipoidal permeability, protein binding, and fractional formation from parent (f_m). Our analysis suggests that f_m is the major determinant of total drug M/P ratio for amine, alcohol, N- and S-oxide, and carboxylic acid metabolites. Passage from the hepatocyte to systemic circulation does not appear to be limiting owing to the vast majority of metabolites formed being relatively lipid permeable. In some cases, active transport plays an important role in this process (e.g., carboxylic acid metabolites). Differences in total parent drug clearance and metabolite clearance are attenuated by the reduction in lipophilicity introduced by the metabolic step and resultant compensatory changes in unbound clearance and protein binding. A small sub-class of these drugs (e.g. terfenadine) are unintentional pro-drugs with very high parent drug clearance resulting in very high M/P ratios. In contrast, arecol metabolites show a more complex relationship with f_m due largely to the new metabolic routes (conjugation) available to the metabolite compared to the parent drug molecule. For these metabolites, a more thorough understanding of the elimination clearance of the metabolite is critical to discern the likelihood of whether the phenol will constitute a major circulating metabolite.

DMD #50278

1. Introduction

An early understanding of the key metabolites for a new chemical entity is important in drug discovery and development. At the lead optimization stage, this can help to support medicinal chemistry efforts in designing more metabolically stable candidates via identification of soft spots in lead candidates that are rapidly cleared (Lin and Lu, 1997; Baillie, 2008). From a development perspective, characterization of metabolites provides a systematic understanding of the metabolic pathways of a drug and therefore the enzymes that are involved in their formation. This can aid in the proper design of drug-drug interaction (DDI) studies to evaluate the impact of inhibition or induction of the metabolic enzyme(s) contributing to the major pathway of clearance on systemic drug exposure. Assessment of circulating metabolites in humans also contributes to an overall understanding of their potential impact on the safety and efficacy of a drug. From a safety point of view, an effort towards gathering knowledge of the *in vivo* metabolic profile of a drug in humans has gained more importance since the publication of position papers on drug metabolites in safety testing (MIST) as well as recently issued ICH Guidance M3(R2) on nonclinical safety studies for the conduct of human clinical trials and marketing authorization for pharmaceuticals (FDA guidance; <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM073246.pdf>); (Davis-Bruno and Atrakchi, 2006; Nedderman et al., 2011). This guidance recommends that nonclinical safety evaluation is warranted if the exposure of a human circulating metabolite is observed at greater than 10% of the total drug-related exposure and at significantly greater levels in humans than the maximum exposure seen in the toxicity studies. Assessment of circulating metabolites in humans is also essential for a comprehensive understanding of their contributions to pharmacologic activities. Structural alteration of drugs

DMD #50278

via metabolism can sometimes lead to products that contribute to the therapeutic effect of the drug (on target activity) or cause an undesirable effect as a consequence of off-target activity (Fura, 2006). More recently, efforts have been directed in evaluating the potential involvement of metabolites in inhibitory DDI. Several clinically important drugs (especially clinically recognized CYP inhibitors) have been shown to have circulating metabolites that act as potent inhibitors of CYP enzymes and therefore contribute to in vivo DDIs (Isoherranen et al., 2009; Yeung et al., 2011). The most recent guidance by the European Medicines Agency (EMA) and US Food and Drug Administration (FDA) also suggest evaluation of potential inhibitory effect of metabolites on the common drug metabolizing enzymes. These guidances recommend that circulating metabolite with an AUC of $\geq 25\%$ relative to parent AUC (Draft FDA Guidance; <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM292362.pdf>) and phase 1 metabolites with an AUC both larger than one fourth of the AUC of parent drug and larger than 10% of the drug-related exposure (EMA Guidance; http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/07/WC500129606.pdf) should be investigated for their inhibitory potential. These recommendations necessitate characterization and quantification of the metabolites in plasma early on to allow a full assessment of their contributions to the pharmacological or toxicological effects and DDI potential.

In most cases, principal metabolites are identified by performing radiolabeled ADME studies where ^{14}C or ^3H analogs of the drugs are administered to humans (Roffey et al., 2007; Isin et al., 2012; Penner et al., 2012). The biological matrices collected from this study are then analyzed to yield a quantitative profile of metabolites in circulation or in the excreta. While the

DMD #50278

abundance of metabolites in the excreta provides a means of assessing the clearance pathways and the fraction of the drug metabolized via a given pathway (f_m), the exposure of metabolites in plasma obtained from this study serves as a starting point for prioritizing their further evaluation. The major metabolites are then synthesized and evaluated for pharmacological on-target and off-target interactions that may result in unpredicted toxicity or DDI. However, radiolabeled ADME studies in humans are expensive and resource-intensive and hence, not routinely feasible in early development. This can hinder the knowledge about relevant circulating plasma metabolites and hence their evaluation in a timely manner. Hence, until definitive data are obtained from human studies, one may need to rely on the *in vitro* systems for an assessment of potential metabolites that are generated. Human-derived reagents such as the liver microsomes, hepatocytes or S9 fractions, are the commonly used *in vitro* systems to study human drug metabolism and identify the principal metabolites, preferably prior to the human radiolabel ADME study (Dalvie et al., 2009). A comprehensive analysis of these *in vitro* systems by Dalvie and co-workers provide sufficient confidence in using these systems to reliably produce primary human metabolites (Dalvie et al., 2009). Although primary human metabolites can be predicted readily by *in vitro* systems, it is important to understand which metabolites formed in these systems are likely to circulate *in vivo*. This can help in prioritizing the synthesis of important metabolites for further characterization with respect to their pharmacological activity and/or their role as inhibitors or inducers of important drug metabolizing enzymes.

Smith and Dalvie have recently attempted to analyze the influence of physicochemical properties on the metabolites that circulate. This review showed that metabolites with high lipoidal permeability will passively diffuse from liver to plasma and circulate (Smith and Dalvie,

DMD #50278

2012). The aim of the current analysis was to extend this investigation and explore how the structures of the metabolite (e.g., amines, alcohols, etc.) and their associated physicochemical properties, as well as the fraction of dose converted to a given metabolite from the parent (f_m) may influence their abundance in circulation.

We conducted a literature review to collect data on the plasma or serum area-under-the-concentration time curve (AUC) values from 125 drugs and their metabolites after oral administration of single or multiple doses of the parent, and then determine the relative abundance of metabolite to parent based on the ratio of plasma/serum AUC values (M/P ratio). Although several secondary and tertiary metabolites were detected circulating for several drugs, no attempt was made to classify and categorize these metabolites separately. In the vast majority of cases, the ratio of plasma or serum AUC of the metabolite to the corresponding parent was used, although in a few cases (as indicated by the asterisk in Tables 1-6), single time-point concentrations were used due to paucity of the overall AUC data. In a few cases (clozapine, dextromethorphan, dothiepin, flecainide, and zolpidone) where plasma concentration-time profiles were presented in the publication but the AUC values were not calculated, the data were captured using the software DigitizeIt version 1.5 (Eden Prairie, MN) and the AUC values were calculated by noncompartmental analysis using WinNonlin version 5.2 (Pharsight Corp., Cary, NC). The M/P ratio for these metabolites was then matched with the structure and physicochemical properties of the corresponding parent and metabolite pair to discern any possible relationship between the two. The physicochemical properties for the parent and metabolites were calculated using the ACD software version 12.01 (ACD Labs, Advanced Chemistry Development Inc., Toronto, Ontario, Canada), and the chemical structures of the

DMD #50278

parent and corresponding metabolites are shown in the Supplementary file (SuppTable 1-6). In addition, where data exists, information on the f_m value was also estimated from the published human radiolabeled ADME studies. The fraction of dose for the metabolite of interest and its sequential metabolic products recovered in excreta (both urine and feces) was assumed to be representative of the in vivo f_m value. An analysis was conducted in an attempt to identify parameter values for the physicochemical properties and f_m that are associated with metabolites considered to have met the criteria for follow-up DDI evaluations according to the regulatory guidance.

2. Theoretical Considerations of Determining In Vivo Metabolite Exposure

Theoretical models related to metabolite kinetics after oral or intravenous administration of the parent drug have been described extensively (Houston, 1982; Lutz et al., 2010) and will only be discussed briefly in this review. In vivo, a metabolite can display either a formation-rate limited or an elimination-rate limited kinetics. In the former case, the half-life of the terminal elimination phase of a metabolite plasma/serum concentration versus time curve will be equal to that of the parent drug; while in the latter case, the half-life of the terminal elimination phase of the metabolite will be greater than that of the parent. Most assessment of circulating concentrations of drugs and their metabolites are made without reference to protein binding, certainly historically and also at the initial profiling and identification stage of drug discovery and development. Regulatory guidelines reflect this in that the identification of “major” metabolite rely on total concentration compared to parent or drug-derived moieties, although there is recognition that certain metabolites may have very different binding characteristics to their parent. In order to understand the dynamics of metabolites, protein binding must be

DMD #50278

measured to derive unbound concentrations. This is reinforced by the tendency for a reduced lipophilicity for the metabolite, which may increase the fraction unbound and lowers unbound clearance leading to disproportionate concentrations of unbound metabolite compared to unbound concentrations of parent. Unfortunately, protein binding data is lacking in many cases to provide more sophisticated analysis. In terms of the effects of changes in chemical structures and how they may impact the ratio of total drug to metabolite, the influence of protein binding may be attenuated. Most metabolic steps typically result in reduced lipophilicity, which will tend to decrease protein binding (or increase fraction unbound) and intrinsic metabolic clearance (unless the metabolic step introduces a route by which the parent cannot be metabolized), thereby attenuating the effect on the difference between parent and metabolite clearance (See Section 9.3). Similarly, the lowered lipophilicity will tend to decrease tissue or membrane affinity, which will be compensated for by the increased free fraction in plasma, thereby attenuating the effects on total volume of distribution. Thus, the ratio of total parent to metabolite is probably more sensitive to other factors as detailed below.

The systemic exposure (total, bound plus free), as represented by the AUC of a metabolite (AUC_m), will be dependent on the dose of the parent drug (D), the fraction of dose absorbed (after oral administration, F_a), the fraction of the absorbed drug dose that is converted to the metabolite (f_m), the systemic availability of the metabolite ($F_{h,m}$), and the clearance of the metabolite (CL_m), i.e., $AUC_m = (f_m \cdot F_{h,m} \cdot F_a \cdot D) / CL_m$ (Houston, 1982). As illustrated in Figure 1, the systemic availability of a metabolite can be conceptualized as the net fraction of the formed metabolite partitioning back into the circulation. Physiologically, this reflects the permeability rate of the metabolite across the sinusoid for passive diffusion (lipoidal

DMD #50278

permeability, f_p) and the effect of sinusoidal efflux transporters (active transport) compared to the rate of further processing of the metabolite by metabolism within the hepatocyte or clearance directly from the hepatocyte or by canalicular transporters (biliary excretion). Lipoidal permeability decreases with decreasing lipophilicity and increasing polar surface area (tPSA). Unless the metabolite is a substrate for sinusoidal transporters, it is possible that its residence time in the liver will be increased by metabolism favoring further processing and attenuating systemic exposure.

In vivo, after oral dosing, the ratio of AUC for the metabolite to parent (M/P ratio) will be determined by $f_m \cdot F_{h,m} \cdot CL_{parent} / CL_m \cdot F_h$ (where F_h is the fraction of absorbed parent drug dose that escape the liver) since dose (parent) and F_a are the same. For a drug cleared by metabolism to a single metabolite which diffuses completely from the hepatocyte to systemic circulation (e.g., a metabolite with high lipoidal permeability), this simplifies to $CL_{parent} / CL_m \cdot F_h$, assuming linear processes. For a drug that undergoes metabolism to more than one metabolite and assuming complete diffusion of the metabolite(s) to systemic circulation, the M/P ratio is expressed as $f_m \cdot CL_{parent} / CL_m \cdot F_h$, which indicates that the formation and elimination clearance (CL_f and CL_m , respectively) of the metabolite are major determinants of its M/P ratio. The differences between CL_{parent} and CL_m likely reflect actual chemical-structural change(s) as a result of the metabolism of a more labile functional group to a more stable functional group rather than simple reduction in lipophilicity (see above discussion on protein binding). These principles for highly diffusible metabolites have been established for some time (Levy et al. 1983), and provide a useful conceptual framework to guide the assessment and interpretation of our analysis regarding the potential relationship between structural and physicochemical

DMD #50278

properties of the metabolites and their abundance in circulation relative to the parent drug based on ratio of AUC.

3. Secondary or Primary Amines as Metabolites

N-Dealkylation represents one of the most common biotransformation pathways for drugs containing secondary and tertiary amines. This oxidative pathway results in the formation of metabolites that contain primary or secondary amines or in some cases cleaved products as primary metabolites. In this dataset, 52 out of a total of 125 drugs (42%) were *N*-dealkylated as part of the overall clearance process. Of these 52 drugs, 40 (77%) had an amine metabolite with an abundance, based on the M/P ratio, of >0.25 in human plasma (Table 1). This trend is not surprising since these metabolites retain lipophilicity, as represented by cLogD, which is within the range of -1 to 5 and is optimal for passive diffusion (Figure 2). The physicochemical changes associated with *N*-dealkylation usually showed a small increase in tPSA of $\sim 9 \text{ \AA}$ and a reduction in cLogD of up to 1.0 log unit when simple alkyl substituents were cleaved (Table 1). However, when *N*-dealkylation reaction cleaved a molecule into smaller fragments, more pronounced changes in cLogD and tPSA were observed. For instance, cleavage of nefadozone to *m*-chlorophenylpiperazine (Figure 3) was accompanied by a reduction in cLogD of ~ 3 log units along with a decrease in tPSA of 36 \AA . Interestingly, some drugs in this dataset were exceptions to this trend in that high abundance of amine metabolites were present in circulation despite their high polarity (i.e., cLogD < -1). For example, metabolites of ranolazine, tramadol, zolmitriptan, and sildenafil (the cleaved piperazine metabolite) displayed cLogD values ranging from -1.11 to -2.97 (Table 1). Further, many cleaved metabolites such as 1-pyrimidinylpiperazine metabolite of buspirone and D617 or D620 metabolites of verapamil were also found circulating in human

DMD #50278

plasma with M/P ratios of >0.25 despite their high polarity. While the reason for this is not known, one possibility for the high circulating concentrations of these metabolites relative to the parent could be attributed to transporter-mediated uptake into the blood. These hypotheses will require further experimental evaluation for confirmation.

Our analysis also identified some amine metabolites that circulated in relatively low abundance despite having optimal cLogD values of between -1 and 5 (Table 1). Based on the conceptual framework discussed in Section 2, this was ascribed to either a low CL_f and/or a high CL_m for the metabolite. For instance, low circulating concentrations of *N*-desalkyl metabolites of azimilide (Riley et al., 2005), elzasonan (Kamel et al., 2010), imatinib (Gschwind et al., 2005), itraconazole (Isoherranen et al., 2004), olanzapine (Kassahun et al., 1997), propafenone (Kroemer et al., 1989), and repaglinide (van Heiningen et al., 1999) can be attributed to the relatively minor contribution of *N*-dealkylation to the overall clearance of these drugs. On the other hand, low circulating concentrations of *m*-chlorophenylpiperazine (Figure 3) may be ascribed to a high metabolic rate of this metabolite to the corresponding hydroxylated chlorophenylpiperazine (i.e., high CL_m of this metabolite) (von Moltke et al., 1999). Ranitidine represents an interesting example in that its *N*-desmethyl metabolite showed both high polarity (cLogD of -2.22) and low f_m (0.01) for this metabolic pathway (Table 1). Thus, it is not surprising that the abundance of this metabolite in circulation is negligible.

The differences in the M/P ratio of *N*-deisopropylidelavirdine (Figure 3) following single and multiple dose administration of delavirdine are also informative to our analysis. After oral administration of a single dose, the M/P ratio of *N*-deisopropylidelavirdine to parent drug was ~ 1.1 (Morse et al., 1997). However, after multiple dosing, this ratio was decreased to ~ 0.12 to

DMD #50278

0.15 (Borin et al., 1997a; Borin et al., 1997b). This disconnect in M/P ratios after single and multiple doses could be possibly attributed to decreased CYP3A-mediated formation of *N*-desisopropyl metabolite of delavirdine leading to the lower concentration in the systemic circulation. *In vitro*, formation of the *N*-desisopropyl delavirdine is mediated by CYP3A (Voorman et al., 1998a; Voorman et al., 1998b). However, delavirdine has also been shown to inactivate CYP3A (Voorman et al., 1998b), which may account for the reduction in oral clearance of delavirdine and hence, a decrease in formation of *N*-desisopropyl delavirdine with repeated dose administration.

Even though most aliphatic tertiary amines can undergo secondary metabolism to primary amines via the corresponding secondary amines, only two drugs, namely sibutramine and tamoxifen, yielded primary amines (didesethylsibutramine and didesethyltamoxifen, respectively, Figure 3) that were circulating in greater than 25% of the parent AUC (Kim et al., 2009; Yeung et al., 2011). Moreover, other drugs (Table 1) that theoretically could be converted to primary amines (based on their structures and lipophilicity) showed very little or no didesalkyl metabolites in circulation. For instance, the M/P ratios for the didesalkyl metabolite of amiodarone (McDonald et al., 2012) or citalopram (Herrlin et al., 2003) were less than 0.1 despite having similar cLogD values to that of the secondary amine metabolites. It is likely that this disconnect can be ascribed to a low fraction of the secondary amine being metabolized to the corresponding primary amine and is indicative of increased metabolic stability of the secondary amine with respect to the subsequent *N*-demethylation/*N*-dealkylation pathway, compared to the parent tertiary amine. Alternatively, the structural change from a tertiary amine to a secondary amine may impart a different biotransformation pathway such as hydroxylation on the secondary

DMD #50278

amine (e.g., desipramine to hydroxydesipramine and nortriptyline to hydroxynortriptyline, Figure 3) rather than formation of a primary amine.

Overall, this dataset suggests that the amine metabolites can be present in circulation that exceeds the threshold of 25% parent AUC over a wide range of cLogD values (Figure 2). Thus, the f_m or CL_m appear to be the more important factors in determining the abundance of the amine metabolites in circulation. For those amine metabolites that have cLogD values below -1, they may still be present in circulation in abundance that exceeds the M/P ratio of 0.25, possibly related to transport-mediated processes. Hence, assessment of f_m , CL_m , cLogD, and transport properties will provide a useful guide to estimate the potential for the amine metabolites to circulate in abundance that may exceed the threshold of 0.25 relative to that for the parent AUC.

4. Alcohols and Ketones as Metabolites

In this dataset, oxygenation of compounds constitutes the second most common biotransformation process. The corresponding hydroxylated metabolites could be formed either by oxygen insertion into the alkyl or alicyclic and aromatic groups or via *O*-dealkylation of aliphatic or aromatic ethers. Analysis of the data set for hydroxylation of aliphatic or alicyclic moieties revealed that of the 125 drugs examined, 40 drugs were shown to yield metabolites with this modification (Table 2A), some of which were further oxidized to ketones (Table 2B). Examination of physicochemical properties of these metabolites suggested that formation of an aliphatic or alicyclic alcohol was associated with an increase in tPSA of up to $\sim 20\text{\AA}$ and a reduction of cLogD by ~ 1 to 2 log units, thus increasing their polarity as expected. Although a general trend of a decrease in the lipophilicity for these metabolites relative to the parent drug

DMD #50278

was observed, the vast majority of these metabolites retained favorable lipophilicity (cLogD values between -1 and 5) for diffusion into the systemic circulation.

Most of the hydroxylated metabolites in this dataset were found circulating as free aglycones rather than their glucuronide or sulfate conjugates. This was possibly attributed to high pKa values of the hydroxyl groups, which were in the range of ~12 to 16 (Table 2A). Chemically, high pKa value could render conjugation of the hydroxyl group with the glucuronosyl moiety from uridine 5'-diphospho-glucuronic acid (UDPGA) or the sulfate group from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) unfavorable (if the alkoxy anion RO⁻ is considered the reactive species in the conjugation reaction), with a net effect of reducing CL_m of alcohol metabolites via conjugation reactions. The optimal cLogD and pKa properties impart favorable ADME characteristics to these metabolites and therefore enable their entry into the systemic circulation as free aglycones and maintain a sufficiently high exposure (>25% parent AUC). The contribution of the metabolic pathway also constituted an important parameter in determining if the metabolite circulated in the system in high concentrations, as in the case of amine metabolites. For instance, in the case of erlotinib (Figure 4), its *O*-desmethyl metabolite was found in low levels in systemic circulation despite its high pKa (13.9) and high cLogD (2.41). Assessment of the contribution of this metabolic pathway revealed that *O*-demethylation is a relatively minor route of metabolism for this drug and only accounts for 13% of the dose in humans (Ling et al., 2006). Hence, low CL_f most likely accounted for the low abundance of this metabolite (OSI-420, 5% of the parent) in human plasma (Frohna et al., 2006).

Secondary alcohols as metabolites

DMD #50278

Drugs that yielded secondary alcohols as metabolites such as flutamide, itraconazole, metoprolol (5 α -hydroxylation), pioglitazone, taranabant or buspirone also circulated as the aglycones in humans (Table 2A). Although this observation could be explained on the basis of their high pKa and cLogD values, an additional consideration to this observation could be that these metabolites are readily oxidized to their corresponding ketones, as exemplified by pioglitazone or itraconazole for which the ketone metabolite was also primarily detected in plasma (Table 2B) (Eckland and Danhof, 2000; Budde et al., 2003). Although this was not observed in all cases the possible reductase and dehydrogenase catalyzed redox cycle between the alcohol \rightarrow ketone \rightarrow alcohol (Oppermann and Maser, 2000), cannot be ruled out for these drugs. Taken together, these factors may result in reducing CL_m for the secondary alcohols, enabling these metabolites to circulate as free aglycones.

Primary alcohols as metabolites

It is well recognized that many drugs or metabolites containing a primary alcohol tend to generally undergo further oxidation to a carboxylic acid. Consequently, the alcohol metabolite may appear in low abundance, if at all, in circulation with the carboxylic acid as a major component in plasma (See Section 7 below). For example terfenadine (Figure 4) is converted to the t-butyl alcohol metabolite by CYP3A4 followed by subsequent oxidation to the corresponding carboxylic acid (fexofenadine), which is the primary metabolite found in human plasma (Abernethy et al., 2001). Other drugs that exhibit similar behavior include celecoxib, metoprolol and montelukast (Table 5). However there are exceptions where metabolites could circulate as alcohols instead of the corresponding carboxylic acids. For instance, oxidation of the tertiary butyl functionality of bupropion yields hydroxybupropion (Hesse et al., 2000). The

DMD #50278

plasma exposure of this hydroxylated metabolite, unlike those of metoprolol, montelukast, and terfenadine, is 7 to 41 times higher than that of parent. It can be postulated that the lack of subsequent oxidation to the carboxylic acid may be related to the spatial arrangement of the hydroxy group relative to the carbonyl group in the molecule. Intra-molecular cyclization of the hydroxyl metabolite leads to the formation of the corresponding cyclic *R,R*-hydroxybupropion and *S,S*-hydroxybupropion isomers (Figure 4) and possibly reduces its chance of undergoing further oxidation to the carboxylic acid. Additionally, the CL_m of the hydroxyl metabolite via the carboxylic acid pathway is probably low relative to its CL_f .

Midazolam and valdecoxib (Figure 4) represent an example of drugs in which the hydroxylated metabolite is primarily detected as glucuronide conjugate and not as an aglycone or carboxylic acid. Glucuronide conjugate of 1'-hydroxymidazolam was the major metabolite excreted in urine and circulating in plasma even though the plasma AUC of 1'-hydroxymidazolam was ~0.48 of parent after oral administration of midazolam (Heizmann and Ziegler, 1981; Eap et al., 2004). Similarly, although valdecoxib was oxidized to a hydroxymethyl metabolite (Yuan et al., 2002), the abundance of this metabolite relative to parent in plasma was low (~0.1) (Sarapa et al., 2005). Results from a human ADME study indicated that this metabolite appears to favor glucuronidation as evidenced by a recovery of 23% of dose in urine as the glucuronide conjugate of this metabolite (Yuan et al., 2002). Interestingly, in both cases, the methyl group undergoing hydroxylation is attached to 5-membered heterocyclic rings (the isoxazole in the case of valdecoxib and imidazole in the case of midazolam). It is possible that the heterocyclic rings may influence the propensity for these alcohol metabolites to undergo glucuronidation. In fact the pKa values of the hydroxymethyl metabolites suggest that the

DMD #50278

hydroxyl groups are relatively more acidic (13.09 and 13.59, respectively) compared to the primary alcohols formed via oxidation of other drugs (Table 2A). Chemically, decreased pKa could potentially increase the susceptibility of these metabolites to undergo further conjugation reaction relative to other hydroxymethyl derivatives. Additionally, potential intra-molecular interactions of the hydroxyl group with the nitrogen or oxygen atom in the imidazole or isoxazole rings of the two drugs could possibly increase the propensity of conjugation by rendering the hydroxyl group more nucleophilic. One caveat involves the limitation of our current understanding of the interaction between alcohol metabolites and the conjugation enzymes such as UGT to allow *a priori* prediction of how well the metabolites with this structural motif may undergo conjugation (i.e., assessment of CL_m). This parameter still requires further in vitro experimentation to ascertain, such as the approach used by Lutz and Isoherranen (Lutz and Isoherranen, 2012).

Bosentan represents another example of the exception to the trend that a hydroxyl metabolite on the primary carbon will undergo further oxidation to a carboxylic acid. In humans, the major metabolic pathway for bosentan involves oxidation of the t-butyl group to yield the hydroxyl metabolite Ro 48-5033. This pathway constitutes at least 35% of the overall biotransformation for bosentan (Weber et al., 1999). However, after oral administration, plasma AUC of t-butyl hydroxybosentan to parent was only ~0.13, and the corresponding carboxylic acid was not reported. Based on the pharmacokinetic framework described in Section 2, it can be postulated that the elimination clearance of this metabolite (by pathways other than further oxidation to carboxylic acid) is more likely favored relative to its formation. Bosentan is a substrate for efflux transporter such as BSEP (Hartman et al., 2010). It may be hypothesized that

DMD #50278

despite the structural modification, the t-butyl hydroxyl metabolite of bosentan may still retain the affinity to the efflux transporters as with the parent so that biliary elimination rather than further oxidation is favored. If this hypothesis holds true, it will illustrate the importance of understanding the key factors (e.g., structural motifs, physicochemical properties) that impacts the affinity of a metabolite to interact with transporters vs metabolic enzyme(s), which will determine the elimination characteristics of the metabolite. Overall, the learnings from these empiric observations can be taken into consideration when formulating a rational strategy to anticipate and prioritize which alcohol metabolite(s) should be considered for synthesis and quantitation early on to address the potential MIST or DDI issues.

5. Arenol Metabolites

Metabolites formed via hydroxylation of aromatic rings or *O*-dealkylation of aromatic ethers (arenols) represented very different characteristics when compared to their alcoholic counterparts. Thirty six drugs in this dataset revealed arenol formation as a part of their biotransformation pathway (Table 3). The physicochemical properties of the arenol metabolites were moderately changed compared to the parent drug, with increase in tPSA by ~11 to 20Å and decrease in cLogD by ~0.5 to 1 log unit associated with these metabolic steps (Table 3). Although the cLogD values of arenols, which typically ranged from -0.6 to 4, showed considerable overlap with those for the alcohol metabolites, the pKa value of the hydroxyl group in these metabolites ranged from ~4 to 11 (Table 2) and were more acidic compared to the alcohol functionality (pKa of 12-16). As seen in Table 3, in contrast to alcohol metabolites, many of the arenol metabolites circulated as glucuronide or sulfate conjugates in humans. Chemically, lower pKa of arenols would make these metabolites more ionizable to yield a

DMD #50278

phenoxy anion and hence favor the nucleophilic addition to UDPGA or PAPS. Thus, it is not surprising that these metabolites are susceptible to further conjugation and circulated either as glucuronide (e.g., apremilast, atomoxetine, chlorpromazine, codeine to morphine, desloratadine, desipramine, dextromethorphan, duloxetine, efavirenz, imipramine, loxapine, and propranolol) or sulfate (e.g., apixaban, pantoprazole, and rosiglitazone) conjugates (Table 3). While in vivo data is lacking, in vitro data suggests that 7-OH S-warfarin likely undergoes extensive glucuronide conjugation as well (Bratton et al., 2012).

Empirically, our meta-analysis suggested that conjugates of *para*-hydroxylated arensols were major circulating metabolites rather than the aglycones while the *ortho* and possibly *meta*-hydroxylated metabolites were primarily found as aglycones in circulation. For instance, the hydroxylated metabolites of atorvastatin, ranolazine, gefitinib and tramadol (Figure 5) primarily circulated as the aglycones with an M/P ratio of 0.25 or greater (Table 3). One reason for this could be limited formation of the conjugates of *ortho*-hydroxyarensols. It is also possible that the position of the hydroxy group on the aromatic ring may influence the efficiency of conjugation. Alternatively, the substrate affinity of the *ortho*-hydroxylated metabolites for various efflux transporters (e.g., MRP2 vs MRP3) may alter the disposition of these metabolites (i.e., being transported to the bile vs to the systemic circulation). The validity of these hypotheses will need to be confirmed with further research.

Some examples from this dataset illustrate the utility of the pharmacokinetic framework discussed in Section 2 in rationalizing the relative abundance of the arensol metabolite to parent. For instance, the ratio of *O*-dealkylamidarone to amidarone is low (0.1) (McDonald et al., 2012), suggesting that the CL_f is low relative to the CL_m of this metabolite. It can be postulated

DMD #50278

that since N-deethylation and 3'-hydroxy monodeethylation (Figure 5) constitutes the major metabolic pathways rather than O-dealkylation of amiodarone, the low ratio may possibly reflect a low CL_f for this metabolite. Another example may include the O-desmethyl metabolite for bosentan (Ro 47-8634) (Figure 5), which also showed low abundance relative to the parent in plasma (~0.05) (van Giersbergen et al., 2002). The relatively low fraction of dose attributable to the O-demethylation pathway in the excreta for this drug (~6.4% of dose) is consistent with this interpretation (Weber et al., 1999).

Another drug that displayed an exception to the general trend was venlafaxine. O-dealkylation of venlafaxine resulted in the formation of desvenlafaxine (Figure 5) (Howell et al., 1993). The systemic exposure of this metabolite was 4 to 6-fold higher than that of parent in CYP2D6 extensive metabolizers (EM) (Patat et al., 1998; Hynninen et al., 2008). Further, while a significant portion of the dose (26%) was excreted in urine as the glucuronide conjugate of desvenlafaxine, 29% of the dose was excreted as the aglycone (Howell et al., 1993). Also, following oral administration of desvenlafaxine, renal clearance constitutes a major clearance mechanism for the parent (DeMaio et al., 2011). This example illustrates that a change in physicochemical properties associated with the metabolite may alter its clearance mechanism, in this case, to renal elimination. This can result in a significant impact on the elimination clearance of the metabolite and thus affecting the anticipated AUC ratio of metabolite to parent. Overall this analysis suggested the complexity of predicting the exact nature of the metabolites that undergo aromatic hydroxylation or O-dealkylation and emphasized the importance to understand the formation and elimination (e.g., conjugation) characteristics of the metabolites. It

DMD #50278

also highlighted the importance of obtaining good estimates of both CL_f and CL_m in order to yield a more accurate assessment of the AUC ratio of metabolite (aglycone) to parent.

6. S- and N-Oxides Metabolites

S-Oxides

Twelve drugs in this dataset were metabolized to the corresponding sulfoxide and/or sulfone metabolites (Table 4). Formation of a sulfoxide via oxidation of the sulfur moiety lowered the cLogD by ~1 to 2 log units and increased tPSA by ~11 Å relative to the parent drug. Despite this, most of the metabolic products from sulfoxidation retained sufficient lipophilicity, as reflected by the cLogD values, in the range of -1 to 3 (Table 4). This favorable physicochemical property probably enabled these entities to reach systemic circulation since most drugs (6 out of 8) that were converted to the sulfoxide metabolite had an M/P ratio exceeding 0.25. On the other hand, relatively low amounts of the circulating sulfoxide metabolite of montelukast and ranitidine were detected in human plasma. Montelukast is primarily metabolized in humans via hydroxylation and subsequent conversion to the carboxylic acid while S-oxidation constitutes a minor pathway of metabolism for this drug. Low levels of the montelukast sulfoxide in human plasma despite the favorable cLogD and tPSA values (1.74 and 107, respectively) were therefore likely attributed to a low CL_f , while the sulfoxide metabolite of ranitidine exhibited both low f_m (0.01) and high polarity (cLogD of -3.43).

Drugs containing a sulfoxide moiety can generally undergo further oxidation to a sulfone, which can also circulate as the major metabolite. For instance, flosequinan, omeprazole, lansoprazole in CYP2C19 PM, and thioridazine were converted to their corresponding sulfone

DMD #50278

metabolites and had M/P ratios >0.25 (Table 4). This was ascribed to their favorable cLog D values since in most cases the conversion of a sulfoxide to a sulfone metabolite increased its cLogD by up to ~ 1 log unit and decreased tPSA by $\sim 6\text{\AA}$ (Table 4). The only outlier to this trend was pantoprazole which showed an M/P ratio < 0.25 . While the reason for this is not known, the discrepancy could also be ascribed to relatively less contribution of this metabolic pathway towards clearance of this drug since the major pathway appears to be *O*-dealkylation of the methoxy ether to the corresponding arenol metabolite. Of interest, in this analysis, for the parent drugs that contain sulfide in its structure, oxidative metabolism yields sulfoxide as the major circulating metabolite (e.g., albendazole, axitinib, chlorpromazine, dothiepin, thioridazine). However, the corresponding sulfone tends to be either not present as major circulating metabolite, or is present in abundance considerably lower than the corresponding sulfoxide (e.g., thioridazine). Like the amines (Section 3), this may also be attributed to the low CL_f for this sequential oxidation step.

N-Oxides

In this analysis, 11 drugs were converted to aliphatic or aromatic *N*-oxides (Table 4). Three drugs showed the formation of aromatic *N*-oxide, two of which are pyridine *N*-oxide (roflumilast and regorafenib), and one is a pyrimidine *N*-oxide (voriconazole). The remaining eight drugs formed aliphatic or cyclic *N*-oxide. Changes in physicochemical properties from *N*-oxidation are of a similar order to S-oxidation to a sulfoxide with an increase of tPSA of 14\AA and a lowering of lipophilicity of ~ 0.5 to 2 log units. All except ranitidine *N*-oxide and zolmitriptan *N*-oxide showed a cLogD value of greater than zero (Table 4), which conferred a favorable lipophilic property to enable these metabolites to reach the systemic circulation provided that a

DMD #50278

sufficient amount was formed. A notable feature of *N*-oxide metabolites is their apparent rapid interconversion with parent leading to formation of both parent drug and other primary and secondary metabolites. Interconversion after administration of the metabolite to man and animals has been shown for chlorpromazine (Jaworski et al., 1990), clozapine (Chang et al., 1998) and voriconazole (Roffey et al., 2003). The interconversion will tend to lower the M/P ratio of this type of metabolite.

7. Carboxylic Acid Metabolites

Eleven drugs in this dataset were oxidized to a carboxylic acid metabolite via various metabolic reactions (Table 5). Seven of these drugs including celecoxib, losartan, metoprolol (O-demethylation), montelukast, quetiapine, N-dealkyl product of ranolazine (CVT-2534), and terfenadine underwent oxidation of either the parent drug or the intermediate alcohol metabolite to the corresponding carboxylic acid (see Section 4). As observed from the M/P ratios, in all cases, the carboxylic acid was the major circulating metabolic product possibly due to its small volume of distribution and high plasma protein binding (Obach et al., 2008). The data from these seven drugs/metabolites was consistent with this hypothesis in that the AUC of the carboxylic acid metabolite was considerably higher compared to their corresponding alcohol metabolite. Another factor that may also influence the high M/P ratio for the carboxylic acid metabolites is related to the new functional group being introduced to the metabolite, which is metabolically more stable than the parent (i.e., reduced CL_m). This is illustrated by the example of losartan. A comparison of the pharmacokinetics of losartan to its carboxylic acid metabolite indicates that the volume of distribution (~0.45 L/kg) and plasma clearance (~9 mL/min/kg) of losartan was greater than that of carboxylosartan (plasma clearance ~0.7 mL/min/kg and volume

DMD #50278

of distribution ~ 0.14 L/Kg) (Christ, 1995; Lo et al., 1995). Additionally, the renal clearance (~ 1 mL/min/kg) of losartan accounted for 11% of its total plasma clearance while the renal clearance of carboxylosartan (~ 0.4 mL/min/kg) accounted for 55% of its total plasma clearance. The free fraction for the metabolite (~ 0.0035) showed ~ 4 -fold decrease compared to the parent (~ 0.0135), suggesting that the effect of the additional acidic function in the metabolite has little effect on unbound volume of distribution or renal clearance. However, there was a 7-fold reduction (593 mL/min/kg for losartan and 86 mL/min/kg for carboxylosartan) in unbound non-renal clearance, which can be postulated to be mainly due to the oxidation of the more labile hydroxyl group in losartan to a substantially more metabolically stable carboxyl function. Besides losartan, metabolic conversion of terfenadine to fexofenadine likely represents another example in support of this postulate.

In this dataset, 8 of 11 carboxylic acid metabolites showed negative cLogD values ranging from -0.06 (carboxylosartan) to -3.4 (zolmitriptan indoleacetic acid), suggesting that these metabolites have high polarity (Table 5). However, 7 out of 8 of these metabolites had an M/P ratio of 0.25 or higher, with repaglinide carboxylic acid (Figure 6) as the only exception. Although the mechanism(s) by which passage of these polar carboxylic acid metabolites into systemic circulation is not well understood, one possible explanation may involve transporter-mediated processes. For instance, fexofenadine is a substrate for hMRP3 (Matsushima et al., 2008), which mediates the sinusoidal efflux of a variety of organic anions. This transporter therefore may play a role in mediating the efflux of fexofenadine from the liver to the systemic circulation, as demonstrated in the Mrp3 (-/-) mouse model (Matsushima et al., 2008). Moreover, the example from the carboxylic acid metabolite of repaglinide illustrates the

DMD #50278

potential complexity of these active processes. In humans, the M/P ratio is low (0.05) despite a large fraction of the repaglinide dose being metabolized to this metabolite as evidenced by the recovery of ~60% of the dose as this metabolite in feces (presumably via biliary excretion since the formation of this metabolite involves oxidative metabolic process) and ~2% of the dose in urine (van Heiningen et al., 1999, Honkalammi et al., 2011). This suggests that the CL_m for this metabolite likely is considerably higher relative to its CL_f . Thus, it is possible that the structural motif, physicochemical and/or ADME properties for repaglinide carboxylic acid metabolite favor an interaction with the transporters involved in biliary excretion, while those for the other 7 carboxylic acid metabolites may have properties that favor an interaction with transporters that enables the passage of these metabolites into the systemic circulation. Interestingly, repaglinide and its carboxylic acid metabolite also showed one of the largest physicochemical change amongst the parent/metabolites reviewed in this analysis, with an increase in tPSA from ~80 to 125 Å and a ~4 log unit reduction in cLogD to ~-2 (Table 5). It is possible that another contributing factor to the low plasma concentration of repaglinide carboxylic acid metabolite may be attributed to low f_p , which, in this case, may become limiting for partitioning from the hepatocytes to systemic circulation. As our knowledge in transporter-mediated processes in relation to structural and physicochemical properties is relatively nascent, further research into this area is needed to elucidate these mechanisms.

8. Novel Metabolites

We have included in Table 6 novel metabolites that are formed in a “unique” manner, although in fact they are still formed mainly via oxidations and reductions. In most cases, the changes in physicochemical properties of metabolites are relatively small compared to the parent

DMD #50278

drug (Table 6). Most metabolites exhibited cLogD values within the range of -1 to 5, which conferred favorable lipophilic characteristics for their passage from the liver to the systemic circulation. For most of these metabolites, unless the structural modifications render the CL_m to be substantially different than CL_{parent} , the M/P ratios will mostly be determined by CL_f . The exception to this was methadone. Conversion of methadone to EDDP metabolite (Figure 6) showed a reduction in cLogD value by almost 6 log units from 2.29 to -3.51 and the tPSA of this metabolite is also very small ($\sim 3 \text{ \AA}$). Thus low M/P ratio of this metabolite (~ 0.16), could be ascribed, at least in part, to the high polarity and hence, low lipoidal permeability for diffusion into the systemic circulation.

9. Theoretical Considerations: Which Factors Appear Dominant?

9.1 Lipoidal Permeability

The liver is the principal site of metabolism and receives approximately 25% of the cardiac output. This equates to a blood flow of around 0.8 mL/min/g in human and 3-4 mL/min/g in rat. The capillaries supplying the hepatocytes are sinusoidal with a broken basement membrane and very leaky endothelial openings of 100-300 nm in diameter (Balaz, 2009). Drug and metabolite molecules are free to access or leave the surface of the hepatocyte without regard to physicochemistry. The membrane permeability of the hepatocyte to drugs and their metabolites has been explored in a series of studies by Chou et al. (Chou et al., 1995). The authors' analysis of the data concludes that the lower limit of PS is 0.3-0.5 mL/min/g liver for compounds with logD values below -3. Permeability then increases dramatically with increases of logD from -3 to 1.5; and at still greater logD values a plateau rate of PS of 200-300 mL/min/g

DMD #50278

liver is reached. In our data set, we have ascribed a lower cLogD value of -1 as a threshold at or above which free passive diffusion of drug and metabolites across the membrane is likely to occur and will not be a limiting factor in determining the presence of circulating metabolites. Around this value PS exceeds liver blood flow, suggesting blood perfusion rate limitations (rather than membrane permeability limitations) on influx and efflux of metabolites.

We have also considered the abundance of circulating concentrations of metabolite relative to the parent drug from the viewpoint of f_m and f_p . The term f_p was thought to be important as a low permeability rate from the hepatocyte would attenuate appearance in the circulation if further metabolism or canalicular biliary excretion occurs. It appears that whether considered from in vitro or in vivo excretion data, f_m is a much more dominant feature in determining the circulating M/P ratios relative to f_p . This is further substantiated by an additional analysis showing that there was no apparent relationship between M/P ratio and change in cLogD or tPSA values from parent drug to its corresponding metabolites (data not shown). This partly reflects the observation that most metabolites still possess favorable lipoidal permeability characteristics due to a limited change in lipophilicity and tPSA. As illustrated by the drugs evaluated in this dataset, in most cases the metabolic steps associated with circulating metabolites lower lipophilicity (cLogD) by up to ~2 log units and raise tPSA by up to ~20Å. While these metabolites typically have lower lipoidal permeability than their parent drugs, the changes are unlikely to be significant enough to render the permeability rate from the hepatocytes be a limiting factor in restricting the passage of metabolites from the liver to the systemic circulation. It is also reasonable to assume that lowered lipophilicity will result in lowered rates of metabolism and canalicular biliary excretion since the binding sites of most drug

DMD #50278

metabolizing enzymes and transporters are hydrophobic in nature. Thus, it can be conceived that the net effect of these changes associated with lower lipophilicity for the metabolites would render f_m to be a more dominant determinant relative to f_p .

9.2. Metabolite to Parent Ratio in Systemic Circulation for High Clearance Drugs

In most cases, in vivo data is lacking for CL_m so that the ratio of CL_{parent}/CL_m typically cannot be directly calculated. As discussed previously (Section 2), CL_m is an important parameter in determining the M/P ratio; thus, it represents an opportunity for further investigations to develop better predictive models for this parameter. Some of the drugs examined in this dataset are very rapidly and extensively metabolized in humans, and act virtually as pro-drugs. These include flutamide (oral clearance value of >600 mL/min/kg), buspirone (intravenous clearance value of 28 mL/min/kg and 4% bioavailability), and terfenadine (oral clearance value of >800 mL/min/kg) (Anjum et al., 1999; Mahmood and Sahajwalla, 1999; Abernethy et al., 2001). These drugs all have very high M/P ratios in systemic circulation. Although characterization of the in vivo CL_f and CL_m values for these metabolites are incomplete, the high M/P ratio after oral administration of the parent drug suggests that the fraction of the parent drug escaping the liver is low, and/or that the CL_f far exceeds the CL_m for these metabolites (see Section 2). Such drugs probably will not be a major factor in current drug discovery programs due to the emphasis on bioavailability and metabolic stability in compound design.

9.3 Protein Binding

DMD #50278

In Section 2, it was postulated that overall the reduction in lipophilicity seen in most cases of metabolism would generally result in a higher free fraction and a lower intrinsic (unbound) metabolic clearance, thus attenuating the effects on total drug clearance versus total metabolite clearance. Data are available for several drugs that undergo different metabolic routes to their corresponding metabolites to illustrate this hypothesis (Table 7). In all cases, metabolism lowered lipophilicity and increased free fraction. For those metabolites which are cleared predominantly by metabolism (5/6, desmethylclozapine being predominantly renal), their total clearances are within 3-fold of that for the parent even though their unbound metabolic or hepato-biliary (non-renal) clearance differences can be much larger. This attenuation of differences in clearance is exemplified by darifenacin and its hydroxyl metabolite. The total clearance of the parent and metabolite is similar but there is a 10-fold difference in unbound clearance and fraction unbound. Similar trends are observed for chlorpromazine, diltiazem, and tolterodine (Table 7). These examples illustrate that the effects of the reduction in lipophilicity seen with metabolism does not have pronounced effects on the ratio of $CL_{\text{parent}}/CL_{\text{m}}$.

Although the effect of reduction in lipophilicity generally tends to attenuate the influence on total drug clearance versus total metabolite clearance, an exception to this has been observed and is related to the new functional group being introduced into the metabolite, which is itself highly labile to a metabolic route unavailable to the parent. In this case, the metabolite may display a higher clearance compared to the parent. This exception is illustrated by the metabolic conversion of propranolol to hydroxypropranolol, which is cleared predominantly by conjugation of the newly introduced phenolic group. The clearance of this metabolite is higher than its parent (Table 7), consistent with its very low concentration after systemic administration and its presence being observed only after the rapid first-pass effects seen for parent after oral

DMD #50278

administration. This reinforces the observation that besides protein binding, additional factors such as the new structural motif(s) introduced into the metabolite, should be taken into consideration when determining the potential influence of structural-chemical changes on the CL_m of the metabolite.

9.4 *Pivotal role of f_m*

Given the pivotal role that f_m plays in determining the AUC of a given metabolite (see Section 2), and hence its relative abundance in relation to the parent drug, an additional analysis was conducted to ascertain if there is a threshold value that can be identified to be associated with a M/P ratio of >25%. The data (M/P ratios and f_m values) used in this analysis are listed in Tables 1-5. The f_m values were estimated from data obtained in human ADME studies with radiolabeled drugs. The fraction of dose for the metabolite of interest and its sequential metabolic products recovered in excreta (both urine and feces) was assumed to be representative of the in vivo f_m value. Data were available for a total of 60 metabolites in this dataset, and the results are presented in Figure 7. Visual inspection of this data suggests that f_m value of 0.15 appears to be a reasonable threshold value to distinguish the metabolites that may circulate with abundance exceeding the threshold of 25% of parent AUC compared to those that do not. Using this threshold f_m value, 4 metabolites have been identified as “false negative” in that while the f_m values are below 0.15, the M/P ratios exceed the threshold of 0.25. These 4 metabolites are cyclized indole product of elzasonan, 2-hydroxycyclopentyl ruxolitinib, zolmitriptan N-oxide, and N-desmethylzolmitriptan. The reason(s) for this is not readily apparent, but it could be postulated that underestimation of f_m (possibly related to incomplete collection of radioactive dose and/or characterization of the metabolite profiles for elzasonan (79% of dose recovered),

DMD #50278

and potential back-conversion of N-oxide to parent in feces for zolmitriptan) may have contributed to this discordance.

When the M/P ratio for the aglycone of the arenol metabolites were used in this analysis, there were 14 “false positives” identified (i.e., f_m of >0.15 but M/P of <0.25). Amongst these 14 “false positives”, the most common structures are the arenol metabolites where the circulating level of the aglycone is negligible. However, when the M/P values for the conjugates of these arenol metabolites were plotted instead (Figure 7), the number of “false positive” is reduced to 6. Three of these 6 metabolites appear to be conjugated (5-hydroxyelzasonan, O-desalkyl flecainide, and 5-methylhydroxy valdecocixib) and excreted; however, the conjugate was not reported to be present in plasma. Two additional metabolites (t-butyl hydroxybosentan and repaglinide carboxylic acid) may also display similar properties in that once formed, these metabolites likely favors excretion into the bile rather than being transported to the systemic circulation. N-desmethylzopiclone represents a marginal case in that the f_m value is estimated to be 0.17 while the M/P ratio is 0.2.

Using the threshold f_m value of 0.15, the false negative and the false positive rates (when the M/P ratios of the conjugates of the arenol metabolites are considered) are 10% (4 out of 40 metabolites) and 30% (6 out of 20 metabolites), respectively, which represent a reasonable trade-off since further reduction in the false negative rate will increase the false positive rate considerably. Moreover, this suggested threshold value may be further refined as more data become available in the future.

10. Special Considerations

DMD #50278

10.1 Genetic Polymorphisms

Genetic polymorphisms of the drug metabolizing enzymes play an important role in influencing the disposition of many drugs and metabolites. In this analysis, a few examples illustrate the important influence of genetic polymorphisms on the abundance of major circulating metabolites. The first set of examples involves the differences in the abundance of circulating metabolites between extensive (EM) and poor metabolizers (PM) of CYP2C19 and CYP2D6. In the former case, the M/P ratio of lansoprazole sulfone in CYP2C19 PM (0.88) was considerably higher than those in the CYP2C19 homozygous EM and heterozygous EM (<0.1). Another example that showed a similar pattern was atomoxetine, which is metabolized by CYP2D6 (aromatic 4-hydroxylation, major pathway in CYP2D6 EM) and CYP2C19 (N-demethylation) (Sauer et al., 2005). In CYP2D6 EM subjects, 4-OH atomoxetine glucuronide constitutes the major circulating moiety, while the plasma AUC ratios of N-desmethyl atomoxetine and 4-hydroxyatomoxetine to parent were low (<0.1) (Sauer et al., 2003). Moreover, in CYP2D6 PM, the AUC ratio of N-desmethyl atomoxetine to parent drug was 0.33 to 0.42 (Sauer et al., 2003); FDA Clinical Pharmacology Review: http://www.accessdata.fda.gov/drugsatfda_docs/nda/2002/21-411_Strattera_biopharmr_P2.pdf), thus constituting a moiety that needs to be considered for assessment for potential DDI under the current EMA and draft FDA Guidance. These examples highlight the need to evaluate the abundance of circulating metabolites in relation to the genotype and phenotype of the metabolizing enzymes, if genetic polymorphisms are known to exist for the particular enzyme of interest.

DMD #50278

Another set of examples that illustrate the importance of gene dose on the abundance of the circulating metabolites include nortriptyline and codeine. In humans, nortriptyline undergoes CYP2D6-mediated hydroxylation to form the cyclic E-10-hydroxynortriptyline (major form of this metabolite), while the formation of Z-10-hydroxynortriptyline (minor form of this metabolite) is not related to CYP2D6 polymorphism. Dalen and co-workers demonstrated that the ratio of AUC of 10-hydroxynortriptyline to parent increased with the number of functional CYP2D6 genes, ranging from 0.36 in subjects with no functional CYP2D6 genes to 13 in subjects with 13 functional CYP2D6 genes (Dalen et al., 1998). Likewise, in humans, codeine undergoes CYP2D6-mediated demethylation to form morphine, which is glucuronidated to morphine 3-glucuronide and morphine 6-glucuronide. In subjects who are CYP2D6 PM, ratio of plasma AUC for morphine to codeine was minimal (0.008), whereas this ratio was higher in subjects who are CYP2D6 EM (0.06) and CYP2D6 UM (0.08) (Kirchheiner et al., 2007). Furthermore, in these same subjects, the systemic exposure (plasma AUC) of morphine 3-glucuronide and morphine 6-glucuronide (pharmacologically active) in CYP2D6 EM and UM were substantially higher than in CYP2D6 PM. Thus, gene dose can exert a significant influence on the systemic exposure to the circulating metabolites and should be taken into consideration when attempting to project the abundance of circulating metabolites.

10.2. Caveats to this Analysis

This analysis is intended to assess general trends regarding possible relationship of structural motif and the associated physicochemical properties of the metabolites to their relative abundance in systemic circulation. Attempts to rationalize some of the observed trends were made based on existing known biologic and/or chemistry principles. However, there are some

DMD #50278

significant knowledge gaps in certain areas where mechanistic insights are lacking to fully understand the observed results. Plausible hypotheses for further testing are postulated for subsequent confirmation when feasible. Not surprisingly, results of this analysis point to the need for: (1) further understanding of the scaling for conjugation reactions (e.g., glucuronidation and sulfation) to provide a more accurate estimate of CL_m for the alcohol and arenol metabolites in order to better predict the ratio of AUC of metabolite to parent, (2) an integrated approach to combine metabolism and transport data for quantitative prediction, particularly the transport processes in the liver that mediate biliary excretion vs passage into systemic circulation, and (3) a more complete understanding of the interplay between physicochemical properties for passive processes with spatial arrangement of the compound for its affinity to enzymatic/active transport processes.

Another caveat to our results involves the complexity of the metabolic pathways. In this dataset, the vast majority of the circulating metabolites are formed from primary biotransformation pathways. However, in humans, there are examples where the major circulating metabolites are derived from secondary or tertiary biotransformation steps, e.g., torceptrapib (Dalvie et al., 2008). Since the ability for the *in vitro* systems to predict these secondary or tertiary metabolites is poor (Dalvie et al., 2009), there is still the necessity to confirm the major circulating metabolites in a human *in vivo* ADME study with radiolabeled drug. In addition, this analysis only addressed the relative abundance of the metabolite to parent, and there is no attempt to assess structure-physicochemical properties in relation to pharmacologic activity. Clearly, in drug discovery and development programs, an understanding of both aspects is essential.

DMD #50278

11. Summary

To our knowledge, this is one of the first attempts to assess the relative abundance of circulating metabolites using an integrated approach of principles of metabolite kinetics along with structure-physicochemical properties of the parent/metabolites. The analysis was conducted in a dataset comprising 125 drugs with their circulating metabolites. The major factors identified in this analysis that will favor a given circulating metabolite to exceed the threshold of 25% of parent appears be: a f_m value of >15%, a $\log D$ value of > -1, and structural motif that minimizes the CL_m relative to its CL_f . Potential hypotheses are proposed where exceptions to these general trends are observed, which provides opportunities for further scientific investigations to refine these recommendations.

DMD #50278

Authorship Contribution:

Participated in research design: Cho-Ming Loi, Dennis Smith, Deepak Dalvie

Performed data analysis: Cho-Ming Loi, Dennis Smith, Deepak Dalvie

Wrote or contributed to the writing of the manuscript: Cho-Ming Loi, Dennis Smith, Deepak

Dalvie

DMD #50278

References

- Abernethy DR, Barbey JT, Franc J, Brown KS, Feirrer I, Ford N, and Salazar DE (2001) Loratadine and terfenadine interaction with nefazodone: Both antihistamines are associated with QTc prolongation. *Clin Pharmacol Ther* **69**:96-103.
- Anjum S, Swan SK, Lambrecht LJ, Radwanski E, Cutler DL, Affrime MB, and Halstenson CE (1999) Pharmacokinetics of flutamide in patients with renal insufficiency. *Br J Clin Pharmacol* **47**:43-47.
- Baillie TA (2008) Metabolism and toxicity of drugs. Two decades of progress in industrial drug metabolism. *Chem Res Toxicol* **21**:129-137.
- Balaz S (2009) Modeling kinetics of subcellular disposition of chemicals. *Chem Rev* **109**:1793-1899.
- Borin MT, Chambers JH, Carel BJ, Gagnon S, and Freimuth WW (1997a) Pharmacokinetic study of the interaction between rifampin and delavirdine mesylate. *Clin Pharmacol Ther* **61**:544-553.
- Borin MT, Cox SR, Herman BD, Carel BJ, Anderson RD, and Freimuth WW (1997b) Effect of fluconazole on the steady-state pharmacokinetics of delavirdine in human immunodeficiency virus-positive patients. *Antimicrob Agents Chemother* **41**:1892-1897.
- Boyd RA, Chin SK, Don-Pedro O, Verotta D, Sheiner LB, Williams RL, and Giacomini KM (1989) The pharmacokinetics and pharmacodynamics of diltiazem and its metabolites in healthy adults after a single oral dose. *Clin Pharmacol Ther* **46**:408-419.
- Bratton SM, Mosher CM, Khallouki F, Finel M, Court MH, Moran JH, and Radominska-Pandya A (2012) Analysis of R- and S-hydroxywarfarin glucuronidation catalyzed by human liver microsomes and recombinant UDP-glucuronosyltransferases. *J Pharmacol Exp Ther* **340**:46-55.
- Budde K, Neumayer HH, Fritsche L, Sulowicz W, Stompor T, and Eckland D (2003) The pharmacokinetics of pioglitazone in patients with impaired renal function. *Br J Clin Pharmacol* **55**:368-374.
- Chang WH, Lin SK, Lane HY, Wei FC, Hu WH, Lam YW, and Jann MW (1998) Reversible metabolism of clozapine and clozapine N-oxide in schizophrenic patients. *Prog Neuropsychopharmacol Biol Psychiatry* **22**:723-739.
- Chou C, McLachlan AJ, and Rowland M (1995) Membrane permeability and lipophilicity in the isolated perfused rat liver: 5-ethyl barbituric acid and other compounds. *J Pharmacol Exp Ther* **275**:933-940.

DMD #50278

- Christ DD (1995) Human plasma protein binding of the angiotensin II receptor antagonist losartan potassium (DuP 753/MK 954) and its pharmacologically active metabolite EXP3174. *J Clin Pharmacol* **35**:515-520.
- Dalen P, Dahl ML, Bernal Ruiz ML, Nordin J, and Bertilsson L (1998) 10-Hydroxylation of nortriptyline in white persons with 0, 1, 2, 3, and 13 functional CYP2D6 genes. *Clin Pharmacol Ther* **63**:444-452.
- Dalvie D, Chen W, Zhang C, Vaz AD, Smolarek TA, Cox LM, Lin J, and Obach RS (2008) Pharmacokinetics, metabolism, and excretion of torcetrapib, a cholesteryl ester transfer protein inhibitor, in humans. *Drug Metab Dispos* **36**:2185-2198.
- Dalvie D, Obach RS, Kang P, Prakash C, Loi CM, Hurst S, Nedderman A, Goulet L, Smith E, Bu HZ, and Smith DA (2009) Assessment of three human in vitro systems in the generation of major human excretory and circulating metabolites. *Chem Res Toxicol* **22**:357-368.
- Davis-Bruno KL and Atrakchi A (2006) A regulatory perspective on issues and approaches in characterizing human metabolites. *Chem Res Toxicol* **19**:1561-1563.
- DeMaio W, Kane CP, Nichols AI, and Jordan R (2011) Metabolism Studies of Desvenlafaxine. *J Bioequiv Availab* **3**:151-160.
- Eap CB, Buclin T, Cucchia G, Zullino D, Hustert E, Bleiber G, Golay KP, Aubert AC, Baumann P, Telenti A, and Kerb R (2004) Oral administration of a low dose of midazolam (75 microg) as an in vivo probe for CYP3A activity. *Eur J Clin Pharmacol* **60**:237-246.
- Eckland DA and Danhof M (2000) Clinical Pharmacokinetics of pioglitazone. *Exp Clin Endocrinol Diabetes* **108**:S234-S-242.
- Freedberg KA, Innis RB, Creese I, and Snyder SH (1979) Antischizophrenic drugs: differential plasma protein binding and therapeutic activity. *Life Sci* **24**:2467-2473.
- Frohna P, Lu J, Eppler S, Hamilton M, Wolf J, Rakhit A, Ling J, Kenkare-Mitra SR, and Lum BL (2006) Evaluation of the absolute oral bioavailability and bioequivalence of erlotinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in a randomized, crossover study in healthy subjects. *J Clin Pharmacol* **46**:282-290.
- Fura A (2006) Role of pharmacologically active metabolites in drug discovery and development. *Drug Discov Today* **11**:133-142.
- Gschwind HP, Pfaar U, Waldmeier F, Zollinger M, Sayer C, Zbinden P, Hayes M, Pokorny R, Seiberling M, Ben-Am M, Peng B, and Gross G (2005) Metabolism and disposition of imatinib mesylate in healthy volunteers. *Drug Metab Dispos* **33**:1503-1512.

DMD #50278

Hartman JC, Brouwer K, Mandagere A, Melvin L, and Gorczynski R (2010) Evaluation of the endothelin receptor antagonists ambrisentan, darusentan, bosentan, and sitaxsentan as substrates and inhibitors of hepatobiliary transporters in sandwich-cultured human hepatocytes. *Can J Physiol Pharmacol* **88**:682-691.

Heizmann P and Ziegler WH (1981) Excretion and metabolism of 14C-midazolam in humans following oral dosing. *Arzneimittelforschung* **31**:2220-2223.

Herrlin K, Yasui-Furukori N, Tybring G, Widen J, Gustafsson LL, and Bertilsson L (2003) Metabolism of citalopram enantiomers in CYP2C19/CYP2D6 phenotyped panels of healthy Swedes. *Br J Clin Pharmacol* **56**:415-421.

Hesse LM, Venkatakrishnan K, Court MH, von Moltke LL, Duan SX, Shader RI, and Greenblatt DJ (2000) CYP2B6 mediates the in vitro hydroxylation of bupropion: potential drug interactions with other antidepressants. *Drug Metab Dispos* **28**:1176-1183.

Honkalammi J, Niemi M, Neuvonen PJ, and Backman JT (2011) Dose-dependent interaction between gemfibrozil and repaglinide in humans: strong inhibition of CYP2C8 with subtherapeutic gemfibrozil doses. *Drug Metab Dispos* **39**:1977-1986.

Houston JB (1982) Drug Metabolite Kinetics. *Pharmacol Ther* **15**:521-552.

Howell SR, Husbands GE, Scatina JA, and Sisenwine SF (1993) Metabolic disposition of 14C-venlafaxine in mouse, rat, dog, rhesus monkey and man. *Xenobiotica* **23**:349-359.

Hynninen VV, Olkkola KT, Bertilsson L, Kurkinen K, Neuvonen PJ, and Laine K (2008) Effect of terbinafine and voriconazole on the pharmacokinetics of the antidepressant venlafaxine. *Clin Pharmacol Ther* **83**:342-348.

Isin EM, Elmore CS, Nilsson GN, Thompson RA, and Weidolf L (2012) Use of radiolabeled compounds in drug metabolism and pharmacokinetic studies. *Chem Res Toxicol* **25**:532-542.

Isoherranen N, Hachad H, Yeung CK, and Levy RH (2009) Qualitative analysis of the role of metabolites in inhibitory drug-drug interactions: literature evaluation based on the metabolism and transport drug interaction database. *Chem Res Toxicol* **22**:294-298.

Isoherranen N, Kunze KL, Allen KE, Nelson WL, and Thummel KE (2004) Role of itraconazole metabolites in CYP3A4 inhibition. *Drug Metab Dispos* **32**:1121-1131.

Jaworski TJ, Hawes EM, McKay G, and Midha KK (1990) The metabolism of chlorpromazine N-oxide in man and dog. *Xenobiotica* **20**:107-115.

Kamel A, Obach RS, Colizza K, Wang W, O'Connell TN, Coelho RV, Jr., Kelley RM, and Schildknecht K (2010) Metabolism, pharmacokinetics, and excretion of the 5-

DMD #50278

- hydroxytryptamine1b receptor antagonist elzasonan in humans. *Drug Metab Dispos* **38**:1984-1999.
- Kassahun K, Mattiuz E, Nyhart E, Jr., Obermeyer B, Gillespie T, Murphy A, Goodwin RM, Tupper D, Callaghan JT, and Lemberger L (1997) Disposition and biotransformation of the antipsychotic agent olanzapine in humans. *Drug Metab Dispos* **25**:81-93.
- Kerbusch T, Milligan PA, and Karlsson MO (2004) Assessment of the relative in vivo potency of the hydroxylated metabolite of darifenacin in its ability to decrease salivary flow using pooled population pharmacokinetic-pharmacodynamic data. *Br J Clin Pharmacol* **57**:170-180.
- Kim KA, Song WK, and Park JY (2009) Association of CYP2B6, CYP3A5, and CYP2C19 genetic polymorphisms with sibutramine pharmacokinetics in healthy Korean subjects. *Clin Pharmacol Ther* **86**:511-518.
- Kirchheiner J, Schmidt H, Tzvetkov M, Keulen JT, Lotsch J, Roots I, and Brockmoller J (2007) Pharmacokinetics of codeine and its metabolite morphine in ultra-rapid metabolizers due to CYP2D6 duplication. *Pharmacogenomics J* **7**:257-265.
- Kroemer HK, Funck-Brentano C, Silberstein DJ, Wood AJ, Eichelbaum M, Woosley RL, and Roden DM (1989) Stereoselective disposition and pharmacologic activity of propafenone enantiomers. *Circulation* **79**:1068-1076.
- Levy RH, Lane EA, Guyot M, Brachet-Liermain A, Cenraud B, and Loiseau P (1983) Analysis of parent drug-metabolite relationship in the presence of an inducer. Application to the carbamazepine-clobazam interaction in normal man. *Drug Metab Dispos* **11**:286-292.
- Lin JH and Lu AY (1997) Role of pharmacokinetics and metabolism in drug discovery and development. *Pharmacol Rev* **49**:403-449.
- Ling J, Johnson KA, Miao Z, Rakhit A, Pantze MP, Hamilton M, Lum BL, and Prakash C (2006) Metabolism and excretion of erlotinib, a small molecule inhibitor of epidermal growth factor receptor tyrosine kinase, in healthy male volunteers. *Drug Metab Dispos* **34**:420-426.
- Lo MW, Goldberg MR, McCrea JB, Lu H, Furtek CI, and Bjornsson TD (1995) Pharmacokinetics of losartan, an angiotensin II receptor antagonist, and its active metabolite EXP3174 in humans. *Clin Pharmacol Ther* **58**:641-649.
- Lutz JD, Fujioka Y, and Isoherranen N (2010) Rationalization and prediction of in vivo metabolite exposures: the role of metabolite kinetics, clearance predictions and in vitro parameters. *Expert Opin Drug Metab Toxicol* **6**:1095-1109.

DMD #50278

- Lutz JD and Isoherranen N (2012) Prediction of relative in vivo metabolite exposure from in vitro data using two model drugs: dextromethorphan and omeprazole. *Drug Metab Dispos* **40**:159-168.
- Mahmood I and Sahajwalla C (1999) Clinical pharmacokinetics and pharmacodynamics of buspirone, an anxiolytic drug. *Clin Pharmacokinet* **36**:277-287.
- Matsushima S, Maeda K, Hayashi H, Debori Y, Schinkel AH, Schuetz JD, Kusuhara H, and Sugiyama Y (2008) Involvement of multiple efflux transporters in hepatic disposition of fexofenadine. *Mol Pharmacol* **73**:1474-1483.
- McDonald MG, Au NT, Wittkowsky AK, and Rettie AE (2012) Warfarin-amiodarone drug-drug interactions: determination of $[I](u)/K(I,u)$ for amiodarone and its plasma metabolites. *Clin Pharmacol Ther* **91**:709-717.
- Morse GD, Fischl MA, Shelton MJ, Cox SR, Driver M, DeRemer M, and Freimuth WW (1997) Single-dose pharmacokinetics of delavirdine mesylate and didanosine in patients with human immunodeficiency virus infection. *Antimicrob Agents Chemother* **41**:169-174.
- Nedderman AN, Dear GJ, North S, Obach RS, and Higton D (2011) From definition to implementation: a cross-industry perspective of past, current and future MIST strategies. *Xenobiotica* **41**:605-622.
- Obach RS, Lombardo F, and Waters NJ (2008) Trend analysis of a database of intravenous pharmacokinetic parameters in humans for 670 drug compounds. *Drug Metab Dispos* **36**:1385-1405.
- Oppermann UC and Maser E (2000) Molecular and structural aspects of xenobiotic carbonyl metabolizing enzymes. Role of reductases and dehydrogenases in xenobiotic phase I reactions. *Toxicology* **144**:71-81.
- Pahlman I and Gozzi P (1999) Serum protein binding of tolterodine and its major metabolites in humans and several animal species. *Biopharm Drug Dispos* **20**:91-99.
- Patat A, Troy S, Burke J, Trocherie S, Danjou P, Le Coz F, Allain H, and Gandon JM (1998) Absolute bioavailability and electroencephalographic effects of conventional and extended-release formulations of venlafaxine in healthy subjects. *J Clin Pharmacol* **38**:256-267.
- Penner N, Xu L, and Prakash C (2012) Radiolabeled absorption, distribution, metabolism, and excretion studies in drug development: why, when, and how? *Chem Res Toxicol* **25**:513-531.

DMD #50278

- Riley P, Figary PC, Entwisle JR, Roe AL, Thompson GA, Ohashi R, Ohashi N, and Moorehead TJ (2005) The metabolic profile of azimilide in man: in vivo and in vitro evaluations. *J Pharm Sci* **94**:2084-2095.
- Rochester CL, Gammon DE, Shane E, and Bilezikian JP (1980) A radioreceptor assay for propranolol and 4-hydroxypropranolol. *Clin Pharmacol Ther* **28**:32-39.
- Roffey SJ, Cole S, Comby P, Gibson D, Jezequel SG, Nedderman AN, Smith DA, Walker DK, and Wood N (2003) The disposition of voriconazole in mouse, rat, rabbit, guinea pig, dog, and human. *Drug Metab Dispos* **31**:731-741.
- Roffey SJ, Obach RS, Gedge JI, and Smith DA (2007) What is the objective of the mass balance study? A retrospective analysis of data in animal and human excretion studies employing radiolabeled drugs. *Drug Metab Rev* **39**:17-43.
- Sarapa N, Britto MR, Mainka MB, and Parivar K (2005) The effect of mild and moderate hepatic impairment on the pharmacokinetics of valdecoxib, a selective COX-2 inhibitor. *Eur J Clin Pharmacol* **61**:247-256.
- Sauer JM, Ponsler GD, Mattiuz EL, Long AJ, Witcher JW, Thomasson HR, and Desante KA (2003) Disposition and metabolic fate of atomoxetine hydrochloride: the role of CYP2D6 in human disposition and metabolism. *Drug Metab Dispos* **31**:98-107.
- Sauer JM, Ring BJ, and Witcher JW (2005) Clinical pharmacokinetics of atomoxetine. *Clin Pharmacokinet* **44**:571-590.
- Schaber G, Stevens I, Gaertner HJ, Dietz K, and Breyer-Pfaff U (1998) Pharmacokinetics of clozapine and its metabolites in psychiatric patients: plasma protein binding and renal clearance. *Br J Clin Pharmacol* **46**:453-459.
- Smith DA and Dalvie D (2012) Why do metabolites circulate? *Xenobiotica* **42**:107-126.
- van Giersbergen PL, Halabi A, and Dingemans J (2002) Single- and multiple-dose pharmacokinetics of bosentan and its interaction with ketoconazole. *Br J Clin Pharmacol* **53**:589-595.
- van Heiningen PN, Hatorp V, Kramer Nielsen K, Hansen KT, van Lier JJ, De Merbel NC, Oosterhuis B, and Jonkman JH (1999) Absorption, metabolism and excretion of a single oral dose of (14)C-repaglinide during repaglinide multiple dosing. *Eur J Clin Pharmacol* **55**:521-525.
- von Moltke LL, Greenblatt DJ, Granda BW, Grassi JM, Schmider J, Harmatz JS, and Shader RI (1999) Nefazodone, meta-chlorophenylpiperazine, and their metabolites in vitro: cytochromes mediating transformation, and P450-3A4 inhibitory actions. *Psychopharmacology (Berl)* **145**:113-122.

DMD #50278

- Voorman RL, Maio SM, Hauer MJ, Sanders PE, Payne NA, and Ackland MJ (1998a) Metabolism of delavirdine, a human immunodeficiency virus type-1 reverse transcriptase inhibitor, by microsomal cytochrome P450 in humans, rats, and other species: probable involvement of CYP2D6 and CYP3A. *Drug Metab Dispos* **26**:631-639.
- Voorman RL, Maio SM, Payne NA, Zhao Z, Koeplinger KA, and Wang X (1998b) Microsomal metabolism of delavirdine: evidence for mechanism-based inactivation of human cytochrome P450 3A. *J Pharmacol Exp Ther* **287**:381-388.
- Weber C, Gasser R, and Hopfgartner G (1999) Absorption, excretion, and metabolism of the endothelin receptor antagonist bosentan in healthy male subjects. *Drug Metab Dispos* **27**:810-815.
- Weiss YA, Safar ME, Lehner JP, Levenson JA, Simon A, and Alexandre JM (1978) (+)-Propranolol clearance, an estimation of hepatic blood flow in man. *Br J Clin Pharmacol* **5**:457-460.
- Yeung CK, Fujioka Y, Hachad H, Levy RH, and Isoherranen N (2011) Are circulating metabolites important in drug-drug interactions?: Quantitative analysis of risk prediction and inhibitory potency. *Clin Pharmacol Ther* **89**:105-113.
- Yeung PK, Hubbard JW, Korchinski ED, and Midha KK (1993) Pharmacokinetics of chlorpromazine and key metabolites. *Eur J Clin Pharmacol* **45**:563-569.
- Yuan JJ, Yang DC, Zhang JY, Bible R, Jr., Karim A, and Findlay JW (2002) Disposition of a specific cyclooxygenase-2 inhibitor, valdecoxib, in human. *Drug Metab Dispos* **30**:1013-1021.

DMD #50278

Legends for Figures

- Figure 1 Scheme depicting potential routes of metabolite disposition
- Figure 2 Relationship between M/P ratio and cLog D values for amine metabolites (A, all metabolites and B, expanded view).
- Figure 3 Structures of selected drugs that form amine metabolites.
- Figure 4 Structures of selected drugs that form alcohol metabolites.
- Figure 5 Structures of selected drugs that form arenol metabolites.
- Figure 6 Scheme showing conversion of repaglinide to repaglinide carboxylic acid and methadone to EDDP metabolite.
- Figure 7 Relationship of metabolite to parent AUC ratio as a function of f_m (A, all metabolites and B, expanded view).

Table 1: Physicochemical properties, abundance and estimated fm of amine metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	cLogD	tPSA	M/P	fm	REF
Amiodarone	5.87	42.7	N-Desethylamiodarone	5.2	51.5	1*		(McDonald et al., 2012)
			Didesethylamiodarone	5.25	65.5	0.05*		
Amitriptyline	2.64	3.24	Nortriptyline	1.49	12	(CYP2C19) EM: 0.43-0.99; PM: 0.13		(Burch and Hullin, 1981; Venkatakrishnan et al., 2001; Jiang et al., 2002)
Atomoxetine	0.77	21.3	Desmethylatomoxetine	1.39	35.3	(CYP2D6) EM:0.06; PM:0.38	0.03	(Sauer et al., 2003; Sauer et al., 2005); FDA Clinical Pharmacology Review ^a
Azelastine	1.57	35.9	Desmethylazelastine	1.63	44.7	0.41		Product Label ^b
Azimilide	2.66	72.6	N-desmethylazilimide	1.13	81.39	NR	0.04	(Riley et al., 2005)
Buspirone	1.24	69.6	1-Pyrimidinylpiperazine	-0.46	41.1	44 to 57		(Jajoo et al., 1989; Dockens et al., 2006)
Citalopram	1.35	36.3	Desmethylcitalopram	0.014	45.1	(CYP2C19) PM:0.24; EM:0.49		(Herrlin et al., 2003; Yu et al., 2003)
			Didesmethylcitalopram	0.003	45.1	(CYP2C19) EM:0.07; PM:0.05		
Clobazam	1.22	40.6	Norclobazam	1.42	49.4	2.3		(Levy et al., 1983)

DMD #50278

Clomipramine	2.95	6.48	Desmethylclomipramine	1.87	15.3	0.37		(Nagy and Johansson, 1977)
Clozapine	3.5	30.9	Desmethylclozapine	3.27	39.7	0.32	0.17	(Chung et al., 1993)
Delavirdine	1.77	119	Desisopropyldeavirdine	1.59	133	SD:1.1; MD:0.14		(Borin et al., 1997a; Borin et al., 1997b; Morse et al., 1997);
Diazepam	2.8	32.7	Desmethyldiazepam	2.78	41.5	0.35 - 1.1		(Perucca et al., 1994; Kosuge et al., 2001; Saari et al., 2007)
Diltiazem	3.36	84.4	N-Desmethyldiltiazem	2.07	93.2	0.35	0.49	(Sugihara et al., 1984; Yeung et al., 1990)
Dothiepin	2.54	28.5	N-Desmethyldothiepin	1.39	37.3	0.54		(Maguire et al., 1981)
Doxepin	2.06	12.5	Desmethyldoxepin	0.91	21.3	0.6 - 1.4		(Kirchheiner et al., 2005)
Enzalutamide	3.17	99.74	Desmethylenzalutamide	2.98	109	1.60		Product Label ^c
Elzasonan	3.24	52.1	Desmethyl elzasonan	2.19	60.9	0.19	NR	(Kamel et al., 2010)
Fenfluramine	0.91	12	Desethylfenfluramine	0.33	26	0.36 - 0.37		(Caccia et al., 1985)
Fentanyl	2.61	23.5	Norfentanyl	-0.44	32.3	2.9		(Yeung et al., 2011)
Fluoxetine	1.41	21.3	Norfluoxetine	2.12	35.3	0.4 - 9.4		(Hamelin et al., 1996; Ouellet et al., 1998; Fjordside et al., 1999; Moraes et al., 1999; Gupta et al., 2004)
Halofantrine	6.93	23.5	Desbutylhalofantrine	3.79	32.3	1.3		(Yeung et al., 2011)
Imatinib	2.49	86.3	Desmethylinatinib	1	95.1	0.16	0.15	(Gschwind et al., 2005)

Imipramine	2.34	6.48	Desipramine	1.27	15.3	1.1		(Suffin et al., 1984; Wells et al., 1986; Suffin et al., 1988; Spina et al., 1997)
Itraconazole	4.93	101	Dealkylitraconazole	3.51	110	0.05		(Templeton et al., 2008)
Ivabradine	3.32	60.1	Desmethylyvabradine	2.37	69.3	0.38 - 0.49		(Portoles et al., 2006a; Portoles et al., 2006b)
Loratadine	3.9	42.4	Desloratidine	0.84	24.3	2.3	0.55	(Brannan et al., 1995) (Ramanathan et al., 2007a)
Loxapine	2.23	28.1	Desmethyloxapine	1.79	36.7	0.24		(Simpson et al., 1978; Cheung et al., 1991)
Lumefantrine	7.42	23.5	Desbutylumefantrine	4.76	32.3	0.05		(McGready et al., 2006)
Mianserin	0.76	6.48	Desmethylmianserin	0.49	15.3	0.61		(Pinder and Van Delft, 1983b; Pinder and van Delft, 1983a)
Mirtazapine	0.28	19.4	Desmethylmirtazapine	-0.35	28.2	0.47 - 0.75	0.25	(Timmer et al., 2000)
Nefazodone	3.89	51.6	m-Chlorophenylpiperazine	0.56	15.3	0.09 - 0.19		(Barbhaiya et al., 1995; Barbhaiya et al., 1996a; Barbhaiya et al., 1996b)
Olanzapine	2.35	59.1	Desmethyloanzapine	2.21	67.9	NR	0.04	(Kassahun et al., 1997)
Otenabant (CP-945598)	4.71	102	N-desethylotenabant	3.63	116	4.6		(Miao et al., 2012)
Oxybutynin	4.15	49.8	Desethyloxybutynin	4.39	58.6	R:21.5; S:8.3		(Zobrist et al., 2001)
Propafenone	1.37	58.6	Despropylprofenanone	0.68	72.6	0.06 - 0.18		(Kates et al., 1985; Dilger et al., 1999)

Ranitidine	-1.07	112	Desmethylranitidine	-2.22	120	0.06	0.01	(Prueksaritanont et al., 1989); Product Label ^d
Ranolazine	-1.28	74.3	Desalkylranolazine	-2.16	53.9	0.33 - 0.37		(Abdallah and Jerling, 2005; Jerling and Abdallah, 2005; Jerling et al., 2005)
Regalinide	1.99	78.87	N-dealkylregaplinide (M1)	-0.58	101.65	0.06	0.04	(Honkalammi et al., 2011)
Rosiglitazone	1.93	96.8	Desmethylrosiglitazone	1.39	106	0.58 - 1.2	0.21	(Cox et al., 2000; Kirchheiner et al., 2006)
Sertraline	3.04	12	Desmethylsertraline	3.17	26	1.2 - 1.7		(Hamelin et al., 1996)
Sibutramine	3.24	3.24	Desmethylsibutramine	2.21	12	3.8		(Kim et al., 2009; Bae et al., 2011)
			Didesmethylsibutramine	1.89	26	12		
Sildenafil	2.45	118	Desmethylsildenafil	0.97	126	0.5	0.22	(Muirhead et al., 2002)
			Cleaved piperazine ring	-1.24	135	0.27	0.22	
Sunitinib	0.44	77.2	Desethylsunitinib	-0.91	86	0.4	0.32	(Speed et al., 2012)
Tamoxifen	3.83	12.5	Desmethyltamoxifen	3.19	21.3	1.8		(Yeung et al., 2011)
			Didesmethyltamoxifen	3.11	35.3	0.37		
Terbinafine	5.46	3.24	Desmethylterbinafine	5.1	12	0.9		(Robbins et al., 1996)
Tramadol	0.29	32.7	Desmethyltramadol	-1.11	41.5	0.27		(Ardakani et al., 2008; de Moraes et al., 2012)

DMD #50278

Udenafil	-1.21	126	Desalkyludenafil	0.77	137	1.2 - 1.7		(Shin et al., 2010)
Vardenafil	3.45	118	Desethylvardenafil	1.64	126	0.73		(Ku et al., 2009)
Verapamil	2.46	63.9	Norverapamil	1.35	72.7	1.0 - 1.3		(Abernethy et al., 2000)
			D617	-0.69	54.3	0.85 - 1.2		
			D620	-1.29	30.5	0.32 - 0.46		
Zimelidine	2.44	16.1	Desmethylzimelidine	1.38	24.9	3		(Brown et al., 1980)
Zolmitriptan	-1.47	57.4	Desmethylzolmitriptan	-2.97	66.2	0.6	0.04	(Dixon et al., 1997; Seaber et al., 1997)
Zopiclone	2.63	91.8	Desmethylzopiclone	1.57	101	0.2	0.17	(Fernandez et al., 1995; Mistri et al., 2008)

1 Structures of drugs and its metabolites are provided in SupplTable 1

2 References provided in Supplementary File

PM, Poor metabolizers; EM, Extensive metabolizers; SD, Single Dose; MD, Multiple dose; R, R-enantiomer; S, S-enantiomer; NR, Not reported.

* Asterisk denotes M/P ratio from single time point concentration

^a http://www.accessdata.fda.gov/drugsatfda_docs/nda/2002/21-411_Strattera_biopharmr_P2.pdf

Table 2A: Physicochemical properties, abundance and estimated fm of alcohol metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	pKa	cLogD	tPSA	M/P	Conjugate	fm	Ref
Amiodarone	5.87	42.7	3'-OH-Desethyl amiodarone	14.97	3.69	71.7	0.39			(McDonald et al., 2012)
Bosentan	2.36	154	Hydroxybosentan	14.63	0.65	174	0.11 to 0.13		0.3	(Weber et al., 1999; van Giersbergen et al., 2002)
Bromhexine	3.69	29.3	Hydroxybromhexine	15.11	2.38	49.5	E4-OH:0.42; E3-OH:0.17			(Liu et al., 2010)
Bufuralol	1.46	45.4	1'-Hydroxybufuralol	14.3	0.066	65.6	1			(Pringle et al., 1986)
Bupropion	2.13	29.1	Hydroxybupropion	15.43	1.28	49.3	7 to 41			(Kirchheiner et al., 2003a; Loboz et al., 2006)
			Dihydrobupropion	13.67	0.94	32.1	Threo: 6.4; Erthro:1.8			(Yeung et al., 2011)
Buspirone	1.24	69.6	Hydroxybuspirone	12.5	-0.4	89.9	34 to 44			(Jajoo et al., 1989; Dockens et al., 2006)
Celecoxib	2.59	83.4	Hydroxycelecoxib	14.16	0.93	107	0.05		0.73	(Paulson et al., 2000; Kirchheiner et al., 2003b)
Cerivastatin	0.071	99.9	Hydroxycerivastatin	14.4	-1.36	120	0.24 to 0.37			(Muck, 2000)

			O-Desmethyl cerivastatin	13.38	-0.7	111	0.08 to 0.11		
Clarithromycin	2.06	183	Hydroxyclearithromycin	14.1	0.99	203	0.67 (SD); 0.42 (MD)	0.2	(Ferrero et al., 1990; Cheng et al., 1998)
Dolansetron	2.54	62.4	Dehydrodolasetron	14.59	1.23	65.6	>1	0.32	(Reith et al., 1995; Lerman et al., 1996)
Darifenacin	1.95	55.6	Hydroxydarifenacin	13.52	0.86	75.6	1 to 1.3	0.32	(Beaumont et al., 1998; Skerjanec, 2006)
Erlotinib	3	74.7	Desmethylelotinib	13.9	2.41	85.7	0.05	0.04	(Frohna et al., 2006; Ling et al., 2006)
Flutamide	3.52	74.9	Hydroxyflutamide	12.87	2.67	95.2	73		(Anjum et al., 1999)
Indacaterol	2.33	61.4	Hydroxyindacaterol	14.58	0.39	81.6	0.39	0.2	(Kagan et al., 2012)
Haloperidol	2.93	40.5	Dihydrohaloperidol	14.29	1.56	43.7	0.88		(Chakraborty et al., 1989)
Itraconazole	4.93	102	Hydroxyitraconazole	14.98	3.46	121	1.4		(Templeton et al., 2008)
Ketanserine	3.05	69.7	Ketanserinol	14.06	2.07	72.9	3.2 to 3.5		(Lebrec et al., 1990)
Metoprolol	-0.47	50.7	α -Hydroxymetoprolol	13.74	-1.68	70.6	0.56		(Cerqueira et al., 2005)
			O-Desmethylmetoprolol	14.9	-1.09	61.7	ND		

Midazolam	3.78	30.2	1'-Hydroxymidazolam	13.59	2.5	50.4	0.48	Yes	0.65	(Heizmann and Ziegler, 1981; Eap et al., 2004)
Montelukast	3.2	95.7	Hydroxymontelukast (M1)	15.86	2.2	116	0.06			(Karonen et al., 2012)
			Hydroxymontelukast (M5a/5b)	13.98	1.82	116	0.77			
Nalidixic Acid	-1.54	70.5	Hydroxynalidixic acid	13.02	-2.3	90.7	0.69			(Cuisinaud et al., 1982)
Naltrexone	1.6	70	6- β -Naltrexol	14.42	0.24	73	29			(Yun et al., 2007)
Nebivolol	2.75	70.6	Hydroxyneбиволol	13.98	1.23	91.2	21			(Kamali et al., 1997)
Nefazadone	3.89	51.6	Hydroxynefazadone	13.8	3.55	71.9	0.35 to 0.55			(Barbhaiya et al., 1995; Barbhaiya et al., 1996a; Barbhaiya et al., 1996b)
Nelfinavir	7.26	127	Hydroxynelfinavir	16.11	6.1	147	0.37			(Zhang et al., 2001)
Nortryptiline	1.49	12	Hydroxynortryptiline	13.9	-0.25	32.3	0.36 -13			(Dalen et al., 1998)
Omeprazole	2.35	96.3	Hydroxyomeprazole	13.15	1.2	117	0.42 to 0.7		0.25	(Renberg et al., 1989; Andersson et al., 1990; Lutz and Isoherranen, 2012)
Pagoclone	3.31	65.2	Hydroxypagoclone	14.98	1.64	83.4	65% of total plasma radioactivity			(Dalvie et al., 2009)

Pioglitazone	2.45	93.6	Hydroxyioglitazone	13.74	0.89	114	SD: 3.8; MD:1.8		(Eckland and Danhof, 2000; Budde et al., 2003)
Praziquantel	2.66	40.6	Hydroxypraziquantel	15.02	1.02	60.9	(+)S:9.0; (-) R: 33-38		(Westhoff and Blaschke, 1992; Lima et al., 2009)
Quinidine	0.98	45.6	Hydroxyquinidine	13.93	0.88	65.8	0.26 to 0.29		(Rakhit et al., 1984; Schellens et al., 1991)
Repaglinide	1.99	78.9	Hydroxyrepaglinide	14.79	0.42	99.1	0.02*	0.01	(van Heiningen et al., 1999; Honkalammi et al., 2011)
Risperidone	1.89	61.9	Hydroxyrisperidone	13	0.61	82.2	3 to 6	0.25	(Mannens et al., 1993; Zhou et al., 2006; Mahatthanatrakul et al., 2012)
Ruzolitininib	2.07	83.2	Hydroxyruzolitininib	14.94	0.56	103	0.24 to 0.3	0.05	(Shilling et al., 2010; Shi et al., 2012)
Taranabant	6.29	75	Hydroxytaranabant	15.14	5.58	95.2	2.5 to 3.3		(Addy et al., 2008; Karanam et al., 2010)
Terfenadine	3.6	43.7	Hydroxyterfenadine	14.96	1.88	63.9	ND		(Abernethy et al., 2001)
Tibolone	3.52	37.3	3 α -Hydroxytibolone	15	3.96	40.5	0.44	0.16	(Vos et al., 2002)
Ticagrelor	2.02	164	O-Desalkylticagrelor	14.28	2.02	155	0.4	0.22	(Teng et al.,

										2010)
Tolterodine	2.37	23.5	Hydroxytolteridine	14.71	1.93	43.7	0.82			(Brynne et al., 1999)
Valdecoxib	3.56	94.6	Hydroxyvaldecoxib	13.09	2.36	115	0.08 to 0.1	Yes	0.23	(Yuan et al., 2002; Sarapa et al., 2005)

1. Structures of drugs and its metabolites are provided in SupplTable 2A

2. References provided in Supplementary File

* Asterisk denotes M/P ratio from single time point concentration

Table 2B: Physicochemical properties, abundance and estimated fm of ketone metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	cLogD	tPSA	M/P	fm	Ref
Nefazadone	3.89	51.6	triazolodione	1.07	68.4	2.1		(Barbhaiya et al., 1995; Barbhaiya et al., 1996a; Barbhaiya et al., 1996b)
Itraconazole	3.77	118	Ketoitraconazole	4.93	101	0.04		(Templeton et al., 2008)
Otenabant	4.71	102	Ketone Metabolite	4.55	63.9	1.1		(Miao et al., 2012)
Pioglitazone	2.45	93.6	Ketopioglitazone	0.73	111	0.9 to 2.8 (SD); 0.5 (MD)		(Eckland and Danhof, 2000; Budde et al., 2003)
Bufuralol	1.46	45.4	1'-Oxobufuralol	0.19	62.5	0.1 to 0.2		(Pringle et al., 1986)

1. Structures of drugs and its metabolites are provided in SupplTable 2B

2. References provided in Supplementary File

SD, Single Dose; MD, Multiple Dose

Table 3: Physicochemical properties, abundance and estimated fm of arenol metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	pKa	cLogD	tPSA	M/P (Aglycone)	Conjugate ³	fm	Ref
Amoxapine	1.79	36.9	7-Hydroxyamoxapine	11.16	1.68	57.1	0.22 to 0.44	Yes	0.25	(Jue et al., 1982; Kobayashi et al., 1985; Takeuchi et al., 1993)
			8-Hydroxyamoxapine	10.35	1.02	57.1	2.4 to 4.9	Yes	0.33	
Atomoxetine	0.77	21.3	Hydroxyatomoxetine	10.33	-0.13	41.5	CYP2D6 EM: 0.01	Yes (CYP2D6 EM- 2.5 & PM - 0.11)	CYP2D6 EM:0.87	(Sauer et al., 2003; Sauer et al., 2005); FDA Clinical Pharmacology Review ^a
Atorvastatin	0.74	112	<i>o</i> -Hydroxyatorvastatin	9.34	1	132	1.4	NR		(Whitfield et al., 2011)
			<i>p</i> -Hydroxyatorvastatin	10	0.1	132	0.04	NR		
Chlorpromazine	3.24	31.8	Hydroxychlorpromazine	9.19	2.65	52	0.58 to 1.8	Yes (2.2 to 4.2)		(Yeung et al., 1993)

Desipramine	1.21	15.3	Hydroxydesipramine	10	-0.19	35.5	0.22 to 0.79	Yes (3.3)		(DeVane et al., 1981; Spina et al., 1997; Patroneva et al., 2008)
Desloratadine	0.84	24.9	Hydroxydesloratadine	9.27	0.49	45.2	0.5	Yes	0.51	(Gupta et al., 2004; Ramanathan et al., 2007b)
Duloxetine	2.31	49.5	Hydroxyduloxetine	9.97	1.51	69.7		Yes (12)	0.22	(Lantz et al., 2003; Knadler et al., 2011)
			5-Hydroxy-6-methoxy duloxetine	7.67	1.49	78.9		Yes (5.6)	0.19	
Efavirenz	3.03	38.3	HydroxyEfavirenz	7.18	2.38	58.6	0.04	Yes (1.1)	0.24	(Mutlib et al., 1999; Cho et al., 2011); FDA Clinical Pharmacology Review ^b
Elzasonan	3.24	52.1	5-HydroxyElsazonan	11.5	3.16	72.3	0	Yes	0.34	(Kamel et al., 2010)
Granisetron	0.0166	50.2	Hydroxygranisetron	8.9	-0.59	70.4	0.11*	0.26*	0.25	(Clarke et al., 1994)
Imipramine	2.34	6.48	Hydroxyimipramine	10.2	1.71	26.7	0.37 to 0.89	Yes (1.9)		(Sutfin et al., 1984; Wells et al., 1986; Sutfin et al., 1988; Spina

										et al., 1997)
Lansoprazole	2.58	87.1	Hydroxylansoprazole	11.16	1.9	10.7	CYP2C19 EM:0.1 - 0.19; PM:0.02			(Miura et al., 2004; Zhang et al., 2011)
Loxapine	2.23	28.1	7-Hydroxyloxapine	11.16	2.1	48.3	0.68			(Simpson et al., 1978; Cheung et al., 1991)
			8-Hydroxyloxapine	10.35	1.45	48.3	3			
Nelfinavir	7.26	127	Methoxycathecol	8.57	7.28	136	0.05			(Zhang et al., 2011)
Propafenone	1.37	58.6	Hydroxypropafenone	9.48	1.21	78.8	0.23 - 0.5			(Kates et al., 1985; Dilger et al., 1999)
Propranolol	0.79	41.5	Hydroxypropranolol	10.17	0.25	61.7	0.01 - 0.09 (without hydrolysis) 2.8 (with hydrolysis)	Yes		(Raghuram et al., 1984; Partani et al., 2009)
Quetiapine	2.51	73.6	Hydroxyquetiapine	11.18	1.91	93.8	0.1			(Gefvert et al., 1998; DeVane and Nemeroff, 2001)
Rosiglitazone	1.93	96.8	Hydroxyrosiglitazone	10.3	1.33	117		Yes (2.8)	0.29	(Cox et al., 2000; Kirchheiner et

										al., 2006)
Tamoxifen	3.83	12.5	Hydroxytamoxifen	10.35	3.63	32.7	0.035			(Yeung et al., 2011)
S-Warfarin	0.33	63.6	Hydroxywarfarin	8.51	0.36	83.8	0.15			(Uno et al., 2008)
Amiodarone	5.87	42.7	O-Desalkylamiodarone	4.84	3.33	50.4	0.1	No		(McDonald et al., 2012)
Ampremilast	0.48	127	Demethoxyampremilast	9.8	-0.032	138	ND	Yes (1.24)	0.39	(Hoffmann et al., 2011)
Apixaban	4.66	111	Demethoxyapixaban	8.56	4.02	122	ND	Yes (0.25)	0.18	(Raghavan et al., 2009)
Astemizole	4.12	42.3	Demethoxyastemizole	10.13	3.53	53.3	6.8			(Lefebvre et al., 1997)
Bosentan	2.36	154	Demethoxybosentan	8.36	2.34	165	0.02 – 0.05		0.07	(Weber et al., 1999; van Giersbergen et al., 2002)
Codeine	0.46	41.9	Morphine	9.48	-0.043	52.9	0.003 – 0.008	Yes (M6G:0.33-0.45; M3G:2 - 2.6)		(Kirchheiner et al., 2007)
Dextromethorphan	2.17	12.5	Demethyl-dextromethorphan	10.07	1.71	23.5	CYP2D6 EM: 0.38 (without	Yes		(Bolden et al., 2002; Pope et

							hydrolysis)			al., 2004)
							69 (with hydrolysis)			
Encainide	1.62	41.57	O-Demethyencainide	8.34	1.2	52.6	CYP2D6 EM: 1.2			(Funk-Brentano et al., 1989)
Flecainide	1.05	59.6	Dealkylflecainide	9.28	-0.14	70.6	0.04	Yes	0.23	(Munafo et al., 1990)
Gefitinib	2.56	68.7	Demethylgefitinib	5.5	1.79	79.7	0.7		0.23	(McKillop et al., 2004); FDA Clinical Pharmacology Review ^c
Pantoprazole	1.54	106	Demethylpantoprazole	7.69	0.49	117	ND	Yes (0.2 – 0.5)		(Radhofer-Welte, 1999)
Paroxetine	1.48	39.7	4-Hydroxy-3-Methoxy Paroxetine	10.34	1.04	50.7	NR	Yes		(Segura et al., 2003)
Ranolazine	1.28	74.3	Demethylranolazine	9.59	0.94	85.3	0.36	Yes		(Abdallah and Jerling, 2005; Jerling and Abdallah, 2005; Jerling et al., 2005)
Tadalafil	2.71	74.9	4-Hydroxy-3-Methoxy Tadalafil	9.87	2.85	85.9	SD - 0.58 MD - 1.3	Yes		(Forgue et al., 2006; Fogue et al., 2007)

Tramadol	0.29	32.7	Demethyltramadol	10	-0.25	43.7	0.35 - 0.95	No		(Ardakani et al., 2008; de Moraes et al., 2012)
Venlafaxine	0.76	32.7	Desvenlafaxine	10	0.0015	43.7	EM - 4.0 - 6.0 PM - 0.35	Yes	0.56	(Howell et al., 1993; Patat et al., 1998; Hynninen et al., 2008)

1. Structures of drugs and its metabolites are provided in SupplTable 3
2. References provided in Supplementary File
3. Yes: denotes presence of conjugate of the metabolite in circulation; NR: Not reported; the number represents M/P ratio of the metabolite conjugate to parent

EM, Extensive metabolizers; PM, Poor Metabolizers; SD, Single Dose; MD, Multiple Dose; ND, Not Detected; * Asterik denotes M/P ratio from single time point concentration

^a http://www.accessdata.fda.gov/drugsatfda_docs/nda/2002/21-411_Strattera_biopharmr_P2.pdf

^b http://www.accessdata.fda.gov/drugsatfda_docs/nda/98/20972biopharm_review.pdf

^c http://www.accessdata.fda.gov/drugsatfda_docs/nda/2003/21-399_IRESSA_Clinr.pdf

Table 4: Physicochemical properties, abundance and estimated fm of N and S-oxide metabolites ^{1,2}.

Drug	cLogD	tPSA	Metabolite	cLogD	tPSA	M/P	fm	Ref
S- Oxidation								
Albendazole	2.99	92.3	Albendazole sulfoxide	0.68	103	>1		(Marriner et al., 1986; Delatour et al., 1991)
Axitinib	2.39	96	Axitinib sulfoxide	1.89	107	1		Product label
Chlorpromazine	3.24	31.78	Chlorpromazine sulfoxide	2.32	0.37	0.95 to 2.4		(Yeung et al., 1993)
Dothiepin	2.54	28.5	Dothiepin sulfoxide	0.98	39.52	2.4		(Maguire et al., 1981)
Flosequinan	-1.43	56.6	Flosequinan sulfone	-1.18	62.83	14		(Wynne et al., 1985)
Lansoprazole	2.58	87.1	Lansoprazole sulfone	3.55	93.3	CYP2C19 EM:0.05; PM:0.33 - 0.88		(Miura et al., 2004; Zhang et al., 2011)
Montelukast	3.2	95.7	Montelukast sulfoxide	1.74	107	NR		(Karonen et al., 2012)
Omeprazole	2.35	96.3	Omeprazole sulfone	3.31	103	0.6 – 0.89		(Renberg et al., 1989; Andersson et al., 1990; Lutz and Isoherranen, 2012)

Pantoprazole	1.54	106	Pantoprazole sulfone	2.46	112	Low (<0.25)		(Radhofer-Welte, 1999)
Quetiapine	2.51	73.6	Quetiapine sulfoxide	-0.76	84.6	0.65		(Gefvert et al., 1998; DeVane and Nemeroff, 2001)
Ranitidine	-2.47	117	Ranitidine sulfoxide	-3.43	128	0.09	0.01	(Prueksaritanont et al., 1989); Product label
Thioridazine	3.49	57.1	Thioridazine sulfoxide (Mesoridazine)	0.76	68.1	4.3 – 6.5		(Hartigan-Go et al., 1996)
			Thioridazine sulfoxide- 2	1	68.1	3.9 – 4.2		
			Mesoridazine sulfone	2.15	74.3	0.95 to 1.6		
N-Oxidation								
Azimilide	2.66	72.6	Azimilide N-oxide	2.22	86.43	0.01	0.07	(Riley et al., 2005)
Chlorpromazine	3.24	31.8	Chlorpromazine N-oxide	2.39	45.6	0.8 – 1.8		(Yeung et al., 1993)
Dothiepin	2.54	28.5	Dothiepin N-oxide	1.61	42.4	Not known		(Maguire et al., 1981)
Elzasonan	3.24	52.1	Elzasonan N-oxide	2.47	65.9	0.27		(Kamel et al., 2010)

DMD #50278

Quinidine	0.98	45.6	Quinidine N-oxide	1.26	59.4	0.14 – 0.39		(Rakhit et al., 1984; Schellens et al., 1991)
Ranitidine	-2.47	117	Ranitidine N-oxide	-4.56	131	0.19	0.04	(Prueksaritanont et al., 1989); Product label
Regorafenib	4.2	92.4	Regorafenib N-oxide	1.55	105	0.96 – 1.1		(Mross et al., 2012; Strumberg et al., 2012)
			N-Desmethylregorafenib N-oxide	1.64	119	1.3 – 1.8		
Roflumilast	2.3	60.5	Roflumilast N-oxide	0.21	73	13	0.21	(Lahu et al., 2008); FDA Clinical Pharmacology Review ^a
Voriconazole	1.21	77	Voriconazole N-oxide	0.45	89.3	1.6	0.21	(Roffey et al., 2003; Damle et al., 2008)
Zolmitriptan	-1.47	57	Zolmitriptan N-oxide	-2.43	71.2	0.38 – 0.48	0.07	(Dixon et al., 1997; Seaber et al., 1997)
Zopiclone	2.63	91.8	Zopiclone N-oxide	1.24	106	0.18	0.14	(Mistri et al., 2008)

1. Structures of drugs and its metabolites are provided in SupplTable 4
2. References provided in Supplementary File

^a http://www.accessdata.fda.gov/drugsatfda_docs/nda/2011/022522Orig1s000ClinPharmR.pdf

Table 5: Physicochemical properties, abundance and estimated fm of carboxylic acid metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	cLogD	tPSA	M/P	fm	Ref
Azimilide	2.66	72.6	Carboxylic acid	1.05	50.4	4	0.35	(Riley et al., 2005)
Celecoxib	2.59	83.4	Carboxylic acid Metabolite	-1.77	124	1	0.73	(Paulson et al., 2000; Kirchheiner et al., 2003a)
Losartan	1.48	92.5	Carboxylosartan	-0.06	110	4.4	0.14	(Lo et al., 1995; Neves et al., 2008)
Metoprolol	-0.47	50.7	O-Desmethylnetoprolol Carboxylic acid	-1.65	78.8	6.5		(Cerqueira et al., 2005)
Montelukast	3.2	95.7	Montelukast carboxylic acid	0.21	133	0.24		(Karonen et al., 2012)
Nabumetone	3.14	26.3	6-Methoxynaphthyl carboxylic acid	-0.45	46.5	>1		(Nobilis et al., 2003)
Quetiapine	2.51	73.6	Quetiapine carboxylic acid	-1.13	90.7	0.63		(Gefvert et al., 1998; DeVane and Nemeroff, 2001)
Ranolazine	1.28	74.3	Cleaved Carboxylic acid	-3.12	76	0.25		(Abdallah and Jerling, 2005; Jerling and Abdallah, 2005; Jerling et al., 2005)
Regaplinide	1.99	78.9	Carboxylic Acid Metabolite	-1.86	125	0.05	0.66	(van Heiningen et al., 1999; Honkalammi et al., 2011)
Terfenadine	3.6	43.7	Fexofenadine	1.23	81	98		(Abernethy et al., 2001)
Zolmitriptan	-1.47	57.4	Zolmitriptan indoleacetic acid	-3.4	91.4	1.3	0.31	(Dixon et al., 1997; Seaber et al., 1997)

1. Structures of drugs and its metabolites are provided in SupplTable 5
2. References provided in Supplementary File

Table 6: Physicochemical properties, abundance and estimated fm of novel metabolites^{1,2}.

Drug	cLogD	tPSA	Metabolite	cLogD	tPSA	M/P	fm	Ref
Aripiprazole	3.54	44.8	Dehydroaripiprazole	3.32	44.8	0.5	0.38	(Caccia, 2011); FDA Clinical Pharmacology Review ^a
Carbamazepine	1.89	46.3	Epoxy carbamazepine	0.16	58.9	0.25		(Ragueneau-Majlessi et al., 2004)
Elzasonan	3.24	52.1	Indole Derivative of Elzasonan	3.39	53.8	3.2	0.06	(Kamel et al., 2010)
Indapamide	1.96	101	Dehydroindapamide	2.65	103	Major		(Yeung et al., 2011)
Methadone	2.29	20.3	EDDP Metabolite	-3.51	3.01	0.16		(Kharasch et al., 2009)
Otenabant	4.71	102	Otenabant oxime	4.64	79.4	1.5		(Miao et al., 2012)
Proguanil	0.77	88.8	Cycloguanil	-0.69	80	0.67		(Funck-Brentano et al., 1997)

1 Structures of drugs and its metabolites are provided in SupplTable 6

2 References provided in Supplementary File

^ahttp://www.accessdata.fda.gov/drugsatfda_docs/nda/2002/21-436_Abilify_biopharmr_P2.pdf

Table 7: Drug and metabolite free fraction and calculated total and free drug clearances for selected drugs shown in Tables 1-6.

Drug	Metabolite	Fraction unbound	CLs ¹ (mL/min/kg)	CL _{R, U} ² (mL/min/kg)	CL _{NR, U} ³ (mL/min/kg)	Ref
Clozapine		0.055	5.4	0.1	98	(Schaber et al., 1998)
	desmethyl-clozapine	0.097	0.6	5	1.2	
Chlorpromazine		0.0013	18.3	1	14000	(Freedberg et al., 1979)
	hydroxy-chlorpromazine	0.015	6.1	6	400	(Yeung et al., 1993)
Darifenacin		0.02	12	6	594	(Kerbusch et al., 2004)
	Hydroxyl-darifenacin	0.13	14	47	60	
Diltiazem		0.25	12		48	(Boyd et al., 1989)
	N-desmethyl diltiazem	0.32	8		25	
Propranolol		0.15	20	0.7	132	(Weiss et al., 1978)
	hydroxy-propranolol	0.26	66	5	248	(Rochester et al., 1980)
Tolterodine		0.037	44	1	1190	(Pahlman and Gozzi, 1999)

DMD #50278

	N-dealkyl-tolterodine	0.14				
	hydroxyl-tolterodine	0.36	21	2.9	55	
Thioridazine		0.0016				(Freedberg et al., 1979)
	thioridazine sulfoxide (Mesoridazine)	0.090				

1. CL_s = Total systemic clearance
2. $CL_{R,U}$ = Unbound renal clearance
3. $CL_{NR,U}$ = Unbound non-renal (hepato-biliary) clearance

Figure 1




Figure 2







Figure 3




Nefazadone
cLogD = 3.89




m-Chlorophenylpiperazine
cLogD = 0.56




R = CH(CH₃)₂, Delavirdine
R = H, N-Desisopropyldelavirdine




R & R' = CH₃, Sibutramine
R = H & R' = CH₃, N-Desmethylsibutramine
R & R' = H, Didesmethylsibutramine




R & R' = CH₃, Tamoxifen
R = H & R' = CH₃, N-Desmethyltamoxifen
R & R' = H, Didesmethyltamoxifen



Imipramine




Desipramine




Hydroxydesipramine


Figure 4




R = CH₃, Erlotinib
R = H, O-Desmethylerlotinib




R = CH₂CH₃, Itraconazole
R = HOCHCH₃, Hydroxyitraconazole
R = O=CCH₃, Keto itraconazole




R = CH₃, Midazolam
R = CH₂OH, Hydroxymethylmidazolam
R = CH₂Oglucuronide, Glucuronide conjugate
of Hydroxy metabolite




R = CH₃, Valdecoxib
R = CH₂OH, Hydroxymethylvaldecoxib
R = CH₂Oglucuronide, Glucuronide conjugate
of Hydroxy metabolite



R = CH₃, Terfenadine
R = CO₂H, Fexofenadine




R = CH₃, Bosentan
R = OH, Hydroxybosentan




Bupropion

R, R, and S, S- Hydroxybupropion


Figure 5




R = H Atorvastatin
R = OH *ortho*-Hydroxyatorvastatin




R = CH₃, Ranolazine
R = H, *ortho*-Hydroxy metabolite




R = CH₃, Gefitinib
R = H, *ortho*-Hydroxy metabolite



R = CH₃, Tramadol
R = H, *meta*-Hydroxy metabolite



R = CH₃, Bosentan
R = H, Hydroxy metabolite



R = CH₃, Venlafaxine
R = H, Desvenlafaxine




Figure 6





Figure 7




**Drug Metabolism and Disposition
Which Metabolites Circulate?**

Cho-Ming Loi, Dennis A. Smith and Deepak Dalvie


Pharmacokinetics, Dynamics, and Metabolism, Pfizer Worldwide Research, San Diego, CA 92121 (*C-M L and DD*) and Honorary Professor, Department of Chemistry, University of Capetown, Capetown, South Africa, Visiting Professor, Institute of Translational Medicine, University of Liverpool, UK (*DAS*).

SupplTable 1: Structures of drugs and their amine metabolites


Drugs	Metabolites	
Amiodarone	N-Desethylamiodarone	
Amitriptyline	Didesethylamiodarone	
Nortriptyline	Desmethyatomoxetine	
Atomoxetine	Azelastine	
Desmethylazelastine	Azimilide	
Azimilide	Desmethylazimilide	
Buspirone	1-Pyrimidinylpiperazine	




Citalopram




Desmethylcitalopram




Didesmethylcitalopram




Clobazam




Norclobazam




Clomipramine




Desmethylclomipramine




Clozapine




Desmethylclozapine




Delavirdine




Desisopropyl delavirdine




Diazepam




Desmethyldiazepam




Diltiazem




Desmethyldiltiazem




Dothiepin




Desmethyldothiepin




Doxepin




Desmethyldoxepin




Enzalutamide




Desmethylenzalutamide




Elzasonan




Desmethyl elzasonan




Fenfluramine




Desethylfenfluramine




Fentanyl




Norfentanyl




Fluoxetine




Norfluoxetine




Halofantrine




Desbutylhalofantrine




Imatinib




Desmethylimatinib




Imipramine




Desipramine




Itraconazole




Dealkylitraconazole




Ivabradine




Desmethylivabradine




Loratadine




Desloratadine




Loxapine




Desmethylloxapine




Lumefantrine




Desbutyllumefantrine




Mianserin




Desmethylmianserin




Mirtazapine




Desmethylmirtazapine




Nefazodone




m-Chlorophenylpiperazine




Olanzapine




Desmethylolanzapine




Otenabant (CP-945598)




Desethylotenabant




Oxybutynin




Desethyloxybutynin




Propafenone




Despropylpropafenone




Ranitidine




Desmethylranitidine




Ranolazine




Desalkylranolazine




Regalinide




Dealkylregaplinide (M1)




Rosiglitazone




Desmethylrosiglitazone




Sertraline




Desmethylsertraline




Sibutramine




Desmethylsibutramine




Didesmethylsibutramine




Sildenafil




Desmethylsildenafil




Cleaved piperazine ring




Sunitinib




Desethylsunitinib




Tamoxifen




Desmethyltamoxifen




Didesmethyltamoxifen




Terbinafine




Desmethylterbinafine




Tramadol




Desmethyltramadol




Udenafil




Desalkyludenafil




Vardenafil




Desethylvardenafil




Verapamil




Norverapamil




D617




D620




Zimelidine




Desmethylzimelidine




Zolmitriptan



Desmethylzolmitriptan


















Zopiclone




Desmethylzopiclone

SupplTable 2A: Structures of drugs and their alcohol metabolites


Compounds	Metabolite
	
	
	
	
	
	
	




Flutamide




Hydroxyflutamide




Haloperidol




Dihydrohaloperidol




Indacaterol




Hydroxyindacaterol




Itraconazole




Hydroxyitraconazole




Ketanserine




Ketanserinol




Metoprolol




α -Hydroxymetoprolol
















O-Desmethylnmetoprolol




Midazolam




1'-Hydroxymidazolam







Nortryptiline




Hydroxynortryptiline




Omeprazole




Hydroxyomeprazole




Pagoclone




Hydroxypagoclone




Pioglitazone




Hydroxypioglitazone




Praziquantel




Hydroxypraziquantel




Quinidine




Hydroxyquinidine




Repaglinide

















Hydroxyrepaglinide













Risperidone














Hydroxyrisperidone

















SupplTable 2B: Structures of drugs and their ketone metabolites

Drugs	Metabolites
	
Nefazadone	triazolodione
	
Itraconazole	Ketoitraconazole
	
Otenabant	Ketone Metabolite
	
Pioglitazone	Ketopioglitazone
	
Bufuralol	1'-Oxobufuralol


SupplTable 3: Structures of drugs and their arenol metabolites

Compounds	Metabolite
 <p>Amiodarone</p>	 <p>O-Desalkylamiodarone</p>
 <p>Amoxapine</p>	 <p>7-Hydroxyamoxapine</p>
 <p>8-Hydroxyamoxapine</p>	
 <p>Apixaban</p>	 <p>Demethoxyapixaban</p>
 <p>Apremilast</p>	 <p>Demethoxyapremilast</p>
 <p>Astemizole</p>	 <p>Demethoxyastemizole</p>







Desloratadine




Hydroxydesloratadine




Dextromethorphan




Demethoxydextromethorphan




Duloxetine




Hydroxyduloxetine




5-Hydroxy-6-methoxy duloxetine




Efavirenz




HydroxyEfavirenz




Elzasonan




5-HydroxyElzasonan




Encainide




Demethyencanide




Flecainide




Dealkyloxyflecainide




Gefitinib




Demethoxygefitinib




Granisetron




Hydroxygranisetron




Imipramine




Hydroxyimipramine




Lansoprazole




Hydroxylansoprazole




Loxapine




7-Hydroxyloxapine




8-Hydroxyloxapine




Nelfinavir




Methoxycathecol




Paroxetine




4-Hydroxy-3-Methoxy Paroxetine




Pantoprazole




Demethoxypantoprazole




Propafenone




Hydroxypropafenone




Propranolol




Hydroxypropranolol




Quetiapine




Hydroxyquetiapine




Ranolazine




Demethoxyranolazine




Rosiglitazone




Hydroxyrosiglitazone




Tamoxifen




Hydroxytamoxifen




Tadalafil




4-Hydroxy-3-Methoxy Tadalafil




Tramadol




Demethoxytramadol




Venlafaxine



Desvenlafaxine
















S-Warfarin




Hydroxywarfarin

SupplTable 4: Structures of drugs and their S and N-oxide metabolites


Compounds	Metabolite
S-Oxides	
	
Albendazole	Albendazole sulfoxide
	
Axitinib	Axitinib sulfoxide
	
Chlorpromazine	Chlorpromazine sulfoxide
	
Dothiepin	Dothiepin sulfoxide
	
Montelukast	Montelukast sulfoxide
	
Flosequinan	Flosequinan sulfone




Lansoprazole




Lansoprazole sulfone




Omeprazole




Omeprazole sulfone




Pantoprazole




Pantoprazole sulfone




Quetiapine




Quetiapine sulfoxide




Ranitidine




Ranitidine sulfoxide




Thioridazine

















Thioridazine sulfoxide (Mesoridazine)




Thioridazine sulfoxide-2




N-Oxides







N-Desmethylregorafinib N-oxide




Roflumilast N-oxide




Roflumilast




Voriconazole




Voriconazole N-oxide




Zolmitriptan



Zolmitriptan N-oxide


















Zopiclone




Zopiclone N-oxide

SupplTable 5: Structures of drugs and their carboxylic acid metabolites


Drugs	Metabolites
	
Azimilide	Carboxylic acid
	
Celecoxib	Carboxylic acid Metabolite
	
Losartan	Carboxylosartan
	
Metoprolol	O-Desmethylmetoprolol Carboxylic acid
	
Montelukast	Montelukast carboxylic acid
	
Nabumetone	6-Methoxynaphthyl carboxylic acid
	
Quetiapine	Quetiapine carboxylic acid




Ranolazine




Cleaved Carboxylic acid




Regalinide




Carboxylic Acid Metabolite




Terfenadine



Fexofenadine

















Zolmitriptan



Zolmitriptan indoleacetic acid

SupplTable 6: Structures of drugs and their novel metabolites

Drugs	Metabolites
	
Aripiprazole	Dehydroaripiprazole
	
Carbamazepine	Epoxycarbamazepine
	
Elzasonan	Indole Derivative of Elzasonan
	
Indapamide	Dehydroindapamide
	
Methadone	EDDP Metabolite
	
Otenabant	Otenabant oxime
	
Proguanil	Cycloguanil

References for Citations in Tables 1 to 6:

Table References:

- Abdallah H and Jerling M (2005) Effect of hepatic impairment on the multiple-dose pharmacokinetics of ranolazine sustained-release tablets. *J Clin Pharmacol* **45**:802-809.
- Abernethy DR, Barbey JT, Franc J, Brown KS, Feirrer I, Ford N, and Salazar DE (2001) Loratadine and terfenadine interaction with nefazodone: Both antihistamines are associated with QTc prolongation. *Clin Pharmacol Ther* **69**:96-103.
- Abernethy DR, Wainer IW, and Anacleto AI (2000) Verapamil metabolite exposure in older and younger men during steady-state oral verapamil administration. *Drug Metab Dispos* **28**:760-765.
- Addy C, Rothenberg P, Li S, Majumdar A, Agrawal N, Li H, Zhong L, Yuan J, Maes A, Dunbar S, Cote J, Rosko K, Van Dyck K, De Lepeleire I, de Hoon J, Van Hecken A, Depre M, Knops A, Gottesdiener K, Stoch A, and Wagner J (2008) Multiple-dose pharmacokinetics, pharmacodynamics, and safety of taranabant, a novel selective cannabinoid-1 receptor inverse agonist, in healthy male volunteers. *J Clin Pharmacol* **48**:734-744.
- Andersson T, Cederberg C, Regardh CG, and Skanberg I (1990) Pharmacokinetics of various single intravenous and oral doses of omeprazole. *Eur J Clin Pharmacol* **39**:195-197.
- Anjum S, Swan SK, Lambrecht LJ, Radwanski E, Cutler DL, Affrime MB, and Halstenson CE (1999) Pharmacokinetics of flutamide in patients with renal insufficiency. *Br J Clin Pharmacol* **47**:43-47.
- Ardakani YH, Mehvar R, Foroumadi A, and Rouini MR (2008) Enantioselective determination of tramadol and its main phase I metabolites in human plasma by high-performance liquid chromatography. *J Chromatogr B Analyt Technol Biomed Life Sci* **864**:109-115.
- Bae JW, Jang CG, and Lee SY (2011) Effects of clopidogrel on the pharmacokinetics of sibutramine and its active metabolites. *J Clin Pharmacol* **51**:1704-1711.
- Barbhaiya RH, Brady ME, Shukla UA, and Greene DS (1995) Steady-state pharmacokinetics of nefazodone in subjects with normal and impaired renal function. *Eur J Clin Pharmacol* **49**:229-235.
- Barbhaiya RH, Dandekar KA, and Greene DS (1996a) Pharmacokinetics, absolute bioavailability, and disposition of [¹⁴C]nefazodone in humans. *Drug Metab Dispos* **24**:91-95.
- Barbhaiya RH, Shukla UA, Greene DS, Breul HP, and Midha KK (1996b) Investigation of pharmacokinetic and pharmacodynamic interactions after coadministration of nefazodone and haloperidol. *J Clin Psychopharmacol* **16**:26-34.
- Beaumont KC, Cussans NJ, Nichols DJ, and Smith DA (1998) Pharmacokinetics and metabolism of darifenacin in the mouse, rat, dog and man. *Xenobiotica* **28**:63-75.
- Bolden RD, Hoke SH, 2nd, Eichhold TH, McCauley-Myers DL, and Wehmeyer KR (2002) Semi-automated liquid-liquid back-extraction in a 96-well format to decrease sample preparation time for the determination of dextromethorphan and dextrorphan in human plasma. *J Chromatogr B Analyt Technol Biomed Life Sci* **772**:1-10.

- Borin MT, Chambers JH, Carel BJ, Gagnon S, and Freimuth WW (1997a) Pharmacokinetic study of the interaction between rifampin and delavirdine mesylate. *Clin Pharmacol Ther* **61**:544-553.
- Borin MT, Cox SR, Herman BD, Carel BJ, Anderson RD, and Freimuth WW (1997b) Effect of fluconazole on the steady-state pharmacokinetics of delavirdine in human immunodeficiency virus-positive patients. *Antimicrob Agents Chemother* **41**:1892-1897.
- Brannan MD, Reidenberg P, Radwanski E, Shneyer L, Lin CC, Cayen MN, and Affrime MB (1995) Loratadine administered concomitantly with erythromycin: pharmacokinetic and electrocardiographic evaluations. *Clin Pharmacol Ther* **58**:269-278.
- Brown D, Scott DH, Scott DB, Meyer M, Westerlund D, and Lundstrom J (1980) Pharmacokinetics of zimelidine. Systemic availability of zimelidine and norzimelidine in human volunteers. *Eur J Clin Pharmacol* **17**:111-116.
- Brynne N, Forslund C, Hallen B, Gustafsson LL, and Bertilsson L (1999) Ketoconazole inhibits the metabolism of tolterodine in subjects with deficient CYP2D6 activity. *Br J Clin Pharmacol* **48**:564-572.
- Budde K, Neumayer HH, Fritsche L, Sulowicz W, Stompor T, and Eckland D (2003) The pharmacokinetics of pioglitazone in patients with impaired renal function. *Br J Clin Pharmacol* **55**:368-374.
- Burch JE and Hullin RP (1981) Amitriptyline pharmacokinetics. A crossover study with single doses of amitriptyline and nortriptyline. *Psychopharmacology (Berl)* **74**:35-42.
- Caccia S (2011) Pharmacokinetics and metabolism update for some recent antipsychotics. *Expert Opin Drug Metab Toxicol* **7**:829-846.
- Caccia S, Conforti I, Duchier J, and Garattini S (1985) Pharmacokinetics of fenfluramine and norfenfluramine in volunteers given D- and DL-fenfluramine for 15 days. *Eur J Clin Pharmacol* **29**:221-224.
- Cerqueira PM, Coelho EB, Geleilate TJ, Goldman GH, and Lanchote VL (2005) Influence of chronic renal failure on stereoselective metoprolol metabolism in hypertensive patients. *J Clin Pharmacol* **45**:1422-1433.
- Chakraborty BS, Hubbard JW, Hawes EM, McKay G, Cooper JK, Gurnsey T, Korchinski ED, and Midha KK (1989) Interconversion between haloperidol and reduced haloperidol in healthy volunteers. *Eur J Clin Pharmacol* **37**:45-48.
- Cheng KL, Nafziger AN, Peloquin CA, and Amsden GW (1998) Effect of grapefruit juice on clarithromycin pharmacokinetics. *Antimicrob Agents Chemother* **42**:927-929.
- Cheung SW, Tang SW, and Remington G (1991) Simultaneous quantitation of loxapine, amoxapine and their 7- and 8-hydroxy metabolites in plasma by high-performance liquid chromatography. *J Chromatogr* **564**:213-221.
- Cho DY, Ogburn ET, Jones D, and Desta Z (2011) Contribution of N-glucuronidation to efavirenz elimination in vivo in the basal and rifampin-induced metabolism of efavirenz. *Antimicrob Agents Chemother* **55**:1504-1509.
- Chung MC, Lin SK, Chang WH, and Jann MW (1993) Determination of clozapine and desmethylclozapine in human plasma by high-performance liquid chromatography with ultraviolet detection. *J Chromatogr* **613**:168-173.
- Clarke SE, Austin NE, Bloomer JC, Haddock RE, Higham FC, Hollis FJ, Nash M, Shardlow PC, Tasker TC, Woods FR, and et al. (1994) Metabolism and disposition of ¹⁴C-granisetron in rat, dog and man after intravenous and oral dosing. *Xenobiotica* **24**:1119-1131.

- Cox PJ, Ryan DA, Hollis FJ, Harris AM, Miller AK, Vousden M, and Cowley H (2000) Absorption, disposition, and metabolism of rosiglitazone, a potent thiazolidinedione insulin sensitizer, in humans. *Drug Metab Dispos* **28**:772-780.
- Cuisinaud G, Ferry N, Pozet N, Zech PY, and Sassard J (1982) Nalidixic acid kinetics in renal insufficiency. *Br J Clin Pharmacol* **14**:489-493.
- Dalen P, Dahl ML, Bernal Ruiz ML, Nordin J, and Bertilsson L (1998) 10-Hydroxylation of nortriptyline in white persons with 0, 1, 2, 3, and 13 functional CYP2D6 genes. *Clin Pharmacol Ther* **63**:444-452.
- Dalvie D, Obach RS, Kang P, Prakash C, Loi CM, Hurst S, Nedderman A, Goulet L, Smith E, Bu HZ, and Smith DA (2009) Assessment of three human in vitro systems in the generation of major human excretory and circulating metabolites. *Chem Res Toxicol* **22**:357-368.
- Damle B, LaBadie R, Crownover P, and Glue P (2008) Pharmacokinetic interactions of efavirenz and voriconazole in healthy volunteers. *Br J Clin Pharmacol* **65**:523-530.
- de Moraes NV, Lauretti GR, Napolitano MN, Santos NR, Godoy AL, and Lanchote VL (2012) Enantioselective analysis of unbound tramadol, O-desmethyltramadol and N-desmethyltramadol in plasma by ultrafiltration and LC-MS/MS: application to clinical pharmacokinetics. *J Chromatogr B Analyt Technol Biomed Life Sci* **880**:140-147.
- Delatour P, Benoit E, Besse S, and Boukraa A (1991) Comparative enantioselectivity in the sulphoxidation of albendazole in man, dogs and rats. *Xenobiotica* **21**:217-221.
- DeVane CL and Nemeroff CB (2001) Clinical pharmacokinetics of quetiapine: an atypical antipsychotic. *Clin Pharmacokinet* **40**:509-522.
- DeVane CL, Savett M, and Jusko WJ (1981) Desipramine and 2-hydroxy-desipramine pharmacokinetics in normal volunteers. *Eur J Clin Pharmacol* **19**:61-64.
- Dilger K, Greiner B, Fromm MF, Hofmann U, Kroemer HK, and Eichelbaum M (1999) Consequences of rifampicin treatment on propafenone disposition in extensive and poor metabolizers of CYP2D6. *Pharmacogenetics* **9**:551-559.
- Dixon R, Gillotin C, Gibbens M, Posner J, and Peck RW (1997) The pharmacokinetics and effects on blood pressure of multiple doses of the novel anti-migraine drug zolmitriptan (311C90) in healthy volunteers. *Br J Clin Pharmacol* **43**:273-281.
- Dockens RC, Salazar DE, Fulmor IE, Wehling M, Arnold ME, and Croop R (2006) Pharmacokinetics of a newly identified active metabolite of buspirone after administration of buspirone over its therapeutic dose range. *J Clin Pharmacol* **46**:1308-1312.
- Eap CB, Buclin T, Cucchia G, Zullino D, Hustert E, Bleiber G, Golay KP, Aubert AC, Baumann P, Telenti A, and Kerb R (2004) Oral administration of a low dose of midazolam (75 microg) as an in vivo probe for CYP3A activity. *Eur J Clin Pharmacol* **60**:237-246.
- Eckland DA and Danhof M (2000) Clinical Pharmacokinetics of pioglitazone. *Exp Clin Endocrinol Diabetes* **108**:S234-S-242.
- Fernandez C, Martin C, Gimenez F, and Farinotti R (1995) Clinical pharmacokinetics of zopiclone. *Clin Pharmacokinet* **29**:431-441.
- Ferrero JL, Bopp BA, Marsh KC, Quigley SC, Johnson MJ, Anderson DJ, Lamm JE, Tolman KG, Sanders SW, Cavanaugh JH, and et al. (1990) Metabolism and disposition of clarithromycin in man. *Drug Metab Dispos* **18**:441-446.

- Fjordside L, Jeppesen U, Eap CB, Powell K, Baumann P, and Brosen K (1999) The stereoselective metabolism of fluoxetine in poor and extensive metabolizers of sparteine. *Pharmacogenetics* **9**:55-60.
- Forgue ST, Patterson BE, Bedding AW, Payne CD, Phillips DL, Wrishko RE, and Mitchell MI (2006) Tadalafil pharmacokinetics in healthy subjects. *Br J Clin Pharmacol* **61**:280-288.
- Forgue ST, Phillips DL, Bedding AW, Payne CD, Jewell H, Patterson BE, Wrishko RE, and Mitchell MI (2007) Effects of gender, age, diabetes mellitus and renal and hepatic impairment on tadalafil pharmacokinetics. *Br J Clin Pharmacol* **63**:24-35.
- Frohna P, Lu J, Eppler S, Hamilton M, Wolf J, Rakhit A, Ling J, Kenkare-Mitra SR, and Lum BL (2006) Evaluation of the absolute oral bioavailability and bioequivalence of erlotinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in a randomized, crossover study in healthy subjects. *J Clin Pharmacol* **46**:282-290.
- Funck-Brentano C, Becquemont L, Leneveu A, Roux A, Jaillon P, and Beaune P (1997) Inhibition by omeprazole of proguanil metabolism: mechanism of the interaction in vitro and prediction of in vivo results from the in vitro experiments. *J Pharmacol Exp Ther* **280**:730-738.
- Funck-Brentano C, Turgeon J, Woosley RL, and Roden DM (1989) Effect of low dose quinidine on encainide pharmacokinetics and pharmacodynamics. Influence of genetic polymorphism. *J Pharmacol Exp Ther* **249**:134-142.
- Gefvert O, Bergstrom M, Langstrom B, Lundberg T, Lindstrom L, and Yates R (1998) Time course of central nervous dopamine-D2 and 5-HT2 receptor blockade and plasma drug concentrations after discontinuation of quetiapine (Seroquel) in patients with schizophrenia. *Psychopharmacology (Berl)* **135**:119-126.
- Gschwind HP, Pfaar U, Waldmeier F, Zollinger M, Sayer C, Zbinden P, Hayes M, Pokorny R, Seiberling M, Ben-Am M, Peng B, and Gross G (2005) Metabolism and disposition of imatinib mesylate in healthy volunteers. *Drug Metab Dispos* **33**:1503-1512.
- Gupta S, Banfield C, Kantesaria B, Flannery B, and Herron J (2004) Pharmacokinetics/pharmacodynamics of desloratadine and fluoxetine in healthy volunteers. *J Clin Pharmacol* **44**:1252-1259.
- Hamelin BA, Turgeon J, Vallee F, Belanger PM, Paquet F, and LeBel M (1996) The disposition of fluoxetine but not sertraline is altered in poor metabolizers of debrisoquin. *Clin Pharmacol Ther* **60**:512-521.
- Hartigan-Go K, Bateman DN, Nyberg G, Martensson E, and Thomas SH (1996) Concentration-related pharmacodynamic effects of thioridazine and its metabolites in humans. *Clin Pharmacol Ther* **60**:543-553.
- Heizmann P and Ziegler WH (1981) Excretion and metabolism of ¹⁴C-midazolam in humans following oral dosing. *Arzneimittelforschung* **31**:2220-2223.
- Herrlin K, Yasui-Furukori N, Tybring G, Widen J, Gustafsson LL, and Bertilsson L (2003) Metabolism of citalopram enantiomers in CYP2C19/CYP2D6 phenotyped panels of healthy Swedes. *Br J Clin Pharmacol* **56**:415-421.
- Hoffmann M, Kumar G, Schafer P, Cedzik D, Capone L, Fong KL, Gu Z, Heller D, Feng H, Surapaneni S, Laskin O, and Wu A (2011) Disposition, metabolism and mass balance of [¹⁴C]apremilast following oral administration. *Xenobiotica* **41**:1063-1075.
- Honkalammi J, Niemi M, Neuvonen PJ, and Backman JT (2011) Dose-dependent interaction between gemfibrozil and repaglinide in humans: strong inhibition of CYP2C8 with subtherapeutic gemfibrozil doses. *Drug Metab Dispos* **39**:1977-1986.

- Howell SR, Husbands GE, Scatina JA, and Sisenwine SF (1993) Metabolic disposition of ¹⁴C-venlafaxine in mouse, rat, dog, rhesus monkey and man. *Xenobiotica* **23**:349-359.
- Hynninen VV, Olkkola KT, Bertilsson L, Kurkinen K, Neuvonen PJ, and Laine K (2008) Effect of terbinafine and voriconazole on the pharmacokinetics of the antidepressant venlafaxine. *Clin Pharmacol Ther* **83**:342-348.
- Jajoo HK, Mayol RF, LaBudde JA, and Blair IA (1989) Metabolism of the antianxiety drug buspirone in human subjects. *Drug Metab Dispos* **17**:634-640.
- Jerling M and Abdallah H (2005) Effect of renal impairment on multiple-dose pharmacokinetics of extended-release ranolazine. *Clin Pharmacol Ther* **78**:288-297.
- Jerling M, Huan BL, Leung K, Chu N, Abdallah H, and Hussein Z (2005) Studies to investigate the pharmacokinetic interactions between ranolazine and ketoconazole, diltiazem, or simvastatin during combined administration in healthy subjects. *J Clin Pharmacol* **45**:422-433.
- Jiang ZP, Shu Y, Chen XP, Huang SL, Zhu RH, Wang W, He N, and Zhou HH (2002) The role of CYP2C19 in amitriptyline N-demethylation in Chinese subjects. *Eur J Clin Pharmacol* **58**:109-113.
- Jue SG, Dawson GW, and Brogden RN (1982) Amoxapine: a review of its pharmacology and efficacy in depressed states. *Drugs* **24**:1-23.
- Kagan M, Dain J, Peng L, and Reynolds C (2012) Metabolism and pharmacokinetics of indacaterol in humans. *Drug Metab Dispos* **40**:1712-1722.
- Kamali F, Howes A, Thomas SH, Ford GA, and Snoeck E (1997) A pharmacokinetic and pharmacodynamic interaction study between nebivolol and the H₂-receptor antagonists cimetidine and ranitidine. *Br J Clin Pharmacol* **43**:201-204.
- Kamel A, Obach RS, Colizza K, Wang W, O'Connell TN, Coelho RV, Jr., Kelley RM, and Schildknecht K (2010) Metabolism, pharmacokinetics, and excretion of the 5-hydroxytryptamine_{1b} receptor antagonist elzasonan in humans. *Drug Metab Dispos* **38**:1984-1999.
- Karanam B, Addy C, Bateman T, Reddy VB, Li S, Dean D, Li H, Jones A, Schenk D, Zhang AS, Braun M, Freeman A, Flach S, Stoch A, Chodakewitz J, Wagner JA, and Kumar S (2010) Metabolism and excretion of [¹⁴C]taranabant, a cannabinoid-1 inverse agonist, in humans. *Xenobiotica* **40**:691-700.
- Karonen T, Neuvonen PJ, and Backman JT (2012) CYP2C8 but not CYP3A4 is important in the pharmacokinetics of montelukast. *Br J Clin Pharmacol* **73**:257-267.
- Kassahun K, Mattiuz E, Nyhart E, Jr., Obermeyer B, Gillespie T, Murphy A, Goodwin RM, Tupper D, Callaghan JT, and Lemberger L (1997) Disposition and biotransformation of the antipsychotic agent olanzapine in humans. *Drug Metab Dispos* **25**:81-93.
- Kates RE, Yee YG, and Winkle RA (1985) Metabolite cumulation during chronic propafenone dosing in arrhythmia. *Clin Pharmacol Ther* **37**:610-614.
- Kharasch ED, Hoffer C, Whittington D, Walker A, and Bedynek PS (2009) Methadone pharmacokinetics are independent of cytochrome P4503A (CYP3A) activity and gastrointestinal drug transport: insights from methadone interactions with ritonavir/indinavir. *Anesthesiology* **110**:660-672.
- Kim KA, Song WK, and Park JY (2009) Association of CYP2B6, CYP3A5, and CYP2C19 genetic polymorphisms with sibutramine pharmacokinetics in healthy Korean subjects. *Clin Pharmacol Ther* **86**:511-518.

- Kirchheiner J, Henckel HB, Franke L, Meineke I, Tzvetkov M, Uebelhack R, Roots I, and Brockmoller J (2005) Impact of the CYP2D6 ultra-rapid metabolizer genotype on doxepin pharmacokinetics and serotonin in platelets. *Pharmacogenet Genomics* **15**:579-587.
- Kirchheiner J, Klein C, Meineke I, Sasse J, Zanger UM, Murdter TE, Roots I, and Brockmoller J (2003a) Bupropion and 4-OH-bupropion pharmacokinetics in relation to genetic polymorphisms in CYP2B6. *Pharmacogenetics* **13**:619-626.
- Kirchheiner J, Schmidt H, Tzvetkov M, Keulen JT, Lotsch J, Roots I, and Brockmoller J (2007) Pharmacokinetics of codeine and its metabolite morphine in ultra-rapid metabolizers due to CYP2D6 duplication. *Pharmacogenomics J* **7**:257-265.
- Kirchheiner J, Stormer E, Meisel C, Steinbach N, Roots I, and Brockmoller J (2003b) Influence of CYP2C9 genetic polymorphisms on pharmacokinetics of celecoxib and its metabolites. *Pharmacogenetics* **13**:473-480.
- Kirchheiner J, Thomas S, Bauer S, Tomalik-Scharte D, Hering U, Doroshenko O, Jetter A, Stehle S, Tsahuridu M, Meineke I, Brockmoller J, and Fuhr U (2006) Pharmacokinetics and pharmacodynamics of rosiglitazone in relation to CYP2C8 genotype. *Clin Pharmacol Ther* **80**:657-667.
- Knadler MP, Lobo E, Chappell J, and Bergstrom R (2011) Duloxetine: clinical pharmacokinetics and drug interactions. *Clin Pharmacokinet* **50**:281-294.
- Kobayashi A, Sugita S, and Nakazawa K (1985) Determination of amoxapine and its metabolites in human serum by high-performance liquid chromatography. *Neuropharmacology* **24**:1253-1256.
- Kosuge K, Jun Y, Watanabe H, Kimura M, Nishimoto M, Ishizaki T, and Ohashi K (2001) Effects of CYP3A4 inhibition by diltiazem on pharmacokinetics and dynamics of diazepam in relation to CYP2C19 genotype status. *Drug Metab Dispos* **29**:1284-1289.
- Ku HY, Shon JH, Liu KH, Shin JG, and Bae SK (2009) Liquid chromatography/tandem mass spectrometry method for the simultaneous determination of vardenafil and its major metabolite, N-desethylvardenafil, in human plasma: application to a pharmacokinetic study. *J Chromatogr B Analyt Technol Biomed Life Sci* **877**:95-100.
- Lahu G, Huennemeyer A, von Richter O, Hermann R, Herzog R, McCracken N, and Zech K (2008) Effect of single and repeated doses of ketoconazole on the pharmacokinetics of roflumilast and roflumilast N-oxide. *J Clin Pharmacol* **48**:1339-1349.
- Lantz RJ, Gillespie TA, Rash TJ, Kuo F, Skinner M, Kuan HY, and Knadler MP (2003) Metabolism, excretion, and pharmacokinetics of duloxetine in healthy human subjects. *Drug Metab Dispos* **31**:1142-1150.
- Lebrec D, Hadengue A, Gaudin C, Levron JC, Fraitag B, Berthelot P, and Benhamou JP (1990) Pharmacokinetics of ketanserin in patients with cirrhosis. *Clin Pharmacokinet* **19**:160-166.
- Lefebvre RA, Van Peer A, and Woestenborghs R (1997) Influence of itraconazole on the pharmacokinetics and electrocardiographic effects of astemizole. *Br J Clin Pharmacol* **43**:319-322.
- Lerman J, Sims C, Sikich N, Gow R, Chin C, Dempsey E, Howard DR, and Keung AC (1996) Pharmacokinetics of the active metabolite (MDL 74,156) of dolasetron mesylate after oral or intravenous administration to anesthetized children. *Clin Pharmacol Ther* **60**:485-492.

- Levy RH, Lane EA, Guyot M, Brachet-Liermain A, Cenraud B, and Loiseau P (1983) Analysis of parent drug-metabolite relationship in the presence of an inducer. Application to the carbamazepine-clobazam interaction in normal man. *Drug Metab Dispos* **11**:286-292.
- Lima RM, Ferreira MA, Ponte TM, Marques MP, Takayanagui OM, Garcia HH, Coelho EB, Bonato PS, and Lanchote VL (2009) Enantioselective analysis of praziquantel and trans-4-hydroxypraziquantel in human plasma by chiral LC-MS/MS: Application to pharmacokinetics. *J Chromatogr B Analyt Technol Biomed Life Sci* **877**:3083-3088.
- Ling J, Johnson KA, Miao Z, Rakhit A, Pantze MP, Hamilton M, Lum BL, and Prakash C (2006) Metabolism and excretion of erlotinib, a small molecule inhibitor of epidermal growth factor receptor tyrosine kinase, in healthy male volunteers. *Drug Metab Dispos* **34**:420-426.
- Liu J, Chen X, Hu Y, Cheng G, and Zhong D (2010) Quantification of the major metabolites of bromhexine in human plasma using RRLC-MS/MS and its application to pharmacokinetics. *J Pharm Biomed Anal* **51**:1134-1141.
- Lo MW, Goldberg MR, McCrea JB, Lu H, Furtek CI, and Bjornsson TD (1995) Pharmacokinetics of losartan, an angiotensin II receptor antagonist, and its active metabolite EXP3174 in humans. *Clin Pharmacol Ther* **58**:641-649.
- Loboz KK, Gross AS, Williams KM, Liauw WS, Day RO, Bliedernicht JK, Zanger UM, and McLachlan AJ (2006) Cytochrome P450 2B6 activity as measured by bupropion hydroxylation: effect of induction by rifampin and ethnicity. *Clin Pharmacol Ther* **80**:75-84.
- Lutz JD and Isoherranen N (2012) Prediction of relative in vivo metabolite exposure from in vitro data using two model drugs: dextromethorphan and omeprazole. *Drug Metab Dispos* **40**:159-168.
- Maguire KP, Burrows GD, Norman TR, and Scoggins BA (1981) Metabolism and pharmacokinetics of dothiepin. *Br J Clin Pharmacol* **12**:405-409.
- Mahatthanatrakul W, Sriwiriyan S, Ridditid W, Boonleang J, Wongnawa M, Rujimamahasan N, and Pipatrattanaseree W (2012) Effect of cytochrome P450 3A4 inhibitor ketoconazole on risperidone pharmacokinetics in healthy volunteers. *J Clin Pharm Ther* **37**:221-225.
- Mannens G, Huang ML, Meuldermans W, Hendrickx J, Woestenborghs R, and Heykants J (1993) Absorption, metabolism, and excretion of risperidone in humans. *Drug Metab Dispos* **21**:1134-1141.
- Marriner SE, Morris DL, Dickson B, and Bogan JA (1986) Pharmacokinetics of albendazole in man. *Eur J Clin Pharmacol* **30**:705-708.
- McDonald MG, Au NT, Wittkowsky AK, and Rettie AE (2012) Warfarin-amiodarone drug-drug interactions: determination of [I](u)/K(I,u) for amiodarone and its plasma metabolites. *Clin Pharmacol Ther* **91**:709-717.
- McGready R, Stepniewska K, Lindegardh N, Ashley EA, La Y, Singhasivanon P, White NJ, and Nosten F (2006) The pharmacokinetics of artemether and lumefantrine in pregnant women with uncomplicated falciparum malaria. *Eur J Clin Pharmacol* **62**:1021-1031.
- McKillop D, Hutchison M, Partridge EA, Bushby N, Cooper CM, Clarkson-Jones JA, Herron W, and Swaisland HC (2004) Metabolic disposition of gefitinib, an epidermal growth factor receptor tyrosine kinase inhibitor, in rat, dog and man. *Xenobiotica* **34**:917-934.
- Miao Z, Sun H, Liras J, and Prakash C (2012) Excretion, metabolism, and pharmacokinetics of 1-(8-(2-chlorophenyl)-9-(4-chlorophenyl)-9H-purin-6-yl)-4-(ethylamino)piperidine-4-

- carboxamide, a selective cannabinoid receptor antagonist, in healthy male volunteers. *Drug Metab Dispos* **40**:568-578.
- Mistri HN, Jangid AG, Pudage A, and Shrivastav P (2008) HPLC-ESI-MS/MS validated method for simultaneous quantification of zopiclone and its metabolites, N-desmethyl zopiclone and zopiclone-N-oxide in human plasma. *J Chromatogr B Analyt Technol Biomed Life Sci* **864**:137-148.
- Miura M, Tada H, Yasui-Furukori N, Uno T, Sugawara K, Tateishi T, and Suzuki T (2004) Pharmacokinetic differences between the enantiomers of lansoprazole and its metabolite, 5-hydroxylansoprazole, in relation to CYP2C19 genotypes. *Eur J Clin Pharmacol* **60**:623-628.
- Moraes MO, Lerner FE, Corso G, Bezerra FA, Moraes ME, and De Nucci G (1999) Fluoxetine bioequivalence study: quantification of fluoxetine and norfluoxetine by liquid chromatography coupled to mass spectrometry. *J Clin Pharmacol* **39**:1053-1061.
- Morse GD, Fischl MA, Shelton MJ, Cox SR, Driver M, DeRemer M, and Freimuth WW (1997) Single-dose pharmacokinetics of delavirdine mesylate and didanosine in patients with human immunodeficiency virus infection. *Antimicrob Agents Chemother* **41**:169-174.
- Mross K, Frost A, Steinbild S, Hedbom S, Buchert M, Fasol U, Unger C, Kratzschmar J, Heinig R, Boix O, and Christensen O (2012) A phase I dose-escalation study of regorafenib (BAY 73-4506), an inhibitor of oncogenic, angiogenic, and stromal kinases, in patients with advanced solid tumors. *Clin Cancer Res* **18**:2658-2667.
- Muck W (2000) Clinical pharmacokinetics of cerivastatin. *Clin Pharmacokinet* **39**:99-116.
- Muirhead GJ, Rance DJ, Walker DK, and Wastall P (2002) Comparative human pharmacokinetics and metabolism of single-dose oral and intravenous sildenafil. *Br J Clin Pharmacol* **53 Suppl 1**:13S-20S.
- Munafò A, Raymond-Michel G, and Biollaz J (1990) Altered flecainide disposition in healthy volunteers taking quinine. *Eur J Clin Pharmacol* **38**:269-273.
- Mutlib AE, Chen H, Nemeth GA, Markwalder JA, Seitz SP, Gan LS, and Christ DD (1999) Identification and characterization of efavirenz metabolites by liquid chromatography/mass spectrometry and high field NMR: species differences in the metabolism of efavirenz. *Drug Metab Dispos* **27**:1319-1333.
- Nagy A and Johansson R (1977) The demethylation of imipramine and clomipramine as apparent from their plasma kinetics. *Psychopharmacology (Berl)* **54**:125-131.
- Neves R, Almeida S, Filipe A, Spinola AC, Abolfathi Z, Yritia M, and Ortuno J (2008) Bioequivalence study of two different film-coated tablet formulations of losartan-hydrochlorothiazide in healthy volunteers. *Arzneimittelforschung* **58**:369-375.
- Nobilis M, Kopecky J, Kvetina J, Svoboda Z, Pour M, Kunes J, Holcapek M, and Kolarova L (2003) Comparative biotransformation and disposition studies of nabumetone in humans and minipigs using high-performance liquid chromatography with ultraviolet, fluorescence and mass spectrometric detection. *J Pharm Biomed Anal* **32**:641-656.
- Ouellet D, Hsu A, Qian J, Lamm JE, Cavanaugh JH, Leonard JM, and Granneman GR (1998) Effect of fluoxetine on pharmacokinetics of ritonavir. *Antimicrob Agents Chemother* **42**:3107-3112.
- Partani P, Modhave Y, Gurule S, Khuroo A, and Monif T (2009) Simultaneous determination of propranolol and 4-hydroxy propranolol in human plasma by solid phase extraction and liquid chromatography/electrospray tandem mass spectrometry. *J Pharm Biomed Anal* **50**:966-976.

- Patat A, Troy S, Burke J, Trocherie S, Danjou P, Le Coz F, Allain H, and Gandon JM (1998) Absolute bioavailability and electroencephalographic effects of conventional and extended-release formulations of venlafaxine in healthy subjects. *J Clin Pharmacol* **38**:256-267.
- Patroneva A, Connolly SM, Fatato P, Pedersen R, Jiang Q, Paul J, Guico-Pabia C, Isler JA, Burczynski ME, and Nichols AI (2008) An assessment of drug-drug interactions: the effect of desvenlafaxine and duloxetine on the pharmacokinetics of the CYP2D6 probe desipramine in healthy subjects. *Drug Metab Dispos* **36**:2484-2491.
- Paulson SK, Hribar JD, Liu NW, Hajdu E, Bible RH, Jr., Piergies A, and Karim A (2000) Metabolism and excretion of [(14)C]celecoxib in healthy male volunteers. *Drug Metab Dispos* **28**:308-314.
- Perucca E, Gatti G, Cipolla G, Spina E, Barel S, Soback S, Gips M, and Bialer M (1994) Inhibition of diazepam metabolism by fluvoxamine: a pharmacokinetic study in normal volunteers. *Clin Pharmacol Ther* **56**:471-476.
- Pinder RM and van Delft AM (1983a) Pharmacological aspects of mianserin. *Acta Psychiatr Scand Suppl* **302**:59-71.
- Pinder RM and Van Delft AM (1983b) The potential therapeutic role of the enantiomers and metabolites of mianserin. *Br J Clin Pharmacol* **15 Suppl 2**:269S-276S.
- Pope LE, Khalil MH, Berg JE, Stiles M, Yakatan GJ, and Sellers EM (2004) Pharmacokinetics of dextromethorphan after single or multiple dosing in combination with quinidine in extensive and poor metabolizers. *J Clin Pharmacol* **44**:1132-1142.
- Portoles A, Calvo A, Terleira A, Laredo L, Resplandy G, Gorostiaga C, and Moreno A (2006a) Lack of pharmacokinetic interaction between omeprazole or lansoprazole and ivabradine in healthy volunteers: an open-label, randomized, crossover, pharmacokinetic interaction clinical trial. *J Clin Pharmacol* **46**:1195-1203.
- Portoles A, Terleira A, Calvo A, Martinez I, and Resplandy G (2006b) Effects of Hypericum perforatum on ivabradine pharmacokinetics in healthy volunteers: an open-label, pharmacokinetic interaction clinical trial. *J Clin Pharmacol* **46**:1188-1194.
- Pringle TH, Francis RJ, East PB, and Shanks RG (1986) Pharmacodynamic and pharmacokinetic studies on bufuralol in man. *Br J Clin Pharmacol* **22**:527-534.
- Prueksaritanont T, Sittichai N, Prueksaritanont S, and Vongsaroj R (1989) Simultaneous determination of ranitidine and its metabolites in human plasma and urine by high-performance liquid chromatography. *J Chromatogr* **490**:175-185.
- Radhofer-Welte S (1999) Pharmacokinetics and metabolism of the proton pump inhibitor pantoprazole in man. *Drugs of Today* **35**:765-771.
- Raghavan N, Frost CE, Yu Z, He K, Zhang H, Humphreys WG, Pinto D, Chen S, Bonacorsi S, Wong PC, and Zhang D (2009) Apixaban metabolism and pharmacokinetics after oral administration to humans. *Drug Metab Dispos* **37**:74-81.
- Raghuram TC, Koshakji RP, Wilkinson GR, and Wood AJ (1984) Polymorphic ability to metabolize propranolol alters 4-hydroxypropranolol levels but not beta blockade. *Clin Pharmacol Ther* **36**:51-56.
- Ragueneau-Majlessi I, Levy RH, Bergen D, Garnett W, Rosenfeld W, Mather G, Shah J, and Grundy JS (2004) Carbamazepine pharmacokinetics are not affected by zonisamide: in vitro mechanistic study and in vivo clinical study in epileptic patients. *Epilepsy Res* **62**:1-11.

- Rakhit A, Holford NH, Guentert TW, Maloney K, and Riegelman S (1984) Pharmacokinetics of quinidine and three of its metabolites in man. *J Pharmacokinetic Biopharm* **12**:1-21.
- Ramanathan R, Reyderman L, Kulmatycki K, Su AD, Alvarez N, Chowdhury SK, Alton KB, Wirth MA, Clement RP, Statkevich P, and Patrick JE (2007a) Disposition of loratadine in healthy volunteers. *Xenobiotica* **37**:753-769.
- Ramanathan R, Reyderman L, Su AD, Alvarez N, Chowdhury SK, Alton KB, Wirth MA, Clement RP, Statkevich P, and Patrick JE (2007b) Disposition of desloratadine in healthy volunteers. *Xenobiotica* **37**:770-787.
- Reith MK, Sproles GD, and Cheng LK (1995) Human metabolism of dolasetron mesylate, a 5-HT₃ receptor antagonist. *Drug Metab Dispos* **23**:806-812.
- Renberg L, Simonsson R, and Hoffmann KJ (1989) Identification of two main urinary metabolites of [¹⁴C]omeprazole in humans. *Drug Metab Dispos* **17**:69-76.
- Riley P, Figary PC, Entwisle JR, Roe AL, Thompson GA, Ohashi R, Ohashi N, and Moorehead TJ (2005) The metabolic profile of azimilide in man: in vivo and in vitro evaluations. *J Pharm Sci* **94**:2084-2095.
- Robbins B, Chang CT, Cramer JA, Garreffa S, Hafkin B, Hunt TL, and Meligeni J (1996) Safe coadministration of terbinafine and terfenadine: a placebo-controlled crossover study of pharmacokinetic and pharmacodynamic interactions in healthy volunteers. *Clin Pharmacol Ther* **59**:275-283.
- Roffey SJ, Cole S, Comby P, Gibson D, Jezequel SG, Nedderman AN, Smith DA, Walker DK, and Wood N (2003) The disposition of voriconazole in mouse, rat, rabbit, guinea pig, dog, and human. *Drug Metab Dispos* **31**:731-741.
- Saari TI, Laine K, Bertilsson L, Neuvonen PJ, and Olkkola KT (2007) Voriconazole and fluconazole increase the exposure to oral diazepam. *Eur J Clin Pharmacol* **63**:941-949.
- Sarapa N, Britto MR, Mainka MB, and Parivar K (2005) The effect of mild and moderate hepatic impairment on the pharmacokinetics of valdecoxib, a selective COX-2 inhibitor. *Eur J Clin Pharmacol* **61**:247-256.
- Sauer JM, Ponsler GD, Mattiuz EL, Long AJ, Witcher JW, Thomasson HR, and Desante KA (2003) Disposition and metabolic fate of atomoxetine hydrochloride: the role of CYP2D6 in human disposition and metabolism. *Drug Metab Dispos* **31**:98-107.
- Sauer JM, Ring BJ, and Witcher JW (2005) Clinical pharmacokinetics of atomoxetine. *Clin Pharmacokinetic* **44**:571-590.
- Schellens JH, Ghabrial H, van der Wart HH, Bakker EN, Wilkinson GR, and Breimer DD (1991) Differential effects of quinidine on the disposition of nifedipine, sparteine, and mephenytoin in humans. *Clin Pharmacol Ther* **50**:520-528.
- Seaber E, On N, Dixon RM, Gibbens M, Leavens WJ, Liptrot J, Chittick G, Posner J, Rolan PE, and Pack RW (1997) The absolute bioavailability and metabolic disposition of the novel antimigraine compound zolmitriptan (311C90). *Br J Clin Pharmacol* **43**:579-587.
- Segura M, Ortuno J, Farre M, Pacifici R, Pichini S, Joglar J, Segura J, and de la Torre R (2003) Quantitative determination of paroxetine and its 4-hydroxy-3-methoxy metabolite in plasma by high-performance liquid chromatography/electrospray ion trap mass spectrometry: application to pharmacokinetic studies. *Rapid Commun Mass Spectrom* **17**:1455-1461.
- Shi JG, Chen X, Emm T, Scherle PA, McGee RF, Lo Y, Landman RR, McKeever EG, Jr., Punwani NG, Williams WV, and Yeleswaram S (2012) The effect of CYP3A4 inhibition or induction on the pharmacokinetics and pharmacodynamics of orally administered

- ruxolitinib (INCB018424 phosphate) in healthy volunteers. *J Clin Pharmacol* **52**:809-818.
- Shilling AD, Nedza FM, Emm T, Diamond S, McKeever E, Punwani N, Williams W, Arvanitis A, Galya LG, Li M, Shepard S, Rodgers J, Yue TY, and Yeleswaram S (2010) Metabolism, excretion, and pharmacokinetics of [¹⁴C]INCB018424, a selective Janus tyrosine kinase 1/2 inhibitor, in humans. *Drug Metab Dispos* **38**:2023-2031.
- Shin KH, Chung YJ, Kim BH, Kim TE, Kim HS, Cho JY, Jang IJ, Shin SG, and Yu KS (2010) Effect of ketoconazole on the pharmacokinetics of udenafil in healthy Korean subjects. *Br J Clin Pharmacol* **69**:307-310.
- Simpson GM, Cooper TB, Lee JH, and Young MA (1978) Clinical and plasma level characteristics of intramuscular and oral loxapine. *Psychopharmacology (Berl)* **56**:225-232.
- Skerjanec A (2006) The clinical pharmacokinetics of darifenacin. *Clin Pharmacokinet* **45**:325-350.
- Speed B, Bu HZ, Pool WF, Peng GW, Wu EY, Patyna S, Bello C, and Kang P (2012) Pharmacokinetics, distribution, and metabolism of [¹⁴C]sunitinib in rats, monkeys, and humans. *Drug Metab Dispos* **40**:539-555.
- Spina E, Avenoso A, Campo GM, Scordo MG, Caputi AP, and Perucca E (1997) Effect of ketoconazole on the pharmacokinetics of imipramine and desipramine in healthy subjects. *Br J Clin Pharmacol* **43**:315-318.
- Strumberg D, Scheulen ME, Schultheis B, Richly H, Frost A, Buchert M, Christensen O, Jeffers M, Heinig R, Boix O, and Mross K (2012) Regorafenib (BAY 73-4506) in advanced colorectal cancer: a phase I study. *Br J Cancer* **106**:1722-1727.
- Sugihara J, Sugawara Y, Ando H, Harigaya S, Etoh A, and Kohno K (1984) Studies on the metabolism of diltiazem in man. *J Pharmacobiodyn* **7**:24-32.
- Sutfin TA, DeVane CL, and Jusko WJ (1984) The analysis and disposition of imipramine and its active metabolites in man. *Psychopharmacology (Berl)* **82**:310-317.
- Sutfin TA, Perini GI, Molnar G, and Jusko WJ (1988) Multiple-dose pharmacokinetics of imipramine and its major active and conjugated metabolites in depressed patients. *J Clin Psychopharmacol* **8**:48-53.
- Takeuchi H, Yokota S, Shimada S, Ohtani Y, Miura S, and Kubo H (1993) Pharmacokinetics of Amoxapine and its active metabolites in healthy subjects. *Curr Ther Res* **53**:427-434.
- Templeton IE, Thummel KE, Kharasch ED, Kunze KL, Hoffer C, Nelson WL, and Isoherranen N (2008) Contribution of itraconazole metabolites to inhibition of CYP3A4 in vivo. *Clin Pharmacol Ther* **83**:77-85.
- Teng R, Oliver S, Hayes MA, and Butler K (2010) Absorption, distribution, metabolism, and excretion of ticagrelor in healthy subjects. *Drug Metab Dispos* **38**:1514-1521.
- Timmer CJ, Sitsen JM, and Delbressine LP (2000) Clinical pharmacokinetics of mirtazapine. *Clin Pharmacokinet* **38**:461-474.
- Uno T, Sugimoto K, Sugawara K, and Tateishi T (2008) The effect of CYP2C19 genotypes on the pharmacokinetics of warfarin enantiomers. *J Clin Pharm Ther* **33**:67-73.
- van Giersbergen PL, Halabi A, and Dingemans J (2002) Single- and multiple-dose pharmacokinetics of bosentan and its interaction with ketoconazole. *Br J Clin Pharmacol* **53**:589-595.
- van Heiningen PN, Hatorp V, Kramer Nielsen K, Hansen KT, van Lier JJ, De Merbel NC, Oosterhuis B, and Jonkman JH (1999) Absorption, metabolism and excretion of a single

- oral dose of (14)C-repaglinide during repaglinide multiple dosing. *Eur J Clin Pharmacol* **55**:521-525.
- Venkatakrishnan K, Schmider J, Harmatz JS, Ehrenberg BL, von Moltke LL, Graf JA, Mertzanis P, Corbett KE, Rodriguez MC, Shader RI, and Greenblatt DJ (2001) Relative contribution of CYP3A to amitriptyline clearance in humans: in vitro and in vivo studies. *J Clin Pharmacol* **41**:1043-1054.
- Vos RM, Krebbers SF, Verhoeven CH, and Delbressine LP (2002) The in vivo human metabolism of tibolone. *Drug Metab Dispos* **30**:106-112.
- Weber C, Gasser R, and Hopfgartner G (1999) Absorption, excretion, and metabolism of the endothelin receptor antagonist bosentan in healthy male subjects. *Drug Metab Dispos* **27**:810-815.
- Wells BG, Pieper JA, Self TH, Stewart CF, Waldon SL, Bobo L, and Warner C (1986) The effect of ranitidine and cimetidine on imipramine disposition. *Eur J Clin Pharmacol* **31**:285-290.
- Westhoff F and Blaschke G (1992) High-performance liquid chromatographic determination of the stereoselective biotransformation of the chiral drug praziquantel. *J Chromatogr* **578**:265-271.
- Whitfield LR, Porcari AR, Alvey C, Abel R, Bullen W, and Hartman D (2011) Effect of gemfibrozil and fenofibrate on the pharmacokinetics of atorvastatin. *J Clin Pharmacol* **51**:378-388.
- Wynne RD, Crampton EL, and Hind ID (1985) The pharmacokinetics and haemodynamics of BTS 49 465 and its major metabolite in healthy volunteers. *Eur J Clin Pharmacol* **28**:659-664.
- Yeung CK, Fujioka Y, Hachad H, Levy RH, and Isoherranen N (2011) Are circulating metabolites important in drug-drug interactions?: Quantitative analysis of risk prediction and inhibitory potency. *Clin Pharmacol Ther* **89**:105-113.
- Yeung PK, Hubbard JW, Korchinski ED, and Midha KK (1993) Pharmacokinetics of chlorpromazine and key metabolites. *Eur J Clin Pharmacol* **45**:563-569.
- Yeung PK, Mosher SJ, Quilliam MA, and Montague TJ (1990) Species comparison of pharmacokinetics and metabolism of diltiazem in humans, dogs, rabbits, and rats. *Drug Metab Dispos* **18**:1055-1059.
- Yu BN, Chen GL, He N, Ouyang DS, Chen XP, Liu ZQ, and Zhou HH (2003) Pharmacokinetics of citalopram in relation to genetic polymorphism of CYP2C19. *Drug Metab Dispos* **31**:1255-1259.
- Yuan JJ, Yang DC, Zhang JY, Bible R, Jr., Karim A, and Findlay JW (2002) Disposition of a specific cyclooxygenase-2 inhibitor, valdecoxib, in human. *Drug Metab Dispos* **30**:1013-1021.
- Yun HY, Bang SC, Lee KC, Baek IH, Lee SP, Kang W, and Kwon KI (2007) Simultaneous analysis of naltrexone and its major metabolite, 6-beta-naltrexol, in human plasma using liquid chromatography-tandem mass spectrometry: Application to a parent-metabolite kinetic model in humans. *Talanta* **71**:1553-1559.
- Zhang D, Wang X, Yang M, Wang G, and Liu H (2011) Effects of CYP2C19 polymorphism on the pharmacokinetics of lansoprazole and its main metabolites in healthy Chinese subjects. *Xenobiotica* **41**:511-517.
- Zhang KE, Wu E, Patick AK, Kerr B, Zorbas M, Lankford A, Kobayashi T, Maeda Y, Shetty B, and Webber S (2001) Circulating metabolites of the human immunodeficiency virus

- protease inhibitor nelfinavir in humans: structural identification, levels in plasma, and antiviral activities. *Antimicrob Agents Chemother* **45**:1086-1093.
- Zhou ZL, Li X, Peng HY, Yu XY, Yang M, Su FL, Wang F, Zhu RH, Deng CY, Lin QX, Wang CY, Li WB, Lin SG, and Li HD (2006) Multiple dose pharmacokinetics of risperidone and 9-hydroxyrisperidone in Chinese female patients with schizophrenia. *Acta Pharmacol Sin* **27**:381-386.
- Zobrist RH, Schmid B, Feick A, Quan D, and Sanders SW (2001) Pharmacokinetics of the R- and S-enantiomers of oxybutynin and N-desethyloxybutynin following oral and transdermal administration of the racemate in healthy volunteers. *Pharm Res* **18**:1029-1034.